

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 November 2001 (01.11.2001)

PCT

(10) International Publication Number
WO 01/81379 A2

- (51) International Patent Classification⁷: **C07K 14/195**
- (74) Agents: **POTTER, Jane, E., R.**; Seed Intellectual Property Law Group PLLC, Suite 6300, 701 Fifth Avenue, Seattle, WA 98104-7092 et al. (US).
- (21) International Application Number: PCT/US01/13081
- (22) International Filing Date: 23 April 2001 (23.04.2001)
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/198,853 21 April 2000 (21.04.2000) US
60/219,752 20 July 2000 (20.07.2000) US
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **CORIXA CORPORATION** [US/US]; 1124 Columbia Street, Suite 200, Seattle, WA 98104 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **BHATIA, Ajay** [IN/US]; 1805 Bellevue Avenue #204, Seattle, WA 98104 (US). **PROBST, Peter** [DE/US]; 137 N.W. 77th St., Seattle, WA 98117 (US). **STROMBERG, Erika, Jean** [US/US]; 440 N.E. 73rd Street #201, Seattle, WA 98103 (US).
- Published:**
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/81379 A2

(54) Title: COMPOUNDS AND METHODS FOR TREATMENT AND DIAGNOSIS OF CHLAMYDIAL INFECTION

(57) Abstract: Compounds and methods for the diagnosis and treatment of Chlamydial infection are disclosed. The compounds provided include polypeptides that contain at least one antigenic portion of a *Chlamydia* antigen and DNA sequences encoding such polypeptides. Pharmaceutical compositions and vaccines comprising such polypeptides or DNA sequences are also provided, together with antibodies directed against such polypeptides. Diagnostic kits containing such polypeptides or DNA sequences and a suitable detection reagent may be used for the detection of Chlamydial infection in patients and in biological samples.

COMPOUNDS AND METHODS FOR TREATMENT AND DIAGNOSIS OF CHLAMYDIAL INFECTION

TECHNICAL FIELD

5 The present invention relates generally to the detection and treatment of Chlamydial infection. In particular, the invention is related to polypeptides comprising a *Chlamydia* antigen and the use of such polypeptides for the serodiagnosis and treatment of Chlamydial infection.

10 BACKGROUND OF THE INVENTION

Chlamydiae are intracellular bacterial pathogens that are responsible for a wide variety of important human and animal infections. *Chlamydia trachomatis* is one of the most common causes of sexually transmitted diseases and can lead to pelvic inflammatory disease (PID), resulting in tubal obstruction and infertility. *Chlamydia*
15 *trachomatis* may also play a role in male infertility. In 1990, the cost of treating PID in the US was estimated to be \$4 billion. Trachoma, due to ocular infection with *Chlamydia trachomatis*, is the leading cause of preventable blindness worldwide. *Chlamydia pneumonia* is a major cause of acute respiratory tract infections in humans and is also believed to play a role in the pathogenesis of atherosclerosis and, in
20 particular, coronary heart disease. Individuals with a high titer of antibodies to *Chlamydia pneumonia* have been shown to be at least twice as likely to suffer from coronary heart disease as seronegative individuals. Chlamydial infections thus constitute a significant health problem both in the US and worldwide.

Chlamydial infection is often asymptomatic. For example, by the time a woman
25 seeks medical attention for PID, irreversible damage may have already occurred resulting in infertility. There thus remains a need in the art for improved vaccines and pharmaceutical compositions for the prevention and treatment of *Chlamydia* infections. The present invention fulfills this need and further provides other related advantages.

30 SUMMARY OF THE INVENTION

The present invention provides compositions and methods for the diagnosis and therapy of *Chlamydia* infection. In one aspect, the present invention provides polypeptides comprising an immunogenic portion of a *Chlamydia* antigen, or a

variant of such an antigen. Certain portions and other variants are immunogenic, such that the ability of the variant to react with antigen-specific antisera is not substantially diminished. Within certain embodiments, the polypeptide comprises an amino acid sequence encoded by a polynucleotide sequence selected from the group consisting of

5 (a) a sequence of SEQ ID NO: 1-48, 114-121, and 125-138; (b) the complements of said sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions. In specific embodiments, the polypeptides of the present invention comprise at least a portion of a *Chlamydial* protein that includes an amino acid sequence selected from the group consisting of sequences recited in SEQ ID NO:122-

10 124 and 139-140 and variants thereof.

The present invention further provides polynucleotides that encode a polypeptide as described above, or a portion thereof (such as a portion encoding at least 15 amino acid residues of a *Chlamydial* protein), expression vectors comprising such polynucleotides and host cells transformed or transfected with such expression vectors.

15 In a related aspect, polynucleotide sequences encoding the above polypeptides, recombinant expression vectors comprising one or more of these polynucleotide sequences and host cells transformed or transfected with such expression vectors are also provided.

In another aspect, the present invention provides fusion proteins

20 comprising an inventive polypeptide, or, alternatively, an inventive polypeptide and a known *Chlamydia* antigen, as well as polynucleotides encoding such fusion proteins, in combination with a physiologically acceptable carrier or immunostimulant for use as pharmaceutical compositions and vaccines thereof.

The present invention further provides pharmaceutical compositions that

25 comprise: (a) an antibody, both polyclonal and monoclonal, or antigen-binding fragment thereof that specifically binds to a *Chlamydial* protein; and (b) a physiologically acceptable carrier. Within other aspects, the present invention provides pharmaceutical compositions that comprise one or more *Chlamydia* polypeptides disclosed herein, for example, a polypeptide of SEQ ID NO: 95-109, 122-124 and 139-

30 140, or a polynucleotide molecule encoding such a polypeptide, such as a polynucleotide sequence of SEQ ID NO: 1-48, 80-94, 114-121 and 125-138, and a physiologically acceptable carrier. The invention also provides compositions for

prophylactic and therapeutic purposes comprising one or more of the disclosed polynucleotides and/or polypeptides and an immunostimulant, e.g., an adjuvant.

In yet another aspect, methods are provided for stimulating an immune response in a patient, e.g., for inducing protective immunity in a patient, comprising
5 administering to a patient an effective amount of one or more of the above pharmaceutical compositions or vaccines.

In yet a further aspect, methods for the treatment of *Chlamydia* infection in a patient are provided, the methods comprising obtaining peripheral blood mononuclear cells (PBMC) from the patient, incubating the PBMC with a polypeptide
10 of the present invention (or a polynucleotide that encodes such a polypeptide) to provide incubated T cells and administering the incubated T cells to the patient. The present invention additionally provides methods for the treatment of *Chlamydia* infection that comprise incubating antigen presenting cells with a polypeptide of the present invention (or a polynucleotide that encodes such a polypeptide) to provide incubated antigen
15 presenting cells and administering the incubated antigen presenting cells to the patient. Proliferated cells may, but need not, be cloned prior to administration to the patient. In certain embodiments, the antigen presenting cells are selected from the group consisting of dendritic cells, macrophages, monocytes, B-cells, and fibroblasts. Compositions for the treatment of *Chlamydia* infection comprising T cells or antigen presenting cells that
20 have been incubated with a polypeptide or polynucleotide of the present invention are also provided. Within related aspects, vaccines are provided that comprise: (a) an antigen presenting cell that expresses a polypeptide as described above and (b) an immunostimulant.

The present invention further provides, within other aspects, methods for
25 removing *Chlamydial*-infected cells from a biological sample, comprising contacting a biological sample with T cells that specifically react with a *Chlamydial* protein, wherein the step of contacting is performed under conditions and for a time sufficient to permit the removal of cells expressing the protein from the sample.

Within related aspects, methods are provided for inhibiting the
30 development of *Chlamydial* infection in a patient, comprising administering to a patient a biological sample treated as described above. In further aspects of the subject invention, methods and diagnostic kits are provided for detecting *Chlamydia* infection

in a patient. In one embodiment, the method comprises: (a) contacting a biological sample with at least one of the polypeptides or fusion proteins disclosed herein; and (b) detecting in the sample the presence of binding agents that bind to the polypeptide or fusion protein, thereby detecting *Chlamydia* infection in the biological sample. Suitable biological samples include whole blood, sputum, serum, plasma, saliva, cerebrospinal fluid and urine. In one embodiment, the diagnostic kits comprise one or more of the polypeptides or fusion proteins disclosed herein in combination with a detection reagent. In yet another embodiment, the diagnostic kits comprise either a monoclonal antibody or a polyclonal antibody that binds with a polypeptide of the present invention.

10 The present invention also provides methods for detecting *Chlamydia* infection comprising: (a) obtaining a biological sample from a patient; (b) contacting the sample with at least two oligonucleotide primers in a polymerase chain reaction, at least one of the oligonucleotide primers being specific for a polynucleotide sequence disclosed herein; and (c) detecting in the sample a polynucleotide sequence that
15 amplifies in the presence of the oligonucleotide primers. In one embodiment, the oligonucleotide primer comprises at least about 10 contiguous nucleotides of a polynucleotide sequence peptide disclosed herein, or of a sequence that hybridizes thereto.

 In a further aspect, the present invention provides a method for detecting
20 *Chlamydia* infection in a patient comprising: (a) obtaining a biological sample from the patient; (b) contacting the sample with an oligonucleotide probe specific for a polynucleotide sequence disclosed herein; and (c) detecting in the sample a polynucleotide sequence that hybridizes to the oligonucleotide probe. In one embodiment, the oligonucleotide probe comprises at least about 15 contiguous
25 nucleotides of a polynucleotide sequence disclosed herein, or a sequence that hybridizes thereto.

 These and other aspects of the present invention will become apparent upon reference to the following detailed description. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated
30 individually.

SEQUENCE IDENTIFIERS

SEQ ID NO:1 sets forth a DNA sequence identified for clone E4-A2-39 (CT10 positive) that is 1311 bp and contains the entire ORF for CT460 (SWIB) and a partial ORF for CT461 (yaeI).

5 SEQ ID NO:2 sets forth a DNA sequence for clone E2-B10-52 (CT10 positive) that has a 1516 bp insert that contains partial ORFs for genes CT827 (nrdA-ribonucleoside reductase large chain) and CT828 (nrdB-ribonucleoside reductase small chain). These genes as were not identified in a Ct L2 library screening.

SEQ ID NO:3 sets forth a DNA sequence for clone E1-B1-80 (CT10 positive) (2397bp) that contains partial ORFs for several genes, CT812 (pmpD), CT015 (phoH ATPase), CT016 (hypothetical protein) and pGp1-D (C. trachomatis plasmid gene).

SEQ ID NO:4 sets forth a DNA sequence for clone E4-F9-4 (CT10, CL8, CT1, CT5, CT13, and CHH037 positive) that contains a 1094 bp insert that has a partial ORF for the gene CT316 (L7/L12 ribosomal protein) as well as a partial ORF for gene

15 CT315 (RNA polymerase beta).

SEQ ID NO:5 sets forth a DNA sequence for clone E2-H6-40 (CT3 positive) that has a 2129 bp insert that contains the entire ORF for the gene CT288 and very small fragments of genes CT287 and CT289. Genes in this clone have not been identified in screening with a Ct L2 library.

20 SEQ ID NO:6 sets forth a DNA sequence for clone E5-D4-2 (CT3, CT10, CT1, CT5, CT12, and CHH037 positive) that has a 1828 bp insert that contains a partial ORF for gene CT378 (pgi), complete ORF for gene CT377 (ltuA) and a complete ORF for the gene CT376 (malate dehydrogenase). In addition, the patient lines CT10, CT1, CT5, CT12, and CHH037 also identified this clone.

25 SEQ ID NO:7 sets forth a DNA sequence for clone E6-C1-31 (CT3 positive) that has a 861 bp insert that contains a partial ORF for gene CT858.

SEQ ID NO:8 sets forth a DNA sequence for clone E9-E11-76 (CT3 positive) that contains a 763 bp insert that is an amino terminal region of the gene for CT798 (Glycogen synthase). This gene was not identified in a previous screening with a

30 Ct L2 library.

SEQ ID NO:9 sets forth a DNA sequence for clone E2-A9-26 (CT1-positive) that contains part of the gene for ORF-3 which is found on the plasmid in *Chlamydia trachomatis*.

SEQ ID NO:10 sets forth a DNA sequence for clone E2-G8-94 (CT1-positive) that has the carboxy terminal end of Lpda gene as well as a partial ORF for CT556.

SEQ ID NO: 11 sets forth a DNA sequence for clone E1-H1-14 (CT1-positive) that has a 1474 bp insert that contains the amino terminal part of an Lpda ORF on the complementary strand.

SEQ ID NO: 12 sets forth a DNA sequence for clone E1-A5-53 (CT1-positive) that contains a 2017 bp insert that has an amino terminal portion of the ORF for dnaK gene on the complementary strand, a partial ORF for the grpE gene (CT395) and a partial ORF for CT166 .

SEQ ID NO: 13 sets forth a DNA sequence for clone E3-A1-50 (positive on CT1 line) that is 1199 bp and contains a carboxy terminal portion of the ORF for CT622.

SEQ ID NO: 14 sets forth a DNA sequence for clone E3-E2-22 that has 877 bp, containing a complete ORF for CT610 on the complementary strand, and was positive on both CT3 and CT10 lines.

SEQ ID NO: 15 sets forth the DNA sequence for clone E5-E2-10 (CT10 positive) which is 427 bp and contains a partial ORF for the major outer membrane protein omp1. SEQ ID NO: 16 sets forth the DNA sequence for clone E2-D5-89 (516bp) which is a CT10 positive clone that contains a partial ORF for pmpD gene (CT812).

SEQ ID NO: 17 sets forth the DNA sequence for clone E4-G9-75 (CT10 positive) which is 723 bp and contains a partial ORF for the amino terminal region of the pmpH gene (CT872).

SEQ ID NO: 18 sets forth the DNA sequence for clone E3-F2-37 (CT10, CT3, CT11, and CT13 positive-1377bp insert) which contains a partial ORF for the tRNA-Trp (CT322) gene and a complete ORF for the gene secE (CT321).

SEQ ID NO: 19 sets forth the DNA sequence for clone E5-A11-8 (CT10 positive-1736 bp) which contains the complete ORF for groES (CT111) and a majority of the ORF for groEL (CT110).

SEQ ID NO: 20 sets forth the DNA sequence for clone E7-H11-61 (CT3 positive-1135 bp) which has partial inserts for *fliA* (CT061), *tyrS* (CT062), *TSA* (CT603) and a hypothetical protein (CT602).

5 SEQ ID NO: 21 sets forth a DNA sequence for clone E6-C8-95 which contains a 731 bp insert that was identified using the donor lines CT3, CT1, and CT12 line. This insert has a carboxy terminal half for the gene for the 60 kDa ORF.

SEQ ID NO: 22 sets forth the DNA sequence for clone E4-D2-79 (CT3 positive) which contains a 1181 bp insert that is a partial ORF for *nrdA* gene. The ORF for this gene was also identified from clone E2-B10-52 (CT10 positive).

10 SEQ ID NO: 23 sets forth the DNA sequence for clone E1-F9-79 (167 bp; CT11 positive) which contains a partial ORF for the gene CT133 on the complementary strand. CT133 is a predicted rRNA methylase.

SEQ ID NO: 24 sets forth the DNA sequence for clone E2-G12-52 (1265 bp; CT11 positive) which contains a partial ORF for *clpB*, a protease ATPase.

15 SEQ ID NO: 25 sets forth the DNA sequence for clone E4-H3-56 (463 bp insert; CT1 positive) which contains a partial ORF for the *TSA* gene (CT603) on the complementary strand.

SEQ ID NO: 26 sets forth the DNA sequence for clone E5-E9-3 (CT1 positive) that contains a 636 bp insert partially encoding the ORF for *dnaK* like gene.
20 Part of this sequence was also identified in clone E1-A5-53.

SEQ ID NO:27 sets forth the full-length serovar E DNA sequence of CT875.

SEQ ID NO:28 sets for the full-length serovar E DNA sequence of CT622.

25 SEQ ID NO:29 sets forth the DNA sequence for clone E3-B4-18 (CT1 positive) that contains a 1224 bp insert containing 4 ORFs. The complete ORF for CT772, and the partial ORFs of CT771, CT191, and CT190.

SEQ ID NO:30 sets forth the DNA sequence for the clone E9-E10-51 (CT10 positive) that contains an 883 bp insert containing two partial ORF, CT680 and
30 CT679.

SEQ ID NO:31 sets forth the DNA sequence of the clone E9-D5-8 (CT10, CTCT1, CT4, and CT11 positive) that contains a 393 bp insert containing the partial ORF for CT680.

5 SEQ ID NO:32 sets forth the DNA sequence of the clone E7-B1-16 (CT10, CT3, CT5, CT11, CT13, and CHH037 positive) that contains a 2577 bp insert containing three ORFs, two full length ORFs for CT694 and CT695 and the third containing the N-terminal portion of CT969.

SEQ ID NO:33 sets forth the DNA sequence of the clone E9-G2-93 (CT10 positive) that contains a 554 bp insert containing a partial ORF for CT178.

10 SEQ ID NO:34 sets forth the DNA sequence of the clone E5-A8-85 (CT1 positive) that contains a 1433 bp insert containing two partial ORFs for CT875 and CT001.

SEQ ID NO:35 sets forth the DNA sequence of the clone E10-C6-45 (CT3 positive) that contains a 196 bp insert containing a partial ORF for CT827.

15 SEQ ID NO:36 sets forth the DNA sequence of the clone E7-H11-10 (CT3 positive) that contains a 1990 bp insert containing the partial ORFs of CT610 and CT613 and the complete ORFs of CT611 and CT612.

SEQ ID NO:37 sets forth the DNA sequence of the clone E2-F7-11 (CT3 and CT10 positive) that contains a 2093 bp insert. It contains a large region of CT609, a complete ORF for CT610 and a partial ORF for CT611.

20 SEQ ID NO:38 sets forth the DNA sequence of the clone E3-A3-31 (CT1 positive) that contains an 1834 bp insert containing a large region of CT622.

SEQ ID NO:39 sets forth the DNA sequence of the clone E1-G9-23 (CT3 positive) that contains an 1180 bp insert containing almost the entire ORF for CT798.

25 SEQ ID NO:40 sets forth the DNA sequence of the clone E4-D6-21 (CT3 positive) that contains a 1297 bp insert containing the partial ORFs of CT329 and CT327 and the complete ORF of CT328.

30 SEQ ID NO:41 sets forth the DNA sequence of the clone E3-F3-18 (CT1 positive) that contains an 1141 bp insert containing the partial ORF of CT871.

SEQ ID NO:42 sets forth the DNA sequence of the clone E10-B2-57 (CT10 positive) that contains an 822 bp insert containing the complete ORF of CT066.

SEQ ID NO:43 sets forth the DNA sequence of the clone E3-F3-7 (CT1 positive) that contains a 1643 bp insert containing the partial ORFs of CT869 and CT870.

SEQ ID NO:44 sets forth the DNA sequence of the clone E10-H8-1
5 (CT3 and CT10 positive) that contains an 1862 bp insert containing the partial ORFs of CT871 and CT872.

SEQ ID NO:45 sets forth the DNA sequence of the clone E3-D10-46 (CT1, CT3, CT4, CT11, and CT12 positive) that contains a 1666 bp insert containing the partial ORFs for CT770 and CT773 and the complete ORFs for CT771 and CT722.

10 SEQ ID NO:46 sets forth the DNA sequence of the clone E2-D8-19 (CT1 positive) that contains a 2010 bp insert containing partial ORFs, ORF3 and ORF6, and complete ORFs, ORF4 and ORF5.

SEQ ID NO:47 sets forth the DNA sequence of the clone E4-C3-40 (CT10 positive) that contains a 2044 bp insert containing the partial ORF for CT827
15 and a complete ORF for CT828.

SEQ ID NO:48 sets forth the DNA sequence of the clone E3-H6-10 (CT12 positive) that contains a 3743 bp insert containing the partial ORFs for CT223 and CT229 and the complete ORFs for CT224 and CT224, CT225, CT226, CT227, and CT228.

20 SEQ ID NO:49 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0454 of the Chlamydia trachomatis gene CT872.

SEQ ID NO:50 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0187, of the Chlamydia trachomatis gene CT133.

SEQ ID NO:51 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0075 of the Chlamydia trachomatis gene CT321.
25

SEQ ID NO:52 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0074, of the Chlamydia trachomatis gene CT322.

SEQ ID NO:53 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0948, of the Chlamydia trachomatis gene CT798.

30 SEQ ID NO:54 sets forth the DNA sequence for the Chlamydia pneumoniae homologue, CPn0985, of the Chlamydia trachomatis gene CT828.

SEQ ID NO:55 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0984, of the *Chlamydia trachomatis* gene CT827.

SEQ ID NO:56 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0062, of the *Chlamydia trachomatis* gene CT289.

5 SEQ ID NO:57 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn00065, of the *Chlamydia trachomatis* gene CT288.

SEQ ID NO:58 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0438, of the *Chlamydia trachomatis* gene CT287.

10 SEQ ID NO:59 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0963, of the *Chlamydia trachomatis* gene CT812.

SEQ ID NO:60 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0778, of the *Chlamydia trachomatis* gene CT603.

SEQ ID NO:61 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0503, of the *Chlamydia trachomatis* gene CT396.

15 SEQ ID NO:62 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn1016, of the *Chlamydia trachomatis* gene CT858.

SEQ ID NO:63 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0728, of the *Chlamydia trachomatis* gene CT622.

20 SEQ ID NO:64 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0557, of the *Chlamydia trachomatis* gene CT460.

SEQ ID NO:65 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0454, of the *Chlamydia trachomatis* gene CT872.

SEQ ID NO:66 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0187, of the *Chlamydia trachomatis* gene CT133.

25 SEQ ID NO:67 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0075, of the *Chlamydia trachomatis* gene CT321.

SEQ ID NO:68 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0074, of the *Chlamydia trachomatis* gene CT322.

30 SEQ ID NO:69 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0948, of the *Chlamydia trachomatis* gene CT798.

SEQ ID NO:70 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0985, of the *Chlamydia trachomatis* gene CT828.

SEQ ID NO:71 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0984, of the *Chlamydia trachomatis* gene CT827.

SEQ ID NO:72 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0062, of the *Chlamydia trachomatis* gene CT289.

5 SEQ ID NO:73 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0065, of the *Chlamydia trachomatis* gene CT288.

SEQ ID NO:74 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0438, of the *Chlamydia trachomatis* gene CT287.

10 SEQ ID NO:75 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0963, of the *Chlamydia trachomatis* gene CT812.

SEQ ID NO:76 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0778, of the *Chlamydia trachomatis* gene CT603.

SEQ ID NO:77 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn1016, of the *Chlamydia trachomatis* gene CT858.

15 SEQ ID NO:78 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0728, of the *Chlamydia trachomatis* gene CT622.

SEQ ID NO:79 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0557, of the *Chlamydia trachomatis* gene CT460.

20 SEQ ID NO:80 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT872.

SEQ ID NO:81 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT828.

SEQ ID NO:82 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT827.

25 SEQ ID NO:83 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT812.

SEQ ID NO:84 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT798.

30 SEQ ID NO:85 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT681 (MompF).

SEQ ID NO:86 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT603.

SEQ ID NO:87 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT460.

SEQ ID NO:88 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT322.

5 SEQ ID NO:89 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT321.

SEQ ID NO:90 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT289.

10 SEQ ID NO:91 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT288.

SEQ ID NO:92 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT287.

SEQ ID NO:93 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT133.

15 SEQ ID NO:94 sets forth the full-length serovar D DNA sequence of the *Chlamydia trachomatis* gene CT113.

SEQ ID NO:95 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT872.

20 SEQ ID NO:96 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT828.

SEQ ID NO:97 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT827.

SEQ ID NO:98 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT812.

25 SEQ ID NO:99 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT798.

SEQ ID NO:100 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT681.

30 SEQ ID NO:101 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT603.

SEQ ID NO:102 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT460.

SEQ ID NO:103 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT322.

SEQ ID NO:104 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT321.

5 SEQ ID NO:105 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT289.

SEQ ID NO:106 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT288.

10 SEQ ID NO:107 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT287.

SEQ ID NO:108 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT133.

SEQ ID NO:109 sets forth the full-length serovar D amino acid sequence of the *Chlamydia trachomatis* gene CT113.

15 SEQ ID NO:110 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0695, of the *Chlamydia trachomatis* gene CT681.

SEQ ID NO:111 sets forth the DNA sequence for the *Chlamydia pneumoniae* homologue, CPn0144, of the *Chlamydia trachomatis* gene CT113.

20 SEQ ID NO:112 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0695, of the *Chlamydia trachomatis* gene CT681.

SEQ ID NO:113 sets forth the amino acid sequence for the *Chlamydia pneumoniae* homologue, CPn0144, of the *Chlamydia trachomatis* gene CT113.

SEQ ID NO:114 sets forth the DNA sequence of the clone E7-B12-65 (CHH037 positive) that contains a 1179 bp insert containing complete ORF for 376.

25 SEQ ID NO:115 sets forth the DNA sequence of the clone E4-H9-83 (CHH037 positive) that contains the partial ORF for the heat shock protein GroEL (CT110).

SEQ ID NO:116 sets forth the DNA sequence of the clone E9-B10-52 (CHH037 positive) that contains the partial ORF for the the gene yscC (CT674).

30 SEQ ID NO:117 sets forth the DNA sequence of the clone E7-A7-79 (CHH037 positive) that contains the complete ORF for the histone like development

gene *hctA* (CT743) and a partial ORF for the rRNA methyltransferase gene *ygca* (CT742).

SEQ ID NO:118 sets forth the DNA sequence of the clone E2-D11-18 (CHH037 positive) that contains the partial ORF for *hctA* (CT743).

5 SEQ ID NO:119 sets forth the DNA sequence for the *Chlamydia trachomatis* serovar E hypothetical protein CT694.

SEQ ID NO:120 sets forth the DNA sequence for the *Chlamydia trachomatis* serovar E hypothetical protein CT695.

10 SEQ ID NO:121 sets forth the DNA sequence for the *Chlamydia trachomatis* serovar E L1 ribosomal protein.

SEQ ID NO:122 sets forth the amino acid sequence for the *Chlamydia trachomatis* serovar E hypothetical protein CT694.

SEQ ID NO:123 sets forth the amino acid sequence for the *Chlamydia trachomatis* serovar E hypothetical protein CT695.

15 SEQ ID NO:124 sets forth the amino acid sequence for the *Chlamydia trachomatis* serovar E L1 ribosomal protein.

SEQ ID NO:125 sets forth the DNA sequence of the clone E9-H6-15 (CT3 positive) that contains the partial ORF for the *pmpB* gene (CT413).

20 SEQ ID NO:126 sets forth the DNA sequence of the clone E3-D10-87 (CT1 positive) that contains the partial ORFs for the hypothetical genes CT388 and CT389.

SEQ ID NO:127 sets forth the DNA sequence of the clone E9-D6-43 (CT3 positive) that contains the partial ORF for the CT858.

25 SEQ ID NO:128 sets forth the DNA sequence of the clone E3-D10-4 (CT1 positive) that contains the partial ORF for *pGP3-D*, an ORF encoded on the plasmid *pCHL1*.

SEQ ID NO:129 sets forth the DNA sequence of the clone E3-G8-7 (CT1 positive) that contains the partial ORFs for the CT557 (*LpdA*) and CT558 (*LipA*).

30 SEQ ID NO:130 sets forth the DNA sequence of the clone E3-F11-32 (CT1 positive) that contains the partial ORF for *pmpD* (CT812).

SEQ ID NO:131 sets forth the DNA sequence of the clone E2-F8-5 (CT12 positive) that contains the complete ORF for the 15 kDa ORF (CT442) and a partial ORF for the 60kDa ORF (CT443).

SEQ ID NO:132 sets forth the DNA sequence of the clone E2-G4-39
5 (CT12 positive) that contains the partial ORF for the 60kDa ORF (CT443).

SEQ ID NO:133 sets forth the DNA sequence of the clone E9-D1-16 (CT10 positive) that contains the partial ORF for pmpH (CT872).

SEQ ID NO:134 sets forth the DNA sequence of the clone E3-F3-6 (CT1 positive) that contains the partial ORFs for the genes accB (CT123), L1 ribosomal
10 (CT125) and S9 ribosomal (CT126).

SEQ ID NO:135 sets forth the DNA sequence of the clone E2-D4-70 (CT12 positive) that contains the partial ORF for the pmpC gene (CT414).

SEQ ID NO:136 sets forth the DNA sequence of the clone E5-A1-79 (CT1 positive) that contains the partial ORF for ydhO (CT127), a complete ORF for S9
15 ribosomal gene (CT126), a complete ORF for the L1 ribosomal gene (CT125) and a partial ORF for accC (CT124).

SEQ ID NO:137 sets forth the DNA sequence of the clone E1-F7-16 (CT12, CT3, and CT11 positive) that contains the partial ORF for the ftsH gene (CT841) and the entire ORF for the pnp gene (CT842).

SEQ ID NO:138 sets forth the DNA sequence of the clone E1-D8-62
20 (CT12 positive) that contains the partial ORFs for the ftsH gene (CT841) and for the pnp gene (CT842).

SEQ ID NO:139 sets forth the amino acid sequence for the serovar E protein CT875.

SEQ ID NO:140 sets forth the amino acid sequence for the serovar E
25 protein CT622.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

As noted above, the present invention is generally directed to
30 compositions and methods for the diagnosis and treatment of Chlamydial infection. In one aspect, the compositions of the subject invention include polypeptides that comprise at least one immunogenic portion of a *Chlamydia* antigen, or a variant thereof.

In specific embodiments, the subject invention discloses polypeptides comprising an immunogenic portion of a *Chlamydia* antigen, wherein the *Chlamydia* antigen comprises an amino acid sequence encoded by a polynucleotide molecule including a sequence selected from the group consisting of (a) nucleotide sequences
5 recited in SEQ ID NO:1-48, 114-121, and 125-138 (b) the complements of said nucleotide sequences, and (c) variants of such sequences.

POLYNUCLEOTIDE COMPOSITIONS

As used herein, the terms "DNA segment" and "polynucleotide" refer to a DNA molecule that has been isolated free of total genomic DNA of a particular
10 species. Therefore, a DNA segment encoding a polypeptide refers to a DNA segment that contains one or more coding sequences yet is substantially isolated away from, or purified free from, total genomic DNA of the species from which the DNA segment is obtained. Included within the terms "DNA segment" and "polynucleotide" are DNA segments and smaller fragments of such segments, and also recombinant vectors,
15 including, for example, plasmids, cosmids, phagemids, phage, viruses, and the like.

As will be understood by those skilled in the art, the DNA segments of this invention can include genomic sequences, extra-genomic and plasmid-encoded sequences and smaller engineered gene segments that express, or may be adapted to express, proteins, polypeptides, peptides and the like. Such segments may be naturally
20 isolated, or modified synthetically by the hand of man.

"Isolated," as used herein, means that a polynucleotide is substantially away from other coding sequences, and that the DNA segment does not contain large portions of unrelated coding DNA, such as large chromosomal fragments or other functional genes or polypeptide coding regions. Of course, this refers to the DNA
25 segment as originally isolated, and does not exclude genes or coding regions later added to the segment by the hand of man.

As will be recognized by the skilled artisan, polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. RNA molecules include HnRNA molecules,
30 which contain introns and correspond to a DNA molecule in a one-to-one manner, and mRNA molecules, which do not contain introns. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present

invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials.

Polynucleotides may comprise a native Chlamydia sequence or may comprise a variant, or a biological or antigenic functional equivalent of such a sequence.

5 Polynucleotide variants may contain one or more substitutions, additions, deletions and/or insertions, as further described below, preferably such that the immunogenicity of the encoded polypeptide is not diminished, relative to a native Chlamydia protein. The effect on the immunogenicity of the encoded polypeptide may generally be assessed as described herein. The term "variants" also encompasses homologous genes of
10 xenogenic origin.

When comparing polynucleotide or polypeptide sequences, two sequences are said to be "identical" if the sequence of nucleotides or amino acids in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the
15 sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

20 Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O. (1978) A model of evolutionary change in proteins – Matrices for detecting distant relationships.
25 In Dayhoff, M.O. (ed.) Atlas of Protein Sequence and Structure, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylogenies pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989) *CABIOS* 5:151-153; Myers, E.W. and Muller W. (1988) *CABIOS* 4:11-17; Robinson,
30 E.D. (1971) *Comb. Theor* 11:105; Santou, N. Nes, M. (1987) *Mol. Biol. Evol.* 4:406-425; Sneath, P.H.A. and Sokal, R.R. (1973) *Numerical Taxonomy – the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, CA; Wilbur, W.J. and

Lipman, D.J. (1983) *Proc. Natl. Acad. Sci. USA* 80:726-730.

Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL. Math* 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity methods of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

One preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) *Nucl. Acids Res.* 25:3389-3402 and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410, respectively. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides and polypeptides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. In one illustrative example, cumulative scores can be calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix can be used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments, (B) of 50, expectation (E) of 10, M=5, N=-4 and a comparison of both strands.

Preferably, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide or polypeptide sequence in the comparison

window may comprise additions or deletions (*i.e.*, gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical
5 nucleic acid bases or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (*i.e.*, the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

Therefore, the present invention encompasses polynucleotide and
10 polypeptide sequences having substantial identity to the sequences disclosed herein, for example those comprising at least 50% sequence identity, preferably at least 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher, sequence identity compared to a polynucleotide or polypeptide sequence of this invention using the methods described herein, (e.g., BLAST analysis using standard parameters, as
15 described below). One skilled in this art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like.

In additional embodiments, the present invention provides isolated
20 polynucleotides and polypeptides comprising various lengths of contiguous stretches of sequence identical to or complementary to one or more of the sequences disclosed herein. For example, polynucleotides are provided by this invention that comprise at least about 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 400, 500 or 1000 or more contiguous nucleotides of one or more of the sequences disclosed herein as well as all
25 intermediate lengths there between. It will be readily understood that "intermediate lengths", in this context, means any length between the quoted values, such as 16, 17, 18, 19, *etc.*; 21, 22, 23, *etc.*; 30, 31, 32, *etc.*; 50, 51, 52, 53, *etc.*; 100, 101, 102, 103, *etc.*; 150, 151, 152, 153, *etc.*; including all integers through 200-500; 500-1,000, and the like.

30 The polynucleotides of the present invention, or fragments thereof, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, polyadenylation signals, additional restriction enzyme

sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. For example, illustrative DNA segments with total lengths of about 10,000, about 5000, about 3000, about 2,000, about 1,000, about 500, about 200, about 100, about 50 base pairs in length, and the like, (including all intermediate lengths) are contemplated to be useful in many implementations of this invention.

In other embodiments, the present invention is directed to polynucleotides that are capable of hybridizing under moderately stringent conditions to a polynucleotide sequence provided herein, or a fragment thereof, or a complementary sequence thereof. Hybridization techniques are well known in the art of molecular biology. For purposes of illustration, suitable moderately stringent conditions for testing the hybridization of a polynucleotide of this invention with other polynucleotides include prewashing in a solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5 X SSC, overnight; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS.

Moreover, it will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention. Further, alleles of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention. Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridization, amplification and/or database sequence comparison).

PROBES AND PRIMERS

In other embodiments of the present invention, the polynucleotide sequences provided herein can be advantageously used as probes or primers for nucleic acid hybridization. As such, it is contemplated that nucleic acid segments that comprise
5 a sequence region of at least about 15 nucleotide long contiguous sequence that has the same sequence as, or is complementary to, a 15 nucleotide long contiguous sequence disclosed herein will find particular utility. Longer contiguous identical or complementary sequences, *e.g.*, those of about 20, 30, 40, 50, 100, 200, 500, 1000 (including all intermediate lengths) and even up to full length sequences will also be of
10 use in certain embodiments.

The ability of such nucleic acid probes to specifically hybridize to a sequence of interest will enable them to be of use in detecting the presence of complementary sequences in a given sample. However, other uses are also envisioned, such as the use of the sequence information for the preparation of mutant species
15 primers, or primers for use in preparing other genetic constructions.

Polynucleotide molecules having sequence regions consisting of contiguous nucleotide stretches of 10-14, 15-20, 30, 50, or even of 100-200 nucleotides or so (including intermediate lengths as well), identical or complementary to a polynucleotide sequence disclosed herein, are particularly contemplated as hybridization
20 probes for use in, *e.g.*, Southern and Northern blotting. This would allow a gene product, or fragment thereof, to be analyzed, both in diverse cell types and also in various bacterial cells. The total size of fragment, as well as the size of the complementary stretch(es), will ultimately depend on the intended use or application of the particular nucleic acid segment. Smaller fragments will generally find use in
25 hybridization embodiments, wherein the length of the contiguous complementary region may be varied, such as between about 15 and about 100 nucleotides, but larger contiguous complementarity stretches may be used, according to the length complementary sequences one wishes to detect.

The use of a hybridization probe of about 15-25 nucleotides in length
30 allows the formation of a duplex molecule that is both stable and selective. Molecules having contiguous complementary sequences over stretches greater than 15 bases in length are generally preferred, though, in order to increase stability and selectivity of the

hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained. One will generally prefer to design nucleic acid molecules having gene-complementary stretches of 15 to 25 contiguous nucleotides, or even longer where desired.

5 Hybridization probes may be selected from any portion of any of the sequences disclosed herein. All that is required is to review the sequence set forth in SEQ ID NO:1-48, 114-121, and 125-138, or to any continuous portion of the sequence, from about 15-25 nucleotides in length up to and including the full length sequence, that one wishes to utilize as a probe or primer. The choice of probe and primer sequences
10 may be governed by various factors. For example, one may wish to employ primers from towards the termini of the total sequence.

 Small polynucleotide segments or fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer. Also, fragments may be
15 obtained by application of nucleic acid reproduction technology, such as the PCR™ technology of U. S. Patent 4,683,202 (incorporated herein by reference), by introducing selected sequences into recombinant vectors for recombinant production, and by other recombinant DNA techniques generally known to those of skill in the art of molecular biology.

20 The nucleotide sequences of the invention may be used for their ability to selectively form duplex molecules with complementary stretches of the entire gene or gene fragments of interest. Depending on the application envisioned, one will typically desire to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence. For applications requiring high selectivity,
25 one will typically desire to employ relatively stringent conditions to form the hybrids, *e.g.*, one will select relatively low salt and/or high temperature conditions, such as provided by a salt concentration of from about 0.02 M to about 0.15 M salt at temperatures of from about 50°C to about 70°C. Such selective conditions tolerate little, if any, mismatch between the probe and the template or target strand, and would
30 be particularly suitable for isolating related sequences.

 Of course, for some applications, for example, where one desires to prepare mutants employing a mutant primer strand hybridized to an underlying

template, less stringent (reduced stringency) hybridization conditions will typically be needed in order to allow formation of the heteroduplex. In these circumstances, one may desire to employ salt conditions such as those of from about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20°C to about 55°C. Cross-hybridizing species
5 can thereby be readily identified as positively hybridizing signals with respect to control hybridizations. In any case, it is generally appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide, which serves to destabilize the hybrid duplex in the same manner as increased temperature. Thus, hybridization conditions can be readily manipulated, and thus will generally be a
10 method of choice depending on the desired results.

POLYNUCLEOTIDE IDENTIFICATION AND CHARACTERIZATION

Polynucleotides may be identified, prepared and/or manipulated using any of a variety of well established techniques. For example, a polynucleotide may be
15 identified, by screening a microarray of cDNAs for Chlamydia expression. Such screens may be performed, for example, using a Synteni microarray (Palo Alto, CA) according to the manufacturer's instructions (and essentially as described by Schena et al., *Proc. Natl. Acad. Sci. USA* 93:10614-10619, 1996 and Heller et al., *Proc. Natl. Acad. Sci. USA* 94:2150-2155, 1997). Alternatively, polynucleotides may be amplified
20 from cDNA prepared from cells expressing the proteins described herein. Such polynucleotides may be amplified via polymerase chain reaction (PCR). For this approach, sequence-specific primers may be designed based on the sequences provided herein, and may be purchased or synthesized.

An amplified portion of a polynucleotide of the present invention may be
25 used to isolate a full length gene from a suitable library (e.g., Chlamydia cDNA library) using well known techniques. Within such techniques, a library (cDNA or genomic) is screened using one or more polynucleotide probes or primers suitable for amplification. Preferably, a library is size-selected to include larger molecules. Random primed libraries may also be preferred for identifying 5' and upstream regions of genes.
30 Genomic libraries are preferred for obtaining introns and extending 5' sequences.

For hybridization techniques, a partial sequence may be labeled (e.g., by nick-translation or end-labeling with ³²P) using well known techniques. A bacterial or

bacteriophage library is then generally screened by hybridizing filters containing denatured bacterial colonies (or lawns containing phage plaques) with the labeled probe (see Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989). Hybridizing colonies or plaques are
5 selected and expanded, and the DNA is isolated for further analysis. cDNA clones may be analyzed to determine the amount of additional sequence by, for example, PCR using a primer from the partial sequence and a primer from the vector. Restriction maps and partial sequences may be generated to identify one or more overlapping clones. The complete sequence may then be determined using standard techniques, which may
10 involve generating a series of deletion clones. The resulting overlapping sequences can then be assembled into a single contiguous sequence. A full length cDNA molecule can be generated by ligating suitable fragments, using well known techniques.

Alternatively, there are numerous amplification techniques for obtaining a full length coding sequence from a partial cDNA sequence. Within such techniques,
15 amplification is generally performed via PCR. Any of a variety of commercially available kits may be used to perform the amplification step. Primers may be designed using, for example, software well known in the art. Primers are preferably 22-30 nucleotides in length, have a GC content of at least 50% and anneal to the target sequence at temperatures of about 68°C to 72°C. The amplified region may be
20 sequenced as described above, and overlapping sequences assembled into a contiguous sequence.

One such amplification technique is inverse PCR (see Triglia et al., *Nucl. Acids Res.* 16:8186, 1988), which uses restriction enzymes to generate a fragment in the known region of the gene. The fragment is then circularized by intramolecular ligation
25 and used as a template for PCR with divergent primers derived from the known region. Within an alternative approach, sequences adjacent to a partial sequence may be retrieved by amplification with a primer to a linker sequence and a primer specific to a known region. The amplified sequences are typically subjected to a second round of amplification with the same linker primer and a second primer specific to the known
30 region. A variation on this procedure, which employs two primers that initiate extension in opposite directions from the known sequence, is described in WO 96/38591. Another such technique is known as "rapid amplification of cDNA ends" or

RACE. This technique involves the use of an internal primer and an external primer, which hybridizes to a polyA region or vector sequence, to identify sequences that are 5' and 3' of a known sequence. Additional techniques include capture PCR (Lagerstrom et al., *PCR Methods Applic. 1*:111-19, 1991) and walking PCR (Parker et al., *Nucl. Acids. Res. 19*:3055-60, 1991). Other methods employing amplification may also be employed to obtain a full length cDNA sequence.

In certain instances, it is possible to obtain a full length cDNA sequence by analysis of sequences provided in an expressed sequence tag (EST) database, such as that available from GenBank. Searches for overlapping ESTs may generally be performed using well known programs (e.g., NCBI BLAST searches), and such ESTs may be used to generate a contiguous full length sequence. Full length DNA sequences may also be obtained by analysis of genomic fragments.

POLYNUCLEOTIDE EXPRESSION IN HOST CELLS

In other embodiments of the invention, polynucleotide sequences or fragments thereof which encode polypeptides of the invention, or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct expression of a polypeptide in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences that encode substantially the same or a functionally equivalent amino acid sequence may be produced and these sequences may be used to clone and express a given polypeptide.

As will be understood by those of skill in the art, it may be advantageous in some instances to produce polypeptide-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

Moreover, the polynucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter polypeptide encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. For example, DNA shuffling by random fragmentation and PCR reassembly of gene

fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. In addition, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, or introduce mutations, and so forth.

5 In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of polypeptide activity, it may be useful to encode a chimeric protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site
10 located between the polypeptide-encoding sequence and the heterologous protein sequence, so that the polypeptide may be cleaved and purified away from the heterologous moiety.

Sequences encoding a desired polypeptide may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M. H. et al.
15 (1980) *Nucl. Acids Res. Symp. Ser.* 215-223, Horn, T. et al. (1980) *Nucl. Acids Res. Symp. Ser.* 225-232). Alternatively, the protein itself may be produced using chemical methods to synthesize the amino acid sequence of a polypeptide, or a portion thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J. Y. et al. (1995) *Science* 269:202-204) and automated synthesis may be
20 achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer, Palo Alto, CA).

A newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, T. (1983) *Proteins, Structures and Molecular Principles*, WH Freeman and Co., New York, N.Y.) or other comparable
25 techniques available in the art. The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure). Additionally, the amino acid sequence of a polypeptide, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

30 In order to express a desired polypeptide, the nucleotide sequences encoding the polypeptide, or functional equivalents, may be inserted into appropriate expression

vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described in Sambrook, J. et al. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Plainview, N.Y., and Ausubel, F. M. et al. (1989) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, N.Y.

A variety of expression vector/host systems may be utilized to contain and express polynucleotide sequences. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems.

The "control elements" or "regulatory sequences" present in an expression vector are those non-translated regions of the vector--enhancers, promoters, 5' and 3' untranslated regions--which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the PBLUESCRIPT phagemid (Stratagene, La Jolla, Calif.) or PSPO1 plasmid (Gibco BRL, Gaithersburg, MD) and the like may be used. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are generally preferred. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding a polypeptide, vectors based on SV40 or EBV may be advantageously used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected

depending upon the use intended for the expressed polypeptide. For example, when large quantities are needed, for example for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the sequence encoding the polypeptide of interest may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S. M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509); and the like. pGEX Vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast, *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) *Methods Enzymol.* 153:516-544.

In cases where plant expression vectors are used, the expression of sequences encoding polypeptides may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, for example, Hobbs, S. or Murry, L. E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, N.Y.; pp. 191-196).

An insect system may also be used to express a polypeptide of interest. For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The sequences encoding the polypeptide may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the polypeptide-encoding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda* cells or *Trichoplusia* larvae in which the polypeptide of interest may be expressed (Engelhard, E. K. et al. (1994) *Proc. Natl. Acad. Sci.* 91 :3224-3227).

In mammalian host cells, a number of viral-based expression systems are generally available. For example, in cases where an adenovirus is used as an expression vector, sequences encoding a polypeptide of interest may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing the polypeptide in infected host cells (Logan, J. and Shenk, T. (1984) *Proc. Natl. Acad. Sci.* 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Specific initiation signals may also be used to achieve more efficient translation of sequences encoding a polypeptide of interest. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding the polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation.

5 Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as CHO, HeLa, MDCK, HEK293, and WI38, which have specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

10 For long-term, high-yield production of recombinant proteins, stable expression is generally preferred. For example, cell lines which stably express a polynucleotide of interest may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be

15 allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

20 Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. et al. (1990) *Cell* 22:817-23) genes which can be employed in tk.sup.- or aprt.sup.- cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can

25 be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); npt, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin, F. et al (1981) *J. Mol. Biol.* 150:1-14); and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, *supra*).

30 Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman, S. C. and R. C. Mulligan (1988) *Proc. Natl. Acad. Sci.*

85:8047-51). Recently, the use of visible markers has gained popularity with such markers as anthocyanins, beta-glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C. A. et al. (1995) *Methods Mol. Biol.* 55:121-131).

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding a polypeptide is inserted within a marker gene sequence, recombinant cells containing sequences can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a polypeptide-encoding sequence under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells which contain and express a desired polynucleotide sequence may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products, using either polyclonal or monoclonal antibodies specific for the product are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on a given polypeptide may be preferred for some applications, but a competitive binding assay may also be employed. These and other assays are described, among other places, in Hampton, R. et al. (1990; *Serological Methods, a Laboratory Manual*, APS Press, St Paul, Minn.) and Maddox, D. E. et al. (1983; *J. Exp. Med.* 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to

polynucleotides include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences, or any portions thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits. Suitable reporter molecules or labels, which may be used include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with a polynucleotide sequence of interest may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides of the invention may be designed to contain signal sequences which direct secretion of the encoded polypeptide through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding a polypeptide of interest to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, Calif.) between the purification domain and the encoded polypeptide may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a polypeptide of interest and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography) as described in Porath, J. et al. (1992, *Prot. Exp. Purif.* 3:263-281) while the enterokinase cleavage site provides a means for purifying the desired polypeptide from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D. J.

et al. (1993; *DNA Cell Biol.* 12:441-453).

In addition to recombinant production methods, polypeptides of the invention, and fragments thereof, may be produced by direct peptide synthesis using solid-phase techniques (Merrifield J. (1963) *J. Am. Chem. Soc.* 85:2149-2154). Protein
5 synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Alternatively, various fragments may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

10

SITE-SPECIFIC MUTAGENESIS

Site-specific mutagenesis is a technique useful in the preparation of individual peptides, or biologically functional equivalent polypeptides, through specific mutagenesis of the underlying polynucleotides that encode them. The technique,
15 well-known to those of skill in the art, further provides a ready ability to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as
20 well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Mutations may be employed in a selected polynucleotide sequence to improve, alter, decrease, modify, or otherwise change the properties of the polynucleotide itself, and/or alter the properties, activity, composition,
25 stability, or primary sequence of the encoded polypeptide.

In certain embodiments of the present invention, the inventors contemplate the mutagenesis of the disclosed polynucleotide sequences to alter one or more properties of the encoded polypeptide, such as the antigenicity of a polypeptide vaccine. The techniques of site-specific mutagenesis are well-known in the art, and are
30 widely used to create variants of both polypeptides and polynucleotides. For example, site-specific mutagenesis is often used to alter a specific portion of a DNA molecule. In such embodiments, a primer comprising typically about 14 to about 25 nucleotides or so

in length is employed, with about 5 to about 10 residues on both sides of the junction of the sequence being altered.

As will be appreciated by those of skill in the art, site-specific mutagenesis techniques have often employed a phage vector that exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially-available and their use is generally well-known to those skilled in the art. Double-stranded plasmids are also routinely employed in site directed mutagenesis that eliminates the step of transferring the gene of interest from a plasmid to a phage.

In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector or melting apart of two strands of a double-stranded vector that includes within its sequence a DNA sequence that encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

The preparation of sequence variants of the selected peptide-encoding DNA segments using site-directed mutagenesis provides a means of producing potentially useful species and is not meant to be limiting as there are other ways in which sequence variants of peptides and the DNA sequences encoding them may be obtained. For example, recombinant vectors encoding the desired peptide sequence may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants. Specific details regarding these methods and protocols are found in the teachings of Maloy *et al.*, 1994; Segal, 1976; Prokop and Bajpai, 1991; Kuby, 1994; and Maniatis *et al.*, 1982, each incorporated herein by reference, for that purpose.

As used herein, the term "oligonucleotide directed mutagenesis procedure" refers to template-dependent processes and vector-mediated propagation which result in an increase in the concentration of a specific nucleic acid molecule

relative to its initial concentration, or in an increase in the concentration of a detectable signal, such as amplification. As used herein, the term "oligonucleotide directed mutagenesis procedure" is intended to refer to a process that involves the template-dependent extension of a primer molecule. The term template dependent process refers to nucleic acid synthesis of an RNA or a DNA molecule wherein the sequence of the newly synthesized strand of nucleic acid is dictated by the well-known rules of complementary base pairing (see, for example, Watson, 1987). Typically, vector mediated methodologies involve the introduction of the nucleic acid fragment into a DNA or RNA vector, the clonal amplification of the vector, and the recovery of the amplified nucleic acid fragment. Examples of such methodologies are provided by U. S. Patent No. 4,237,224, specifically incorporated herein by reference in its entirety.

POLYNUCLEOTIDE AMPLIFICATION TECHNIQUES

A number of template dependent processes are available to amplify the target sequences of interest present in a sample. One of the best known amplification methods is the polymerase chain reaction (PCRTM) which is described in detail in U.S. Patent Nos. 4,683,195, 4,683,202 and 4,800,159, each of which is incorporated herein by reference in its entirety. Briefly, in PCRTM, two primer sequences are prepared which are complementary to regions on opposite complementary strands of the target sequence. An excess of deoxynucleoside triphosphates is added to a reaction mixture along with a DNA polymerase (e.g., *Taq* polymerase). If the target sequence is present in a sample, the primers will bind to the target and the polymerase will cause the primers to be extended along the target sequence by adding on nucleotides. By raising and lowering the temperature of the reaction mixture, the extended primers will dissociate from the target to form reaction products, excess primers will bind to the target and to the reaction product and the process is repeated. Preferably reverse transcription and PCRTM amplification procedure may be performed in order to quantify the amount of mRNA amplified. Polymerase chain reaction methodologies are well known in the art.

Another method for amplification is the ligase chain reaction (referred to as LCR), disclosed in Eur. Pat. Appl. Publ. No. 320,308 (specifically incorporated herein by reference in its entirety). In LCR, two complementary probe pairs are

prepared, and in the presence of the target sequence, each pair will bind to opposite complementary strands of the target such that they abut. In the presence of a ligase, the two probe pairs will link to form a single unit. By temperature cycling, as in PCR™, bound ligated units dissociate from the target and then serve as "target sequences" for
5 ligation of excess probe pairs. U.S. Patent No. 4,883,750, incorporated herein by reference in its entirety, describes an alternative method of amplification similar to LCR for binding probe pairs to a target sequence.

Qbeta Replicase, described in PCT Intl. Pat. Appl. Publ. No. PCT/US87/00880, incorporated herein by reference in its entirety, may also be used as
10 still another amplification method in the present invention. In this method, a replicative sequence of RNA that has a region complementary to that of a target is added to a sample in the presence of an RNA polymerase. The polymerase will copy the replicative sequence that can then be detected.

An isothermal amplification method, in which restriction endonucleases
15 and ligases are used to achieve the amplification of target molecules that contain nucleotide 5'-[α-thio]triphosphates in one strand of a restriction site (Walker *et al.*, 1992, incorporated herein by reference in its entirety), may also be useful in the amplification of nucleic acids in the present invention.

Strand Displacement Amplification (SDA) is another method of carrying
20 out isothermal amplification of nucleic acids which involves multiple rounds of strand displacement and synthesis, *i.e.* nick translation. A similar method, called Repair Chain Reaction (RCR) is another method of amplification which may be useful in the present invention and is involves annealing several probes throughout a region targeted for amplification, followed by a repair reaction in which only two of the four bases are
25 present. The other two bases can be added as biotinylated derivatives for easy detection. A similar approach is used in SDA.

Sequences can also be detected using a cyclic probe reaction (CPR). In CPR, a probe having a 3' and 5' sequences of non-target DNA and an internal or "middle" sequence of the target protein specific RNA is hybridized to DNA which is
30 present in a sample. Upon hybridization, the reaction is treated with RNaseH, and the products of the probe are identified as distinctive products by generating a signal that is released after digestion. The original template is annealed to another cycling probe and

the reaction is repeated. Thus, CPR involves amplifying a signal generated by hybridization of a probe to a target gene specific expressed nucleic acid.

Still other amplification methods described in Great Britain Pat. Appl. No. 2 202 328, and in PCT Intl. Pat. Appl. Publ. No. PCT/US89/01025, each of which
5 is incorporated herein by reference in its entirety, may be used in accordance with the present invention. In the former application, "modified" primers are used in a PCR-like, template and enzyme dependent synthesis. The primers may be modified by labeling with a capture moiety (*e.g.*, biotin) and/or a detector moiety (*e.g.*, enzyme). In the latter application, an excess of labeled probes is added to a sample. In the presence of the
10 target sequence, the probe binds and is cleaved catalytically. After cleavage, the target sequence is released intact to be bound by excess probe. Cleavage of the labeled probe signals the presence of the target sequence.

Other nucleic acid amplification procedures include transcription-based amplification systems (TAS) (Kwoh *et al.*, 1989; PCT Intl. Pat. Appl. Publ. No. WO
15 88/10315, incorporated herein by reference in its entirety), including nucleic acid sequence based amplification (NASBA) and 3SR. In NASBA, the nucleic acids can be prepared for amplification by standard phenol/chloroform extraction, heat denaturation of a sample, treatment with lysis buffer and minispin columns for isolation of DNA and RNA or guanidinium chloride extraction of RNA. These amplification techniques
20 involve annealing a primer that has sequences specific to the target sequence. Following polymerization, DNA/RNA hybrids are digested with RNase H while double stranded DNA molecules are heat-denatured again. In either case the single stranded DNA is made fully double stranded by addition of second target-specific primer, followed by polymerization. The double stranded DNA molecules are then multiply
25 transcribed by a polymerase such as T7 or SP6. In an isothermal cyclic reaction, the RNAs are reverse transcribed into DNA, and transcribed once again with a polymerase such as T7 or SP6. The resulting products, whether truncated or complete, indicate target-specific sequences.

Eur. Pat. Appl. Publ. No. 329,822, incorporated herein by reference in its
30 entirety, disclose a nucleic acid amplification process involving cyclically synthesizing single-stranded RNA ("ssRNA"), ssDNA, and double-stranded DNA (dsDNA), which may be used in accordance with the present invention. The ssRNA is a first template

for a first primer oligonucleotide, which is elongated by reverse transcriptase (RNA-dependent DNA polymerase). The RNA is then removed from resulting DNA:RNA duplex by the action of ribonuclease H (RNase H, an RNase specific for RNA in a duplex with either DNA or RNA). The resultant ssDNA is a second template
5 for a second primer, which also includes the sequences of an RNA polymerase promoter (exemplified by T7 RNA polymerase) 5' to its homology to its template. This primer is then extended by DNA polymerase (exemplified by the large "Klenow" fragment of *E. coli* DNA polymerase I), resulting as a double-stranded DNA ("dsDNA") molecule, having a sequence identical to that of the original RNA between the primers and having
10 additionally, at one end, a promoter sequence. This promoter sequence can be used by the appropriate RNA polymerase to make many RNA copies of the DNA. These copies can then re-enter the cycle leading to very swift amplification. With proper choice of enzymes, this amplification can be done isothermally without addition of enzymes at each cycle. Because of the cyclical nature of this process, the starting sequence can be
15 chosen to be in the form of either DNA or RNA.

PCT Intl. Pat. Appl. Publ. No. WO 89/06700, incorporated herein by reference in its entirety, disclose a nucleic acid sequence amplification scheme based on the hybridization of a promoter/primer sequence to a target single-stranded DNA ("ssDNA") followed by transcription of many RNA copies of the sequence. This
20 scheme is not cyclic; i.e. new templates are not produced from the resultant RNA transcripts. Other amplification methods include "RACE" (Frohman, 1990), and "one-sided PCR" (Ohara, 1989) which are well-known to those of skill in the art.

Methods based on ligation of two (or more) oligonucleotides in the presence of nucleic acid having the sequence of the resulting "di-oligonucleotide",
25 thereby amplifying the di-oligonucleotide (Wu and Dean, 1996, incorporated herein by reference in its entirety), may also be used in the amplification of DNA sequences of the present invention.

BIOLOGICAL FUNCTIONAL EQUIVALENTS

30 Modification and changes may be made in the structure of the polynucleotides and polypeptides of the present invention and still obtain a functional molecule that encodes a polypeptide with desirable characteristics. As mentioned

above, it is often desirable to introduce one or more mutations into a specific polynucleotide sequence. In certain circumstances, the resulting encoded polypeptide sequence is altered by this mutation, or in other cases, the sequence of the polypeptide is unchanged by one or more mutations in the encoding polynucleotide.

5 When it is desirable to alter the amino acid sequence of a polypeptide to create an equivalent, or even an improved, second-generation molecule, the amino acid changes may be achieved by changing one or more of the codons of the encoding DNA sequence, according to Table 1.

10 For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid sequence substitutions can be made in a protein sequence, and, of course, its underlying DNA
15 coding sequence, and nevertheless obtain a protein with like properties. It is thus contemplated by the inventors that various changes may be made in the peptide sequences of the disclosed compositions, or corresponding DNA sequences which encode said peptides without appreciable loss of their biological utility or activity.

TABLE 1

Amino Acids			Codons						
Alanine	Ala	A	GCA	GCC	GCG	GCU			
Cysteine	Cys	C	UGC	UGU					
Aspartic acid	Asp	D	GAC	GAU					
Glutamic acid	Glu	E	GAA	GAG					
Phenylalanine	Phe	F	UUC	UUU					
Glycine	Gly	G	GGA	GGC	GGG	GGU			
Histidine	His	H	CAC	CAU					
Isoleucine	Ile	I	AUA	AUC	AUU				
Lysine	Lys	K	AAA	AAG					
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	CUU	
Methionine	Met	M	AUG						
Asparagine	Asn	N	AAC	AAU					
Proline	Pro	P	CCA	CCC	CCG	CCU			
Glutamine	Gln	Q	CAA	CAG					
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGU	
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	UCU	
Threonine	Thr	T	ACA	ACC	ACG	ACU			
Valine	Val	V	GUA	GUC	GUG	GUU			
Tryptophan	Trp	W	UGG						
Tyrosine	Tyr	Y	UAC	UAU					

In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte and Doolittle, 1982, incorporated herein by reference). It is accepted that the relative hydropathic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics (Kyte and Doolittle, 1982). These values are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (−0.4); threonine (−0.7); serine (−0.8);

tryptophan (−0.9); tyrosine (−1.3); proline (−1.6); histidine (−3.2); glutamate (−3.5); glutamine (−3.5); aspartate (−3.5); asparagine (−3.5); lysine (−3.9); and arginine (−4.5).

It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydropathic index or score and still result in a protein with
5 similar biological activity, *i.e.* still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydropathic indices are within ± 2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred. It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U. S. Patent
10 4,554,101 (specifically incorporated herein by reference in its entirety), states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

As detailed in U. S. Patent 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate
15 (+3.0 \pm 1); glutamate (+3.0 \pm 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (−0.4); proline (−0.5 \pm 1); alanine (−0.5); histidine (−0.5); cysteine (−1.0); methionine (−1.3); valine (−1.5); leucine (−1.8); isoleucine (−1.8); tyrosine (−2.3); phenylalanine (−2.5); tryptophan (−3.4). It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a
20 biologically equivalent, and in particular, an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within ± 2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

As outlined above, amino acid substitutions are generally therefore based
25 on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions that take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

30 In addition, any polynucleotide may be further modified to increase stability *in vivo*. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends; the use of phosphorothioate or 2' O-methyl

rather than phosphodiesterase linkages in the backbone; and/or the inclusion of nontraditional bases such as inosine, queosine and wybutosine, as well as acetyl-methyl-, thio- and other modified forms of adenine, cytidine, guanine, thymine and uridine.

5

IN VIVO POLYNUCLEOTIDE DELIVERY TECHNIQUES

In additional embodiments, genetic constructs comprising one or more of the polynucleotides of the invention are introduced into cells *in vivo*. This may be achieved using any of a variety of well known approaches, several of which are outlined
10 below for the purpose of illustration.

1. ADENOVIRUS

One of the preferred methods for *in vivo* delivery of one or more nucleic acid sequences involves the use of an adenovirus expression vector. "Adenovirus expression vector" is meant to include those constructs containing adenovirus sequences
15 sufficient to (a) support packaging of the construct and (b) to express a polynucleotide that has been cloned therein in a sense or antisense orientation. Of course, in the context of an antisense construct, expression does not require that the gene product be synthesized.

The expression vector comprises a genetically engineered form of an
20 adenovirus. Knowledge of the genetic organization of adenovirus, a 36 kb, linear, double-stranded DNA virus, allows substitution of large pieces of adenoviral DNA with foreign sequences up to 7 kb (Grunhaus and Horwitz, 1992). In contrast to retrovirus, the adenoviral infection of host cells does not result in chromosomal integration because adenoviral DNA can replicate in an episomal manner without potential genotoxicity.
25 Also, adenoviruses are structurally stable, and no genome rearrangement has been detected after extensive amplification. Adenovirus can infect virtually all epithelial cells regardless of their cell cycle stage. So far, adenoviral infection appears to be linked only to mild disease such as acute respiratory disease in humans.

Adenovirus is particularly suitable for use as a gene transfer vector
30 because of its mid-sized genome, ease of manipulation, high titer, wide target-cell range and high infectivity. Both ends of the viral genome contain 100-200 base pair inverted repeats (ITRs), which are *cis* elements necessary for viral DNA replication and

packaging. The early (E) and late (L) regions of the genome contain different transcription units that are divided by the onset of viral DNA replication. The E1 region (E1A and E1B) encodes proteins responsible for the regulation of transcription of the viral genome and a few cellular genes. The expression of the E2 region (E2A and E2B) results in the synthesis of the proteins for viral DNA replication. These proteins are involved in DNA replication, late gene expression and host cell shut-off (Renan, 1990). The products of the late genes, including the majority of the viral capsid proteins, are expressed only after significant processing of a single primary transcript issued by the major late promoter (MLP). The MLP, (located at 16.8 m.u.) is particularly efficient during the late phase of infection, and all the mRNA's issued from this promoter possess a 5'-tripartite leader (TPL) sequence which makes them preferred mRNA's for translation.

In a current system, recombinant adenovirus is generated from homologous recombination between shuttle vector and provirus vector. Due to the possible recombination between two proviral vectors, wild-type adenovirus may be generated from this process. Therefore, it is critical to isolate a single clone of virus from an individual plaque and examine its genomic structure.

Generation and propagation of the current adenovirus vectors, which are replication deficient, depend on a unique helper cell line, designated 293, which was transformed from human embryonic kidney cells by Ad5 DNA fragments and constitutively expresses E1 proteins (Graham *et al.*, 1977). Since the E3 region is dispensable from the adenovirus genome (Jones and Shenk, 1978), the current adenovirus vectors, with the help of 293 cells, carry foreign DNA in either the E1, the D3 or both regions (Graham and Prevec, 1991). In nature, adenovirus can package approximately 105% of the wild-type genome (Ghosh-Choudhury *et al.*, 1987), providing capacity for about 2 extra kB of DNA. Combined with the approximately 5.5 kB of DNA that is replaceable in the E1 and E3 regions, the maximum capacity of the current adenovirus vector is under 7.5 kB, or about 15% of the total length of the vector. More than 80% of the adenovirus viral genome remains in the vector backbone and is the source of vector-borne cytotoxicity. Also, the replication deficiency of the E1-deleted virus is incomplete. For example, leakage of viral gene expression has been

observed with the currently available vectors at high multiplicities of infection (MOI) (Mulligan, 1993).

Helper cell lines may be derived from human cells such as human embryonic kidney cells, muscle cells, hematopoietic cells or other human embryonic mesenchymal or epithelial cells. Alternatively, the helper cells may be derived from the cells of other mammalian species that are permissive for human adenovirus. Such cells include, *e.g.*, Vero cells or other monkey embryonic mesenchymal or epithelial cells. As stated above, the currently preferred helper cell line is 293.

Recently, Racher *et al.* (1995) disclosed improved methods for culturing 293 cells and propagating adenovirus. In one format, natural cell aggregates are grown by inoculating individual cells into 1 liter siliconized spinner flasks (Techne, Cambridge, UK) containing 100-200 ml of medium. Following stirring at 40 rpm, the cell viability is estimated with trypan blue. In another format, Fibra-Cel microcarriers (Bibby Sterlin, Stone, UK) (5 g/l) is employed as follows. A cell inoculum, resuspended in 5 ml of medium, is added to the carrier (50 ml) in a 250 ml Erlenmeyer flask and left stationary, with occasional agitation, for 1 to 4 h. The medium is then replaced with 50 ml of fresh medium and shaking initiated. For virus production, cells are allowed to grow to about 80% confluence, after which time the medium is replaced (to 25% of the final volume) and adenovirus added at an MOI of 0.05. Cultures are left stationary overnight, following which the volume is increased to 100% and shaking commenced for another 72 h.

Other than the requirement that the adenovirus vector be replication defective, or at least conditionally defective, the nature of the adenovirus vector is not believed to be crucial to the successful practice of the invention. The adenovirus may be of any of the 42 different known serotypes or subgroups A-F. Adenovirus type 5 of subgroup C is the preferred starting material in order to obtain a conditional replication-defective adenovirus vector for use in the present invention, since Adenovirus type 5 is a human adenovirus about which a great deal of biochemical and genetic information is known, and it has historically been used for most constructions employing adenovirus as a vector.

As stated above, the typical vector according to the present invention is replication defective and will not have an adenovirus E1 region. Thus, it will be most

convenient to introduce the polynucleotide encoding the gene of interest at the position from which the E1-coding sequences have been removed. However, the position of insertion of the construct within the adenovirus sequences is not critical to the invention. The polynucleotide encoding the gene of interest may also be inserted in lieu
5 of the deleted E3 region in E3 replacement vectors as described by Karlsson *et al.* (1986) or in the E4 region where a helper cell line or helper virus complements the E4 defect.

Adenovirus is easy to grow and manipulate and exhibits broad host range *in vitro* and *in vivo*. This group of viruses can be obtained in high titers, *e.g.*, 10^9 - 10^{11}
10 plaque-forming units per ml, and they are highly infective. The life cycle of adenovirus does not require integration into the host cell genome. The foreign genes delivered by adenovirus vectors are episomal and, therefore, have low genotoxicity to host cells. No side effects have been reported in studies of vaccination with wild-type adenovirus (Couch *et al.*, 1963; Top *et al.*, 1971), demonstrating their safety and therapeutic
15 potential as *in vivo* gene transfer vectors.

Adenovirus vectors have been used in eukaryotic gene expression (Levrero *et al.*, 1991; Gomez-Foix *et al.*, 1992) and vaccine development (Grunhaus and Horwitz, 1992; Graham and Prevec, 1992). Recently, animal studies suggested that recombinant adenovirus could be used for gene therapy (Stratford-Perricaudet and
20 Perricaudet, 1991; Stratford-Perricaudet *et al.*, 1990; Rich *et al.*, 1993). Studies in administering recombinant adenovirus to different tissues include trachea instillation (Rosenfeld *et al.*, 1991; Rosenfeld *et al.*, 1992), muscle injection (Ragot *et al.*, 1993), peripheral intravenous injections (Herz and Gerard, 1993) and stereotactic inoculation into the brain (Le Gal La Salle *et al.*, 1993).

25 2. RETROVIRUSES

The retroviruses are a group of single-stranded RNA viruses characterized by an ability to convert their RNA to double-stranded DNA in infected cells by a process of reverse-transcription (Coffin, 1990). The resulting DNA then stably integrates into cellular chromosomes as a provirus and directs synthesis of viral
30 proteins. The integration results in the retention of the viral gene sequences in the recipient cell and its descendants. The retroviral genome contains three genes, *gag*, *pol*, and *env* that code for capsid proteins, polymerase enzyme, and envelope components,

respectively. A sequence found upstream from the gag gene contains a signal for packaging of the genome into virions. Two long terminal repeat (LTR) sequences are present at the 5' and 3' ends of the viral genome. These contain strong promoter and enhancer sequences and are also required for integration in the host cell genome
5 (Coffin, 1990).

In order to construct a retroviral vector, a nucleic acid encoding one or more oligonucleotide or polynucleotide sequences of interest is inserted into the viral genome in the place of certain viral sequences to produce a virus that is replication-defective. In order to produce virions, a packaging cell line containing the gag, pol, and
10 env genes but without the LTR and packaging components is constructed (Mann *et al.*, 1983). When a recombinant plasmid containing a cDNA, together with the retroviral LTR and packaging sequences is introduced into this cell line (by calcium phosphate precipitation for example), the packaging sequence allows the RNA transcript of the recombinant plasmid to be packaged into viral particles, which are then secreted into the
15 culture media (Nicolas and Rubenstein, 1988; Temin, 1986; Mann *et al.*, 1983). The media containing the recombinant retroviruses is then collected, optionally concentrated, and used for gene transfer. Retroviral vectors are able to infect a broad variety of cell types. However, integration and stable expression require the division of host cells (Paskind *et al.*, 1975).

20 A novel approach designed to allow specific targeting of retrovirus vectors was recently developed based on the chemical modification of a retrovirus by the chemical addition of lactose residues to the viral envelope. This modification could permit the specific infection of hepatocytes *via* sialoglycoprotein receptors.

A different approach to targeting of recombinant retroviruses was
25 designed in which biotinylated antibodies against a retroviral envelope protein and against a specific cell receptor were used. The antibodies were coupled *via* the biotin components by using streptavidin (Roux *et al.*, 1989). Using antibodies against major histocompatibility complex class I and class II antigens, they demonstrated the infection of a variety of human cells that bore those surface antigens with an ecotropic virus *in*
30 *vitro* (Roux *et al.*, 1989).

3. ADENO-ASSOCIATED VIRUSES

AAV (Ridgeway, 1988; Hermonat and Muzyczka, 1984) is a parovirus, discovered as a contamination of adenoviral stocks. It is a ubiquitous virus (antibodies are present in 85% of the US human population) that has not been linked to any disease. It is also classified as a dependovirus, because its replications is dependent on the presence of a helper virus, such as adenovirus. Five serotypes have been isolated, of which AAV-2 is the best characterized. AAV has a single-stranded linear DNA that is encapsidated into capsid proteins VP1, VP2 and VP3 to form an icosahedral virion of 20 to 24 nm in diameter (Muzyczka and McLaughlin, 1988).

The AAV DNA is approximately 4.7 kilobases long. It contains two open reading frames and is flanked by two ITRs (FIG. 2). There are two major genes in the AAV genome: *rep* and *cap*. The *rep* gene codes for proteins responsible for viral replications, whereas *cap* codes for capsid protein VP1-3. Each ITR forms a T-shaped hairpin structure. These terminal repeats are the only essential *cis* components of the AAV for chromosomal integration. Therefore, the AAV can be used as a vector with all viral coding sequences removed and replaced by the cassette of genes for delivery. Three viral promoters have been identified and named p5, p19, and p40, according to their map position. Transcription from p5 and p19 results in production of rep proteins, and transcription from p40 produces the capsid proteins (Hermonat and Muzyczka, 1984).

There are several factors that prompted researchers to study the possibility of using rAAV as an expression vector. One is that the requirements for delivering a gene to integrate into the host chromosome are surprisingly few. It is necessary to have the 145-bp ITRs, which are only 6% of the AAV genome. This leaves room in the vector to assemble a 4.5-kb DNA insertion. While this carrying capacity may prevent the AAV from delivering large genes, it is amply suited for delivering the antisense constructs of the present invention.

AAV is also a good choice of delivery vehicles due to its safety. There is a relatively complicated rescue mechanism: not only wild type adenovirus but also AAV genes are required to mobilize rAAV. Likewise, AAV is not pathogenic and not associated with any disease. The removal of viral coding sequences minimizes immune

reactions to viral gene expression, and therefore, rAAV does not evoke an inflammatory response.

4. OTHER VIRAL VECTORS AS EXPRESSION CONSTRUCTS

Other viral vectors may be employed as expression constructs in the present invention for the delivery of oligonucleotide or polynucleotide sequences to a host cell. Vectors derived from viruses such as vaccinia virus (Ridgeway, 1988; Coupar *et al.*, 1988), lentiviruses, polio viruses and herpes viruses may be employed. They offer several attractive features for various mammalian cells (Friedmann, 1989; Ridgeway, 1988; Coupar *et al.*, 1988; Horwich *et al.*, 1990).

With the recent recognition of defective hepatitis B viruses, new insight was gained into the structure-function relationship of different viral sequences. *In vitro* studies showed that the virus could retain the ability for helper-dependent packaging and reverse transcription despite the deletion of up to 80% of its genome (Horwich *et al.*, 1990). This suggested that large portions of the genome could be replaced with foreign genetic material. The hepatotropism and persistence (integration) were particularly attractive properties for liver-directed gene transfer. Chang *et al.* (1991) introduced the chloramphenicol acetyltransferase (CAT) gene into duck hepatitis B virus genome in the place of the polymerase, surface, and pre-surface coding sequences. It was cotransfected with wild-type virus into an avian hepatoma cell line. Culture media containing high titers of the recombinant virus were used to infect primary duckling hepatocytes. Stable CAT gene expression was detected for at least 24 days after transfection (Chang *et al.*, 1991).

5. NON-VIRAL VECTORS

In order to effect expression of the oligonucleotide or polynucleotide sequences of the present invention, the expression construct must be delivered into a cell. This delivery may be accomplished *in vitro*, as in laboratory procedures for transforming cells lines, or *in vivo* or *ex vivo*, as in the treatment of certain disease states. As described above, one preferred mechanism for delivery is *via* viral infection where the expression construct is encapsulated in an infectious viral particle.

Once the expression construct has been delivered into the cell the nucleic acid encoding the desired oligonucleotide or polynucleotide sequences may be positioned and expressed at different sites. In certain embodiments, the nucleic acid

encoding the construct may be stably integrated into the genome of the cell. This integration may be in the specific location and orientation *via* homologous recombination (gene replacement) or it may be integrated in a random, non-specific location (gene augmentation). In yet further embodiments, the nucleic acid may be stably maintained in the cell as a separate, episomal segment of DNA. Such nucleic acid segments or "episomes" encode sequences sufficient to permit maintenance and replication independent of or in synchronization with the host cell cycle. How the expression construct is delivered to a cell and where in the cell the nucleic acid remains is dependent on the type of expression construct employed.

10 In certain embodiments of the invention, the expression construct comprising one or more oligonucleotide or polynucleotide sequences may simply consist of naked recombinant DNA or plasmids. Transfer of the construct may be performed by any of the methods mentioned above which physically or chemically permeabilize the cell membrane. This is particularly applicable for transfer *in vitro* but it may be applied to *in vivo* use as well. Dubensky *et al.* (1984) successfully injected polyomavirus DNA in the form of calcium phosphate precipitates into liver and spleen of adult and newborn mice demonstrating active viral replication and acute infection. Benvenisty and Reshef (1986) also demonstrated that direct intraperitoneal injection of calcium phosphate-precipitated plasmids results in expression of the transfected genes. 15 It is envisioned that DNA encoding a gene of interest may also be transferred in a similar manner *in vivo* and express the gene product.

Another embodiment of the invention for transferring a naked DNA expression construct into cells may involve particle bombardment. This method depends on the ability to accelerate DNA-coated microprojectiles to a high velocity 25 allowing them to pierce cell membranes and enter cells without killing them (Klein *et al.*, 1987). Several devices for accelerating small particles have been developed. One such device relies on a high voltage discharge to generate an electrical current, which in turn provides the motive force (Yang *et al.*, 1990). The microprojectiles used have consisted of biologically inert substances such as tungsten or gold beads.

30 Selected organs including the liver, skin, and muscle tissue of rats and mice have been bombarded *in vivo* (Yang *et al.*, 1990; Zelenin *et al.*, 1991). This may require surgical exposure of the tissue or cells, to eliminate any intervening tissue

between the gun and the target organ, *i.e.* *ex vivo* treatment. Again, DNA encoding a particular gene may be delivered *via* this method and still be incorporated by the present invention.

5 ANTISENSE OLIGONUCLEOTIDES

The end result of the flow of genetic information is the synthesis of protein. DNA is transcribed by polymerases into messenger RNA and translated on the ribosome to yield a folded, functional protein. Thus there are several steps along the route where protein synthesis can be inhibited. The native DNA segment coding for a polypeptide described herein, as all such mammalian DNA strands, has two strands: a sense strand and an antisense strand held together by hydrogen bonding. The messenger RNA coding for polypeptide has the same nucleotide sequence as the sense DNA strand except that the DNA thymidine is replaced by uridine. Thus, synthetic antisense nucleotide sequences will bind to a mRNA and inhibit expression of the protein encoded by that mRNA.

The targeting of antisense oligonucleotides to mRNA is thus one mechanism to shut down protein synthesis, and, consequently, represents a powerful and targeted therapeutic approach. For example, the synthesis of polygalacturonase and the muscarine type 2 acetylcholine receptor are inhibited by antisense oligonucleotides directed to their respective mRNA sequences (U. S. Patent 5,739,119 and U. S. Patent 5,759,829, each specifically incorporated herein by reference in its entirety). Further, examples of antisense inhibition have been demonstrated with the nuclear protein cyclin, the multiple drug resistance gene (MDG1), ICAM-1, E-selectin, STK-1, striatal GABA_A receptor and human EGF (Jaskulski *et al.*, 1988; Vasanthakumar and Ahmed, 1989; Peris *et al.*, 1998; U. S. Patent 5,801,154; U. S. Patent 5,789,573; U. S. Patent 5,718,709 and U. S. Patent 5,610,288, each specifically incorporated herein by reference in its entirety). Antisense constructs have also been described that inhibit and can be used to treat a variety of abnormal cellular proliferations, *e.g.* cancer (U. S. Patent 5,747,470; U. S. Patent 5,591,317 and U. S. Patent 5,783,683, each specifically incorporated herein by reference in its entirety).

Therefore, in exemplary embodiments, the invention provides oligonucleotide sequences that comprise all, or a portion of, any sequence that is

capable of specifically binding to polynucleotide sequence described herein, or a complement thereof. In one embodiment, the antisense oligonucleotides comprise DNA or derivatives thereof. In another embodiment, the oligonucleotides comprise RNA or derivatives thereof. In a third embodiment, the oligonucleotides are modified DNAs comprising a phosphorothioated modified backbone. In a fourth embodiment, the oligonucleotide sequences comprise peptide nucleic acids or derivatives thereof. In each case, preferred compositions comprise a sequence region that is complementary, and more preferably substantially-complementary, and even more preferably, completely complementary to one or more portions of polynucleotides disclosed herein.

Selection of antisense compositions specific for a given gene sequence is based upon analysis of the chosen target sequence (*i.e.* in these illustrative examples the rat and human sequences) and determination of secondary structure, T_m , binding energy, relative stability, and antisense compositions were selected based upon their relative inability to form dimers, hairpins, or other secondary structures that would reduce or prohibit specific binding to the target mRNA in a host cell.

Highly preferred target regions of the mRNA, are those which are at or near the AUG translation initiation codon, and those sequences which were substantially complementary to 5' regions of the mRNA. These secondary structure analyses and target site selection considerations were performed using v.4 of the OLIGO primer analysis software (Rychlik, 1997) and the BLASTN 2.0.5 algorithm software (Altschul *et al.*, 1997).

The use of an antisense delivery method employing a short peptide vector, termed MPG (27 residues), is also contemplated. The MPG peptide contains a hydrophobic domain derived from the fusion sequence of HIV gp41 and a hydrophilic domain from the nuclear localization sequence of SV40 T-antigen (Morris *et al.*, 1997). It has been demonstrated that several molecules of the MPG peptide coat the antisense oligonucleotides and can be delivered into cultured mammalian cells in less than 1 hour with relatively high efficiency (90%). Further, the interaction with MPG strongly increases both the stability of the oligonucleotide to nuclease and the ability to cross the plasma membrane (Morris *et al.*, 1997).

RIBOZYMES

Although proteins traditionally have been used for catalysis of nucleic acids, another class of macromolecules has emerged as useful in this endeavor. Ribozymes are RNA-protein complexes that cleave nucleic acids in a site-specific fashion. Ribozymes have specific catalytic domains that possess endonuclease activity (Kim and Cech, 1987; Gerlach *et al.*, 1987; Forster and Symons, 1987). For example, a large number of ribozymes accelerate phosphoester transfer reactions with a high degree of specificity, often cleaving only one of several phosphoesters in an oligonucleotide substrate (Cech *et al.*, 1981; Michel and Westhof, 1990; Reinhold-Hurek and Shub, 1992). This specificity has been attributed to the requirement that the substrate bind via specific base-pairing interactions to the internal guide sequence ("IGS") of the ribozyme prior to chemical reaction.

Ribozyme catalysis has primarily been observed as part of sequence-specific cleavage/ligation reactions involving nucleic acids (Joyce, 1989; Cech *et al.*, 1981). For example, U. S. Patent No. 5,354,855 (specifically incorporated herein by reference) reports that certain ribozymes can act as endonucleases with a sequence specificity greater than that of known ribonucleases and approaching that of the DNA restriction enzymes. Thus, sequence-specific ribozyme-mediated inhibition of gene expression may be particularly suited to therapeutic applications (Scanlon *et al.*, 1991; Sarver *et al.*, 1990). Recently, it was reported that ribozymes elicited genetic changes in some cells lines to which they were applied; the altered genes included the oncogenes H-ras, c-fos and genes of HIV. Most of this work involved the modification of a target mRNA, based on a specific mutant codon that is cleaved by a specific ribozyme.

Six basic varieties of naturally-occurring enzymatic RNAs are known presently. Each can catalyze the hydrolysis of RNA phosphodiester bonds *in trans* (and thus can cleave other RNA molecules) under physiological conditions. In general, enzymatic nucleic acids act by first binding to a target RNA. Such binding occurs through the target binding portion of a enzymatic nucleic acid which is held in close proximity to an enzymatic portion of the molecule that acts to cleave the target RNA. Thus, the enzymatic nucleic acid first recognizes and then binds a target RNA through complementary base-pairing, and once bound to the correct site, acts enzymatically to cut the target RNA. Strategic cleavage of such a target RNA will destroy its ability to

direct synthesis of an encoded protein. After an enzymatic nucleic acid has bound and cleaved its RNA target, it is released from that RNA to search for another target and can repeatedly bind and cleave new targets.

The enzymatic nature of a ribozyme is advantageous over many technologies, such as antisense technology (where a nucleic acid molecule simply binds to a nucleic acid target to block its translation) since the concentration of ribozyme necessary to affect a therapeutic treatment is lower than that of an antisense oligonucleotide. This advantage reflects the ability of the ribozyme to act enzymatically. Thus, a single ribozyme molecule is able to cleave many molecules of target RNA. In addition, the ribozyme is a highly specific inhibitor, with the specificity of inhibition depending not only on the base pairing mechanism of binding to the target RNA, but also on the mechanism of target RNA cleavage. Single mismatches, or base-substitutions, near the site of cleavage can completely eliminate catalytic activity of a ribozyme. Similar mismatches in antisense molecules do not prevent their action (Woolf *et al.*, 1992). Thus, the specificity of action of a ribozyme is greater than that of an antisense oligonucleotide binding the same RNA site.

The enzymatic nucleic acid molecule may be formed in a hammerhead, hairpin, a hepatitis δ virus, group I intron or RNaseP RNA (in association with an RNA guide sequence) or Neurospora VS RNA motif. Examples of hammerhead motifs are described by Rossi *et al.* (1992). Examples of hairpin motifs are described by Hampel *et al.* (Eur. Pat. Appl. Publ. No. EP 0360257), Hampel and Tritz (1989), Hampel *et al.* (1990) and U. S. Patent 5,631,359 (specifically incorporated herein by reference). An example of the hepatitis δ virus motif is described by Perrotta and Been (1992); an example of the RNaseP motif is described by Guerrier-Takada *et al.* (1983); Neurospora VS RNA ribozyme motif is described by Collins (Saville and Collins, 1990; Saville and Collins, 1991; Collins and Olive, 1993); and an example of the Group I intron is described in (U. S. Patent 4,987,071, specifically incorporated herein by reference). All that is important in an enzymatic nucleic acid molecule of this invention is that it has a specific substrate binding site which is complementary to one or more of the target gene RNA regions, and that it have nucleotide sequences within or surrounding that substrate binding site which impart an RNA cleaving activity to the molecule. Thus the ribozyme constructs need not be limited to specific motifs mentioned herein.

In certain embodiments, it may be important to produce enzymatic cleaving agents which exhibit a high degree of specificity for the RNA of a desired target, such as one of the sequences disclosed herein. The enzymatic nucleic acid molecule is preferably targeted to a highly conserved sequence region of a target mRNA. Such enzymatic nucleic acid molecules can be delivered exogenously to specific cells as required. Alternatively, the ribozymes can be expressed from DNA or RNA vectors that are delivered to specific cells.

Small enzymatic nucleic acid motifs (e.g., of the hammerhead or the hairpin structure) may also be used for exogenous delivery. The simple structure of these molecules increases the ability of the enzymatic nucleic acid to invade targeted regions of the mRNA structure. Alternatively, catalytic RNA molecules can be expressed within cells from eukaryotic promoters (e.g., Scanlon *et al.*, 1991; Kashani-Sabet *et al.*, 1992; Dropulic *et al.*, 1992; Weerasinghe *et al.*, 1991; Ojwang *et al.*, 1992; Chen *et al.*, 1992; Sarver *et al.*, 1990). Those skilled in the art realize that any ribozyme can be expressed in eukaryotic cells from the appropriate DNA vector. The activity of such ribozymes can be augmented by their release from the primary transcript by a second ribozyme (Int. Pat. Appl. Publ. No. WO 93/23569, and Int. Pat. Appl. Publ. No. WO 94/02595, both hereby incorporated by reference; Ohkawa *et al.*, 1992; Taira *et al.*, 1991; and Ventura *et al.*, 1993).

Ribozymes may be added directly, or can be complexed with cationic lipids, lipid complexes, packaged within liposomes, or otherwise delivered to target cells. The RNA or RNA complexes can be locally administered to relevant tissues *ex vivo*, or *in vivo* through injection, aerosol inhalation, infusion pump or stent, with or without their incorporation in biopolymers.

Ribozymes may be designed as described in Int. Pat. Appl. Publ. No. WO 93/23569 and Int. Pat. Appl. Publ. No. WO 94/02595, each specifically incorporated herein by reference) and synthesized to be tested *in vitro* and *in vivo*, as described. Such ribozymes can also be optimized for delivery. While specific examples are provided, those in the art will recognize that equivalent RNA targets in other species can be utilized when necessary.

Hammerhead or hairpin ribozymes may be individually analyzed by computer folding (Jaeger *et al.*, 1989) to assess whether the ribozyme sequences fold

into the appropriate secondary structure. Those ribozymes with unfavorable intramolecular interactions between the binding arms and the catalytic core are eliminated from consideration. Varying binding arm lengths can be chosen to optimize activity. Generally, at least 5 or so bases on each arm are able to bind to, or otherwise
5 interact with, the target RNA.

Ribozymes of the hammerhead or hairpin motif may be designed to anneal to various sites in the mRNA message, and can be chemically synthesized. The method of synthesis used follows the procedure for normal RNA synthesis as described in Usman *et al.* (1987) and in Scaringe *et al.* (1990) and makes use of common nucleic
10 acid protecting and coupling groups, such as dimethoxytrityl at the 5'-end, and phosphoramidites at the 3'-end. Average stepwise coupling yields are typically >98%. Hairpin ribozymes may be synthesized in two parts and annealed to reconstruct an active ribozyme (Chowrira and Burke, 1992). Ribozymes may be modified extensively to enhance stability by modification with nuclease resistant groups, for example, 2'-
15 amino, 2'-C-allyl, 2'-fluoro, 2'-O-methyl, 2'-H (for a review see *e.g.*, Usman and Cedergren, 1992). Ribozymes may be purified by gel electrophoresis using general methods or by high pressure liquid chromatography and resuspended in water.

Ribozyme activity can be optimized by altering the length of the ribozyme binding arms, or chemically synthesizing ribozymes with modifications that
20 prevent their degradation by serum ribonucleases (see *e.g.*, Int. Pat. Appl. Publ. No. WO 92/07065; Perrault *et al.*, 1990; Pieken *et al.*, 1991; Usman and Cedergren, 1992; Int. Pat. Appl. Publ. No. WO 93/15187; Int. Pat. Appl. Publ. No. WO 91/03162; Eur. Pat. Appl. Publ. No. 92110298.4; U. S. Patent 5,334,711; and Int. Pat. Appl. Publ. No. WO 94/13688, which describe various chemical modifications that can be made to the sugar
25 moieties of enzymatic RNA molecules), modifications which enhance their efficacy in cells, and removal of stem II bases to shorten RNA synthesis times and reduce chemical requirements.

Sullivan *et al.* (Int. Pat. Appl. Publ. No. WO 94/02595) describes the general methods for delivery of enzymatic RNA molecules. Ribozymes may be
30 administered to cells by a variety of methods known to those familiar to the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins, biodegradable

nanocapsules, and bioadhesive microspheres. For some indications, ribozymes may be directly delivered *ex vivo* to cells or tissues with or without the aforementioned vehicles. Alternatively, the RNA/vehicle combination may be locally delivered by direct inhalation, by direct injection or by use of a catheter, infusion pump or stent. Other routes of delivery include, but are not limited to, intravascular, intramuscular, subcutaneous or joint injection, aerosol inhalation, oral (tablet or pill form), topical, systemic, ocular, intraperitoneal and/or intrathecal delivery. More detailed descriptions of ribozyme delivery and administration are provided in Int. Pat. Appl. Publ. No. WO 94/02595 and Int. Pat. Appl. Publ. No. WO 93/23569, each specifically incorporated
10 herein by reference.

Another means of accumulating high concentrations of a ribozyme(s) within cells is to incorporate the ribozyme-encoding sequences into a DNA expression vector. Transcription of the ribozyme sequences are driven from a promoter for eukaryotic RNA polymerase I (pol I), RNA polymerase II (pol II), or RNA polymerase
15 III (pol III). Transcripts from pol II or pol III promoters will be expressed at high levels in all cells; the levels of a given pol II promoter in a given cell type will depend on the nature of the gene regulatory sequences (enhancers, silencers, *etc.*) present nearby. Prokaryotic RNA polymerase promoters may also be used, providing that the prokaryotic RNA polymerase enzyme is expressed in the appropriate cells (Elroy-Stein
20 and Moss, 1990; Gao and Huang, 1993; Lieber *et al.*, 1993; Zhou *et al.*, 1990). Ribozymes expressed from such promoters can function in mammalian cells (*e.g.* Kashani-Saber *et al.*, 1992; Ojwang *et al.*, 1992; Chen *et al.*, 1992; Yu *et al.*, 1993; L'Huillier *et al.*, 1992; Lisiewicz *et al.*, 1993). Such transcription units can be incorporated into a variety of vectors for introduction into mammalian cells, including
25 but not restricted to, plasmid DNA vectors, viral DNA vectors (such as adenovirus or adeno-associated vectors), or viral RNA vectors (such as retroviral, semliki forest virus, sindbis virus vectors).

Ribozymes may be used as diagnostic tools to examine genetic drift and mutations within diseased cells. They can also be used to assess levels of the target
30 RNA molecule. The close relationship between ribozyme activity and the structure of the target RNA allows the detection of mutations in any region of the molecule which alters the base-pairing and three-dimensional structure of the target RNA. By using

multiple ribozymes, one may map nucleotide changes which are important to RNA structure and function *in vitro*, as well as in cells and tissues. Cleavage of target RNAs with ribozymes may be used to inhibit gene expression and define the role (essentially) of specified gene products in the progression of disease. In this manner, other genetic
5 targets may be defined as important mediators of the disease. These studies will lead to better treatment of the disease progression by affording the possibility of combinational therapies (*e.g.*, multiple ribozymes targeted to different genes, ribozymes coupled with known small molecule inhibitors, or intermittent treatment with combinations of ribozymes and/or other chemical or biological molecules). Other *in vitro* uses of
10 ribozymes are well known in the art, and include detection of the presence of mRNA associated with an IL-5 related condition. Such RNA is detected by determining the presence of a cleavage product after treatment with a ribozyme using standard methodology.

15 PEPTIDE NUCLEIC ACIDS

In certain embodiments, the inventors contemplate the use of peptide nucleic acids (PNAs) in the practice of the methods of the invention. PNA is a DNA mimic in which the nucleobases are attached to a pseudopeptide backbone (Good and Nielsen, 1997). PNA is able to be utilized in a number methods that traditionally have
20 used RNA or DNA. Often PNA sequences perform better in techniques than the corresponding RNA or DNA sequences and have utilities that are not inherent to RNA or DNA. A review of PNA including methods of making, characteristics of, and methods of using, is provided by Corey (1997) and is incorporated herein by reference. As such, in certain embodiments, one may prepare PNA sequences that are
25 complementary to one or more portions of the ACE mRNA sequence, and such PNA compositions may be used to regulate, alter, decrease, or reduce the translation of ACE-specific mRNA, and thereby alter the level of ACE activity in a host cell to which such PNA compositions have been administered.

PNAs have 2-aminoethyl-glycine linkages replacing the normal
30 phosphodiester backbone of DNA (Nielsen *et al.*, 1991; Hanvey *et al.*, 1992; Hyrup and Nielsen, 1996; Neilsen, 1996). This chemistry has three important consequences: firstly, in contrast to DNA or phosphorothioate oligonucleotides, PNAs are neutral

molecules; secondly, PNAs are achiral, which avoids the need to develop a stereoselective synthesis; and thirdly, PNA synthesis uses standard Boc (Dueholm *et al.*, 1994) or Fmoc (Thomson *et al.*, 1995) protocols for solid-phase peptide synthesis, although other methods, including a modified Merrifield method, have been used
5 (Christensen *et al.*, 1995).

PNA monomers or ready-made oligomers are commercially available from PerSeptive Biosystems (Framingham, MA). PNA syntheses by either Boc or Fmoc protocols are straightforward using manual or automated protocols (Norton *et al.*, 1995). The manual protocol lends itself to the production of chemically modified PNAs
10 or the simultaneous synthesis of families of closely related PNAs.

As with peptide synthesis, the success of a particular PNA synthesis will depend on the properties of the chosen sequence. For example, while in theory PNAs can incorporate any combination of nucleotide bases, the presence of adjacent purines can lead to deletions of one or more residues in the product. In expectation of this
15 difficulty, it is suggested that, in producing PNAs with adjacent purines, one should repeat the coupling of residues likely to be added inefficiently. This should be followed by the purification of PNAs by reverse-phase high-pressure liquid chromatography (Norton *et al.*, 1995) providing yields and purity of product similar to those observed during the synthesis of peptides.

20 Modifications of PNAs for a given application may be accomplished by coupling amino acids during solid-phase synthesis or by attaching compounds that contain a carboxylic acid group to the exposed N-terminal amine. Alternatively, PNAs can be modified after synthesis by coupling to an introduced lysine or cysteine. The ease with which PNAs can be modified facilitates optimization for better solubility or
25 for specific functional requirements. Once synthesized, the identity of PNAs and their derivatives can be confirmed by mass spectrometry. Several studies have made and utilized modifications of PNAs (Norton *et al.*, 1995; Haaima *et al.*, 1996; Stetsenko *et al.*, 1996; Petersen *et al.*, 1995; Ulmann *et al.*, 1996; Koch *et al.*, 1995; Orum *et al.*, 1995; Footer *et al.*, 1996; Griffith *et al.*, 1995; Kremsky *et al.*, 1996; Pardridge *et al.*,
30 1995; Boffa *et al.*, 1995; Landsdorp *et al.*, 1996; Gambacorti-Passerini *et al.*, 1996; Armitage *et al.*, 1997; Seeger *et al.*, 1997; Ruskowski *et al.*, 1997). U.S. Patent No. 5,700,922 discusses PNA-DNA-PNA chimeric molecules and their uses in diagnostics,

modulating protein in organisms, and treatment of conditions susceptible to therapeutics.

In contrast to DNA and RNA, which contain negatively charged linkages, the PNA backbone is neutral. In spite of this dramatic alteration, PNAs
5 recognize complementary DNA and RNA by Watson-Crick pairing (Egholm *et al.*, 1993), validating the initial modeling by Nielsen *et al.* (1991). PNAs lack 3' to 5' polarity and can bind in either parallel or antiparallel fashion, with the antiparallel mode being preferred (Egholm *et al.*, 1993).

Hybridization of DNA oligonucleotides to DNA and RNA is destabilized
10 by electrostatic repulsion between the negatively charged phosphate backbones of the complementary strands. By contrast, the absence of charge repulsion in PNA-DNA or PNA-RNA duplexes increases the melting temperature (T_m) and reduces the dependence of T_m on the concentration of mono- or divalent cations (Nielsen *et al.*, 1991). The enhanced rate and affinity of hybridization are significant because they are responsible
15 for the surprising ability of PNAs to perform strand invasion of complementary sequences within relaxed double-stranded DNA. In addition, the efficient hybridization at inverted repeats suggests that PNAs can recognize secondary structure effectively within double-stranded DNA. Enhanced recognition also occurs with PNAs immobilized on surfaces, and Wang *et al.* have shown that support-bound PNAs can be
20 used to detect hybridization events (Wang *et al.*, 1996).

One might expect that tight binding of PNAs to complementary sequences would also increase binding to similar (but not identical) sequences, reducing the sequence specificity of PNA recognition. As with DNA hybridization, however, selective recognition can be achieved by balancing oligomer length and incubation
25 temperature. Moreover, selective hybridization of PNAs is encouraged by PNA-DNA hybridization being less tolerant of base mismatches than DNA-DNA hybridization. For example, a single mismatch within a 16 bp PNA-DNA duplex can reduce the T_m by up to 15°C (Egholm *et al.*, 1993). This high level of discrimination has allowed the development of several PNA-based strategies for the analysis of point mutations (Wang
30 *et al.*, 1996; Carlsson *et al.*, 1996; Thiede *et al.*, 1996; Webb and Hurskainen, 1996; Perry-O'Keefe *et al.*, 1996).

High-affinity binding provides clear advantages for molecular recognition and the development of new applications for PNAs. For example, 11-13 nucleotide PNAs inhibit the activity of telomerase, a ribonucleo-protein that extends telomere ends using an essential RNA template, while the analogous DNA oligomers do not (Norton *et al.*, 1996).

Neutral PNAs are more hydrophobic than analogous DNA oligomers, and this can lead to difficulty solubilizing them at neutral pH, especially if the PNAs have a high purine content or if they have the potential to form secondary structures. Their solubility can be enhanced by attaching one or more positive charges to the PNA termini (Nielsen *et al.*, 1991).

Findings by Allfrey and colleagues suggest that strand invasion will occur spontaneously at sequences within chromosomal DNA (Boffa *et al.*, 1995; Boffa *et al.*, 1996). These studies targeted PNAs to triplet repeats of the nucleotides CAG and used this recognition to purify transcriptionally active DNA (Boffa *et al.*, 1995) and to inhibit transcription (Boffa *et al.*, 1996). This result suggests that if PNAs can be delivered within cells then they will have the potential to be general sequence-specific regulators of gene expression. Studies and reviews concerning the use of PNAs as antisense and anti-gene agents include Nielsen *et al.* (1993b), Hanvey *et al.* (1992), and Good and Nielsen (1997). Koppelhus *et al.* (1997) have used PNAs to inhibit HIV-1 inverse transcription, showing that PNAs may be used for antiviral therapies.

Methods of characterizing the antisense binding properties of PNAs are discussed in Rose (1993) and Jensen *et al.* (1997). Rose uses capillary gel electrophoresis to determine binding of PNAs to their complementary oligonucleotide, measuring the relative binding kinetics and stoichiometry. Similar types of measurements were made by Jensen *et al.* using BIAcore™ technology.

Other applications of PNAs include use in DNA strand invasion (Nielsen *et al.*, 1991), antisense inhibition (Hanvey *et al.*, 1992), mutational analysis (Orum *et al.*, 1993), enhancers of transcription (Mollegaard *et al.*, 1994), nucleic acid purification (Orum *et al.*, 1995), isolation of transcriptionally active genes (Boffa *et al.*, 1995), blocking of transcription factor binding (Vickers *et al.*, 1995), genome cleavage (Veselkov *et al.*, 1996), biosensors (Wang *et al.*, 1996), *in situ* hybridization (Thisted *et al.*, 1996), and in an alternative to Southern blotting (Perry-O'Keefe, 1996).

POLYPEPTIDE COMPOSITIONS AND USES

The present invention, in other aspects, provides polypeptide compositions. Generally, a polypeptide of the invention will be an isolated polypeptide (or an epitope, variant, or active fragment thereof) derived from a mammalian species. Preferably, the polypeptide is encoded by a polynucleotide sequence disclosed herein or a sequence which hybridizes under moderately stringent conditions to a polynucleotide sequence disclosed herein. Alternatively, the polypeptide may be defined as a polypeptide which comprises a contiguous amino acid sequence from an amino acid sequence disclosed herein, or which polypeptide comprises an entire amino acid sequence disclosed herein.

Likewise, a polypeptide composition of the present invention is understood to comprise one or more polypeptides that are capable of eliciting antibodies that are immunologically reactive with one or more polypeptides encoded by one or more contiguous nucleic acid sequences contained in SEQ ID NO:1-48, 114-121, and 125-138, or to active fragments, or to variants thereof, or to one or more nucleic acid sequences which hybridize to one or more of these sequences under conditions of moderate to high stringency.

As used herein, an active fragment of a polypeptide includes a whole or a portion of a polypeptide which is modified by conventional techniques, *e.g.*, mutagenesis, or by addition, deletion, or substitution, but which active fragment exhibits substantially the same structure function, antigenicity, etc., as a polypeptide as described herein.

In certain illustrative embodiments, the polypeptides of the invention will comprise at least an immunogenic portion of a Chlamydia protein or a variant thereof, as described herein. Proteins that are Chlamydia proteins generally also react detectably within an immunoassay (such as an ELISA) with antisera from a patient with a Chlamydial infection. Polypeptides as described herein may be of any length. Additional sequences derived from the native protein and/or heterologous sequences may be present, and such sequences may (but need not) possess further immunogenic or antigenic properties.

An "immunogenic portion," as used herein is a portion of a protein that

is recognized (*i.e.*, specifically bound) by a B-cell and/or T-cell surface antigen receptor. Such immunogenic portions generally comprise at least 5 amino acid residues, more preferably at least 10, and still more preferably at least 20 amino acid residues of a Chlamydia protein or a variant thereof. Certain preferred immunogenic portions include
5 peptides in which an N-terminal leader sequence and/or transmembrane domain have been deleted. Other preferred immunogenic portions may contain a small N- and/or C-terminal deletion (*e.g.*, 1-30 amino acids, preferably 5-15 amino acids), relative to the mature protein.

Immunogenic portions may generally be identified using well known
10 techniques, such as those summarized in Paul, *Fundamental Immunology*, 3rd ed., 243-247 (Raven Press, 1993) and references cited therein. Such techniques include screening polypeptides for the ability to react with antigen-specific antibodies, antisera and/or T-cell lines or clones. As used herein, antisera and antibodies are "antigen-specific" if they specifically bind to an antigen (*i.e.*, they react with the protein in an
15 ELISA or other immunoassay, and do not react detectably with unrelated proteins). Such antisera and antibodies may be prepared as described herein, and using well known techniques. An immunogenic portion of a native Chlamydia protein is a portion that reacts with such antisera and/or T-cells at a level that is not substantially less than the reactivity of the full length polypeptide (*e.g.*, in an ELISA and/or T-cell reactivity
20 assay). Such immunogenic portions may react within such assays at a level that is similar to or greater than the reactivity of the full length polypeptide. Such screens may generally be performed using methods well known to those of ordinary skill in the art, such as those described in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. For example, a polypeptide may be immobilized on a
25 solid support and contacted with patient sera to allow binding of antibodies within the sera to the immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, ¹²⁵I-labeled Protein A.

As noted above, a composition may comprise a variant of a native Chlamydia protein. A polypeptide "variant," as used herein, is a polypeptide that differs
30 from a native Chlamydia protein in one or more substitutions, deletions, additions and/or insertions, such that the immunogenicity of the polypeptide is not substantially diminished. In other words, the ability of a variant to react with antigen-specific

antisera may be enhanced or unchanged, relative to the native protein, or may be diminished by less than 50%, and preferably less than 20%, relative to the native protein. Such variants may generally be identified by modifying one of the above polypeptide sequences and evaluating the reactivity of the modified polypeptide with
5 antigen-specific antibodies or antisera as described herein. Preferred variants include those in which one or more portions, such as an N-terminal leader sequence or transmembrane domain, have been removed. Other preferred variants include variants in which a small portion (*e.g.*, 1-30 amino acids, preferably 5-15 amino acids) has been removed from the N- and/or C-terminal of the mature protein.

10 Polypeptide variants encompassed by the present invention include those exhibiting at least about 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more identity (determined as described above) to the polypeptides disclosed herein.

Preferably, a variant contains conservative substitutions. A
15 "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. Amino acid substitutions may generally be made on the basis of similarity in polarity, charge, solubility, hydrophobicity,
20 hydrophilicity and/or the amphipathic nature of the residues. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine and valine; glycine and alanine; asparagine and glutamine; and serine, threonine, phenylalanine and tyrosine.
25 Other groups of amino acids that may represent conservative changes include: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. A variant may also, or alternatively, contain nonconservative changes. In a preferred embodiment, variant polypeptides differ from a native sequence by substitution, deletion or addition of five amino acids or fewer.
30 Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the immunogenicity, secondary structure and hydrophobic nature of the polypeptide.

As noted above, polypeptides may comprise a signal (or leader) sequence at the N-terminal end of the protein, which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

Polypeptides may be prepared using any of a variety of well known techniques. Recombinant polypeptides encoded by DNA sequences as described above may be readily prepared from the DNA sequences using any of a variety of expression vectors known to those of ordinary skill in the art. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast, and higher eukaryotic cells, such as mammalian cells and plant cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line such as COS or CHO. Supernatants from suitable host/vector systems which secrete recombinant protein or polypeptide into culture media may be first concentrated using a commercially available filter. Following concentration, the concentrate may be applied to a suitable purification matrix such as an affinity matrix or an ion exchange resin. Finally, one or more reverse phase HPLC steps can be employed to further purify a recombinant polypeptide.

Portions and other variants having less than about 100 amino acids, and generally less than about 50 amino acids, may also be generated by synthetic means, using techniques well known to those of ordinary skill in the art. For example, such polypeptides may be synthesized using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. See Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963. Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, CA), and may be operated according to the manufacturer's instructions.

Within certain specific embodiments, a polypeptide may be a fusion protein that comprises multiple polypeptides as described herein, or that comprises at

least one polypeptide as described herein and an unrelated sequence, such as a known Chlamydia protein. A fusion partner may, for example, assist in providing T helper epitopes (an immunological fusion partner), preferably T helper epitopes recognized by humans, or may assist in expressing the protein (an expression enhancer) at higher yields than the native recombinant protein. Certain preferred fusion partners are both immunological and expression enhancing fusion partners. Other fusion partners may be selected so as to increase the solubility of the protein or to enable the protein to be targeted to desired intracellular compartments. Still further fusion partners include affinity tags, which facilitate purification of the protein.

Fusion proteins may generally be prepared using standard techniques, including chemical conjugation. Preferably, a fusion protein is expressed as a recombinant protein, allowing the production of increased levels, relative to a non-fused protein, in an expression system. Briefly, DNA sequences encoding the polypeptide components may be assembled separately, and ligated into an appropriate expression vector. The 3' end of the DNA sequence encoding one polypeptide component is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide component so that the reading frames of the sequences are in phase. This permits translation into a single fusion protein that retains the biological activity of both component polypeptides.

A peptide linker sequence may be employed to separate the first and second polypeptide components by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S.

Patent No. 4,751,180. The linker sequence may generally be from 1 to about 50 amino acids in length. Linker sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

5 The ligated DNA sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of DNA are located only 5' to the DNA sequence encoding the first polypeptides. Similarly, stop codons required to end translation and transcription termination signals are only present 3' to the DNA sequence encoding the
10 second polypeptide.

Fusion proteins are also provided. Such proteins comprise a polypeptide as described herein together with an unrelated immunogenic protein. Preferably the immunogenic protein is capable of eliciting a recall response. Examples of such proteins include tetanus, tuberculosis and hepatitis proteins (*see, for example, Stoute et al. New Engl. J. Med.*, 336:86-91, 1997).
15

Within preferred embodiments, an immunological fusion partner is derived from protein D, a surface protein of the gram-negative bacterium *Haemophilus influenza B* (WO 91/18926). Preferably, a protein D derivative comprises approximately the first third of the protein (*e.g.*, the first N-terminal 100-110 amino
20 acids), and a protein D derivative may be lipidated. Within certain preferred embodiments, the first 109 residues of a Lipoprotein D fusion partner is included on the N-terminus to provide the polypeptide with additional exogenous T-cell epitopes and to increase the expression level in *E. coli* (thus functioning as an expression enhancer). The lipid tail ensures optimal presentation of the antigen to antigen presenting cells.
25 Other fusion partners include the non-structural protein from influenzae virus, NS1 (hemagglutinin). Typically, the N-terminal 81 amino acids are used, although different fragments that include T-helper epitopes may be used.

In another embodiment, the immunological fusion partner is the protein known as LYTA, or a portion thereof (preferably a C-terminal portion). LYTA is
30 derived from *Streptococcus pneumoniae*, which synthesizes an N-acetyl-L-alanine amidase known as amidase LYTA (encoded by the *LytA* gene; *Gene* 43:265-292, 1986). LYTA is an autolysin that specifically degrades certain bonds in the peptidoglycan

backbone. The C-terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LYTA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LYTA
5 fragment at the amino terminus has been described (*see Biotechnology 10:795-798, 1992*). Within a preferred embodiment, a repeat portion of LYTA may be incorporated into a fusion protein. A repeat portion is found in the C-terminal region starting at residue 178. A particularly preferred repeat portion incorporates residues 188-305.

In general, polypeptides (including fusion proteins) and polynucleotides
10 as described herein are isolated. An "isolated" polypeptide or polynucleotide is one that is removed from its original environment. For example, a naturally-occurring protein is isolated if it is separated from some or all of the coexisting materials in the natural system. Preferably, such polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. A polynucleotide is
15 considered to be isolated if, for example, it is cloned into a vector that is not a part of the natural environment.

ILLUSTRATIVE THERAPEUTIC COMPOSITIONS AND USES

In another aspect, the present invention provides methods for using one
20 or more of the above polypeptides or fusion proteins (or polynucleotides encoding such polypeptides or fusion proteins) to induce protective immunity against Chlamydial infection in a patient. As used herein, a "patient" refers to any warm-blooded animal, preferably a human. A patient may be afflicted with a disease, or may be free of detectable disease and/or infection. In other words, protective immunity may be
25 induced to prevent or treat Chlamydial infection.

In this aspect, the polypeptide, fusion protein or polynucleotide molecule is generally present within a pharmaceutical composition or a vaccine. Pharmaceutical compositions may comprise one or more polypeptides, each of which may contain one or more of the above sequences (or variants thereof), and a physiologically acceptable
30 carrier. Vaccines may comprise one or more of the above polypeptides and an immunostimulant, such as an adjuvant or a liposome (into which the polypeptide is incorporated). Such pharmaceutical compositions and vaccines may also contain other

Chlamydia antigens, either incorporated into a combination polypeptide or present within a separate polypeptide.

Alternatively, a vaccine may contain polynucleotides encoding one or more polypeptides or fusion proteins as described above, such that the polypeptide is generated *in situ*. In such vaccines, the polynucleotides may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacterial and viral expression systems. Appropriate nucleic acid expression systems contain the necessary polynucleotide sequences for expression in the patient (such as a suitable promoter and terminating signal). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an immunogenic portion of the polypeptide on its cell surface. In a preferred embodiment, the polynucleotides may be introduced using a viral expression system (e.g., vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective) virus. Techniques for incorporating polynucleotides into such expression systems are well known to those of ordinary skill in the art. The polynucleotides may also be administered as "naked" plasmid vectors as described, for example, in Ulmer et al., *Science* 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. Techniques for incorporating DNA into such vectors are well known to those of ordinary skill in the art. A retroviral vector may additionally transfer or incorporate a gene for a selectable marker (to aid in the identification or selection of transduced cells) and/or a targeting moiety, such as a gene that encodes a ligand for a receptor on a specific target cell, to render the vector target specific. Targeting may also be accomplished using an antibody, by methods known to those of ordinary skill in the art.

Other formulations for therapeutic purposes include colloidal dispersion systems, such as macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. A preferred colloidal system for use as a delivery vehicle *in vitro* and *in vivo* is a liposome (*i.e.*, an artificial membrane vesicle). The uptake of naked polynucleotides may be increased by incorporating the polynucleotides into and/or onto biodegradable beads, which are efficiently transported into the cells. The preparation and use of such systems is well known in the art.

In a related aspect, a polynucleotide vaccine as described above may be administered simultaneously with or sequentially to either a polypeptide of the present invention or a known *Chlamydia* antigen. For example, administration of polynucleotides encoding a polypeptide of the present invention, either "naked" or in a delivery system as described above, may be followed by administration of an antigen in order to enhance the protective immune effect of the vaccine.

Polypeptides and polynucleotides disclosed herein may also be employed in adoptive immunotherapy for the treatment of *Chlamydial* infection. Adoptive immunotherapy may be broadly classified into either active or passive immunotherapy. In active immunotherapy, treatment relies on the *in vivo* stimulation of the endogenous host immune system with the administration of immune response-modifying agents (for example, vaccines, bacterial adjuvants, and/or cytokines).

In passive immunotherapy, treatment involves the delivery of biologic reagents with established immune reactivity (such as effector cells or antibodies) that can directly or indirectly mediate anti-*Chlamydia* effects and does not necessarily depend on an intact host immune system. Examples of effector cells include T lymphocytes (for example, CD8+ cytotoxic T-lymphocyte, CD4+ T-helper), killer cells (such as Natural Killer cells, lymphokine-activated killer cells), B cells, or antigen presenting cells (such as dendritic cells and macrophages) expressing the disclosed antigens. The polypeptides disclosed herein may also be used to generate antibodies or anti-idiotypic antibodies (as in U.S. Patent No. 4,918,164), for passive immunotherapy.

The predominant method of procuring adequate numbers of T-cells for adoptive immunotherapy is to grow immune T-cells *in vitro*. Culture conditions for expanding single antigen-specific T-cells to several billion in number with retention of antigen recognition *in vivo* are well known in the art. These *in vitro* culture conditions typically utilize intermittent stimulation with antigen, often in the presence of cytokines, such as IL-2, and non-dividing feeder cells. As noted above, the immunoreactive polypeptides described herein may be used to rapidly expand antigen-specific T cell cultures in order to generate sufficient number of cells for immunotherapy. In particular, antigen-presenting cells, such as dendritic, macrophage, monocyte, fibroblast, or B-cells, may be pulsed with immunoreactive polypeptides, or polynucleotide sequence(s) may be introduced into antigen presenting cells, using a

variety of standard techniques well known in the art. For example, antigen presenting cells may be transfected or transduced with a polynucleotide sequence, wherein said sequence contains a promoter region appropriate for increasing expression, and can be expressed as part of a recombinant virus or other expression system. Several viral
5 vectors may be used to transduce an antigen presenting cell, including pox virus, vaccinia virus, and adenovirus; also, antigen presenting cells may be transfected with polynucleotide sequences disclosed herein by a variety of means, including gene-gun technology, lipid-mediated delivery, electroporation, osmotic shock, and particulate delivery mechanisms, resulting in efficient and acceptable expression levels as
10 determined by one of ordinary skill in the art. For cultured T-cells to be effective in therapy, the cultured T-cells must be able to grow and distribute widely and to survive long term *in vivo*. Studies have demonstrated that cultured T-cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (see, for example, Cheever, M., *et al*, "Therapy
15 With Cultured T Cells: Principles Revisited," *Immunological Reviews*, 157:177, 1997).

The polypeptides disclosed herein may also be employed to generate and/or isolate chlamydial-reactive T-cells, which can then be administered to the patient. In one technique, antigen-specific T-cell lines may be generated by *in vivo* immunization with short peptides corresponding to immunogenic portions of the
20 disclosed polypeptides. The resulting antigen specific CD8+ or CD4+ T-cell clones may be isolated from the patient, expanded using standard tissue culture techniques, and returned to the patient.

Alternatively, peptides corresponding to immunogenic portions of the polypeptides may be employed to generate *Chlamydia* reactive T cell subsets by
25 selective *in vitro* stimulation and expansion of autologous T cells to provide antigen-specific T cells which may be subsequently transferred to the patient as described, for example, by Chang *et al*, (*Crit. Rev. Oncol. Hematol.*, 22(3), 213, 1996). Cells of the immune system, such as T cells, may be isolated from the peripheral blood of a patient, using a commercially available cell separation system, such as Isolex™ System,
30 available from Nexell Therapeutics, Inc. Irvine, CA. The separated cells are stimulated with one or more of the immunoreactive polypeptides contained within a delivery vehicle, such as a microsphere, to provide antigen-specific T cells. The population of

antigen-specific T cells is then expanded using standard techniques and the cells are administered back to the patient.

In other embodiments, T-cell and/or antibody receptors specific for the polypeptides disclosed herein can be cloned, expanded, and transferred into other
5 vectors or effector cells for use in adoptive immunotherapy. In particular, T cells may be transfected with the appropriate genes to express the variable domains from chlamydia specific monoclonal antibodies as the extracellular recognition elements and joined to the T cell receptor signaling chains, resulting in T cell activation, specific lysis, and cytokine release. This enables the T cell to redirect its specificity in an MHC-
10 independent manner. See for example, Eshhar, Z., *Cancer Immunol Immunother*, 45(3-4):131-6, 1997 and Hwu, P., et al, *Cancer Res*, 55(15):3369-73, 1995. Another embodiment may include the transfection of chlamydia antigen specific alpha and beta T cell receptor chains into alternate T cells, as in Cole, DJ, et al, *Cancer Res*, 55(4):748-52, 1995.

15 In a further embodiment, syngeneic or autologous dendritic cells may be pulsed with peptides corresponding to at least an immunogenic portion of a polypeptide disclosed herein. The resulting antigen-specific dendritic cells may either be transferred into a patient, or employed to stimulate T cells to provide antigen-specific T cells which may, in turn, be administered to a patient. The use of peptide-pulsed dendritic cells to
20 generate antigen-specific T cells and the subsequent use of such antigen-specific T cells to eradicate disease in a murine model has been demonstrated by Cheever et al, *Immunological Reviews*, 157:177, 1997). Additionally, vectors expressing the disclosed polynucleotides may be introduced into stem cells taken from the patient and clonally propagated *in vitro* for autologous transplant back into the same patient.

25 Within certain aspects, polypeptides, polynucleotides, T cells and/or binding agents disclosed herein may be incorporated into pharmaceutical compositions or immunogenic compositions (*i.e.*, vaccines). Alternatively, a pharmaceutical composition may comprise an antigen-presenting cell (*e.g.* a dendritic cell) transfected with a *Chlamydial* polynucleotide such that the antigen presenting cell expresses a
30 *Chlamydial* polypeptide. Pharmaceutical compositions comprise one or more such compounds and a physiologically acceptable carrier. Vaccines may comprise one or more such compounds and an immunostimulant. An immunostimulant may be any

substance that enhances or potentiates an immune response to an exogenous antigen. Examples of immunostimulants include adjuvants, biodegradable microspheres (*e.g.*, polylactic galactide) and liposomes (into which the compound is incorporated; *see e.g.*, Fullerton, U.S. Patent No. 4,235,877). Vaccine preparation is generally described in, for example, M.F. Powell and M.J. Newman, eds., "Vaccine Design (the subunit and adjuvant approach)," Plenum Press (NY, 1995). Pharmaceutical compositions and vaccines within the scope of the present invention may also contain other compounds, which may be biologically active or inactive. For example, one or more immunogenic portions of other *Chlamydial* antigens may be present, either incorporated into a fusion polypeptide or as a separate compound, within the composition or vaccine.

A pharmaceutical composition or vaccine may contain DNA encoding one or more of the polypeptides as described above, such that the polypeptide is generated *in situ*. As noted above, the DNA may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacteria and viral expression systems. Numerous gene delivery techniques are well known in the art, such as those described by Rolland, *Crit. Rev. Therap. Drug Carrier Systems* 15:143-198, 1998, and references cited therein. Appropriate nucleic acid expression systems contain the necessary DNA sequences for expression in the patient (such as a suitable promoter and terminating signal). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an immunogenic portion of the polypeptide on its cell surface or secretes such an epitope.

In a preferred embodiment, the DNA may be introduced using a viral expression system (*e.g.*, vaccinia or other pox virus, retrovirus, adenovirus, baculovirus, togavirus, bacteriophage, and the like), which often involves the use of a non-pathogenic (defective), replication competent virus.

For example, many viral expression vectors are derived from viruses of the retroviridae family. This family includes the murine leukemia viruses, the mouse mammary tumor viruses, the human foamy viruses, Rous sarcoma virus, and the immunodeficiency viruses, including human, simian, and feline. Considerations when designing retroviral expression vectors are discussed in Comstock *et al.* (1997).

Excellent murine leukemia virus (MLV)-based viral expression vectors have been developed by Kim *et al.* (1998). In creating the MLV vectors, Kim *et al.* found that the entire *gag* sequence, together with the immediate upstream region, could be deleted without significantly affecting viral packaging or gene expression. Further, it was found that nearly the entire U3 region could be replaced with the immediately-early promoter of human cytomegalovirus without deleterious effects. Additionally, MCR and internal ribosome entry sites (IRES) could be added without adverse effects. Based on their observations, Kim *et al.* have designed a series of MLV-based expression vectors comprising one or more of the features described above.

As more has been learned about human foamy virus (HFV), characteristics of HFV that are favorable for its use as an expression vector have been discovered. These characteristics include the expression of pol by splicing and start of translation at a defined initiation codon. Other aspects of HFV viral expression vectors are reviewed in Bodem *et al.* (1997).

Murakami *et al.* (1997) describe a Rous sarcoma virus (RSV)-based replication-competent avian retrovirus vectors, IR1 and IR2 to express a heterologous gene at a high level. In these vectors, the IRES derived from encephalomyocarditis virus (EMCV) was inserted between the *env* gene and the heterologous gene. The IR1 vector retains the splice-acceptor site that is present downstream of the *env* gene while the IR2 vector lacks it. Murakami *et al.* have shown high level expression of several different heterologous genes by these vectors.

Recently, a number of lentivirus-based retroviral expression vectors have been developed. Kafri *et al.* (1997) have shown sustained expression of genes delivered directly into liver and muscle by a human immunodeficiency virus (HIV)-based expression vector. One benefit of the system is the inherent ability of HIV to transduce non-dividing cells. Because the viruses of Kafri *et al.* are pseudotyped with vesicular stomatitis virus G glycoprotein (VSVG), they can transduce a broad range of tissues and cell types.

A large number of adenovirus-based expression vectors have been developed, primarily due to the advantages offered by these vectors in gene therapy applications. Adenovirus expression vectors and methods of using such vectors are the subject of a number of United States patents, including United States Patent No.

5,698,202, United States Patent No. 5,616,326, United States Patent No. 5,585,362, and United States Patent No. 5,518,913, all incorporated herein by reference.

Additional adenoviral constructs are described in Khatri *et al.* (1997) and Tomanin *et al.* (1997). Khatri *et al.* describe novel ovine adenovirus expression vectors and their ability to infect bovine nasal turbinate and rabbit kidney cells as well as a range of human cell type, including lung and foreskin fibroblasts as well as liver, prostate, breast, colon and retinal lines. Tomanin *et al.* describe adenoviral expression vectors containing the T7 RNA polymerase gene. When introduced into cells containing a heterologous gene operably linked to a T7 promoter, the vectors were able to drive gene expression from the T7 promoter. The authors suggest that this system may be useful for the cloning and expression of genes encoding cytotoxic proteins.

Poxviruses are widely used for the expression of heterologous genes in mammalian cells. Over the years, the vectors have been improved to allow high expression of the heterologous gene and simplify the integration of multiple heterologous genes into a single molecule. In an effort to diminish cytopathic effects and to increase safety, vaccinia virus mutant and other poxviruses that undergo abortive infection in mammalian cells are receiving special attention (Oertli *et al.*, 1997). The use of poxviruses as expression vectors is reviewed in Carroll and Moss (1997).

Togaviral expression vectors, which includes alphaviral expression vectors have been used to study the structure and function of proteins and for protein production purposes. Attractive features of togaviral expression vectors are rapid and efficient gene expression, wide host range, and RNA genomes (Huang, 1996). Also, recombinant vaccines based on alphaviral expression vectors have been shown to induce a strong humoral and cellular immune response with good immunological memory and protective effects (Tubulekas *et al.*, 1997). Alphaviral expression vectors and their use are discussed, for example, in Lundstrom (1997).

In one study, Li and Garoff (1996) used Semliki Forest virus (SFV) expression vectors to express retroviral genes and to produce retroviral particles in BHK-21 cells. The particles produced by this method had protease and reverse transcriptase activity and were infectious. Furthermore, no helper virus could be detected in the virus stocks. Therefore, this system has features that are attractive for its use in gene therapy protocols.

Baculoviral expression vectors have traditionally been used to express heterologous proteins in insect cells. Examples of proteins include mammalian chemokine receptors (Wang *et al.*, 1997), reporter proteins such as green fluorescent protein (Wu *et al.*, 1997), and FLAG fusion proteins (Wu *et al.*, 1997; Koh *et al.*, 1997).
5 Recent advances in baculoviral expression vector technology, including their use in virion display vectors and expression in mammalian cells is reviewed by Possee (1997). Other reviews on baculoviral expression vectors include Jones and Morikawa (1996) and O'Reilly (1997).

Other suitable viral expression systems are disclosed, for example, in
10 Fisher-Hoch *et al.*, *Proc. Natl. Acad. Sci. USA* 86:317-321, 1989; Flexner *et al.*, *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner *et al.*, *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld *et al.*, *Science* 252:431-434, 1991; Kolls *et al.*, *Proc. Natl. Acad. Sci. USA*
15 91:215-219, 1994; Kass-Eisler *et al.*, *Proc. Natl. Acad. Sci. USA* 90:11498-11502, 1993; Guzman *et al.*, *Circulation* 88:2838-2848, 1993; and Guzman *et al.*, *Cir. Res.* 73:1202-1207, 1993. Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. In other systems, the DNA may be introduced as "naked" DNA, as described, for example, in Ulmer *et al.*, *Science*
20 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

It will be apparent that a vaccine may comprise a polynucleotide and/or a polypeptide component, as desired. It will also be apparent that a vaccine may contain
25 pharmaceutically acceptable salts of the polynucleotides and/or polypeptides provided herein. Such salts may be prepared from pharmaceutically acceptable non-toxic bases, including organic bases (*e.g.*, salts of primary, secondary and tertiary amines and basic amino acids) and inorganic bases (*e.g.*, sodium, potassium, lithium, ammonium, calcium and magnesium salts). While any suitable carrier known to those of ordinary
30 skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. Compositions of the present invention may be formulated for any appropriate manner of administration,

including for example, topical, oral, nasal, intravenous, intracranial, intraperitoneal, subcutaneous or intramuscular administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactate polyglycolate) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

Such compositions may also comprise buffers (*e.g.*, neutral buffered saline or phosphate buffered saline), carbohydrates (*e.g.*, glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, bacteriostats, chelating agents such as EDTA or glutathione, adjuvants (*e.g.*, aluminum hydroxide), solutes that render the formulation isotonic, hypotonic or weakly hypertonic with the blood of a recipient, suspending agents, thickening agents and/or preservatives. Alternatively, compositions of the present invention may be formulated as a lyophilizate. Compounds may also be encapsulated within liposomes using well known technology.

Any of a variety of immunostimulants may be employed in the vaccines of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis* derived proteins. Suitable adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI); Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ); AS-2 (SmithKline Beecham, Philadelphia, PA); aluminum salts such as aluminum hydroxide gel (alum) or aluminum phosphate; salts of calcium, iron or zinc; an insoluble suspension of acylated tyrosine; acylated sugars; cationically or anionically derivatized polysaccharides; polyphosphazenes; biodegradable microspheres; monophosphoryl lipid A and quil A. Cytokines, such as GM-CSF or interleukin-2, -7, or -12, may also be used as adjuvants.

Within the vaccines provided herein, under select circumstances, the

adjuvant composition may be designed to induce an immune response predominantly of the Th1 type or Th2 type. High levels of Th1-type cytokines (*e.g.*, IFN- γ , TNF α , IL-2 and IL-12) tend to favor the induction of cell mediated immune responses to an administered antigen. In contrast, high levels of Th2-type cytokines (*e.g.*, IL-4, IL-5, 5 IL-6 and IL-10) tend to favor the induction of humoral immune responses. Following application of a vaccine as provided herein, a patient will support an immune response that includes Th1- and Th2-type responses. Within a preferred embodiment, in which a response is predominantly Th1-type, the level of Th1-type cytokines will increase to a greater extent than the level of Th2-type cytokines. The levels of these cytokines may 10 be readily assessed using standard assays. For a review of the families of cytokines, see Mosmann and Coffman, *Ann. Rev. Immunol.* 7:145-173, 1989.

Preferred adjuvants for use in eliciting a predominantly Th1-type response include, for example, a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A (3D-MPL), together with an aluminum salt. 15 MPL adjuvants are available from Corixa Corporation (Seattle, WA; *see* US Patent Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce a predominantly Th1 response. Such oligonucleotides are well known and are described, for example, in WO 96/02555 and WO 99/33488. Immunostimulatory DNA sequences are also described, 20 for example, by Sato et al., *Science* 273:352, 1996. Another preferred adjuvant is a saponin, preferably QS21 (Aquila Biopharmaceuticals Inc., Framingham, MA), which may be used alone or in combination with other adjuvants. For example, an enhanced system involves the combination of a monophosphoryl lipid A and saponin derivative, such as the combination of QS21 and 3D-MPL as described in WO 94/00153, or a less 25 reactogenic composition where the QS21 is quenched with cholesterol, as described in WO 96/33739. Other preferred formulations comprise an oil-in-water emulsion and tocopherol. A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil-in-water emulsion is described in WO 95/17210.

Other preferred adjuvants include Montanide ISA 720 (Seppic, France), 30 SAF (Chiron, California, United States), ISCOMS (CSL), MF-59 (Chiron), the SBAS series of adjuvants (*e.g.*, SBAS-2 or SBAS-4, available from SmithKline Beecham, Rixensart, Belgium), Detox (Corixa Corporation; Seattle, WA), RC-529 (Corixa

Corporation; Seattle, WA) and other aminoalkyl glucosaminide 4-phosphates (AGPs), such as those described in pending U.S. Patent Application Serial Nos. 08/853,826 and 09/074,720, the disclosures of which are incorporated herein by reference in their entireties.

5 Any vaccine provided herein may be prepared using well known methods that result in a combination of antigen, immunostimulant and a suitable carrier or excipient. The compositions described herein may be administered as part of a sustained release formulation (*i.e.*, a formulation such as a capsule, sponge or gel (composed of polysaccharides, for example) that effects a slow release of compound
10 following administration). Such formulations may generally be prepared using well known technology (*see, e.g.*, Coombes et al., *Vaccine* 14:1429-1438, 1996) and administered by, for example, oral, rectal or subcutaneous implantation, or by implantation at the desired target site. Sustained-release formulations may contain a polypeptide, polynucleotide or antibody dispersed in a carrier matrix and/or contained
15 within a reservoir surrounded by a rate controlling membrane.

Carriers for use within such formulations are biocompatible, and may also be biodegradable; preferably the formulation provides a relatively constant level of active component release. Such carriers include microparticles of poly(lactide-co-glycolide), as well as polyacrylate, latex, starch, cellulose and dextran. Other delayed-
20 release carriers include supramolecular biovectors, which comprise a non-liquid hydrophilic core (*e.g.*, a cross-linked polysaccharide or oligosaccharide) and, optionally, an external layer comprising an amphiphilic compound, such as a phospholipid (*see e.g.*, U.S. Patent No. 5,151,254 and PCT applications WO 94/20078, WO/94/23701 and WO 96/06638). The amount of active compound contained within a sustained release
25 formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented.

Any of a variety of delivery vehicles may be employed within pharmaceutical compositions and vaccines to facilitate production of an antigen-specific immune response that targets *Chlamydia*-infected cells. Delivery vehicles include
30 antigen presenting cells (APCs), such as dendritic cells, macrophages, B cells, monocytes and other cells that may be engineered to be efficient APCs. Such cells may, but need not, be genetically modified to increase the capacity for presenting the antigen,

to improve activation and/or maintenance of the T cell response, to have anti-*Chlamydia* effects *per se* and/or to be immunologically compatible with the receiver (*i.e.*, matched HLA haplotype). APCs may generally be isolated from any of a variety of biological fluids and organs, and may be autologous, allogeneic, syngeneic or
5 xenogeneic cells.

Certain preferred embodiments of the present invention use dendritic cells or progenitors thereof as antigen-presenting cells. Dendritic cells are highly potent APCs (Banchereau and Steinman, *Nature* 392:245-251, 1998) and have been shown to be effective as a physiological adjuvant for eliciting prophylactic or therapeutic
10 immunity (*see* Timmerman and Levy, *Ann. Rev. Med.* 50:507-529, 1999). In general, dendritic cells may be identified based on their typical shape (stellate *in situ*, with marked cytoplasmic processes (dendrites) visible *in vitro*), their ability to take up, process and present antigens with high efficiency, and their ability to activate naïve T cell responses. Dendritic cells may, of course, be engineered to express specific cell-
15 surface receptors or ligands that are not commonly found on dendritic cells *in vivo* or *ex vivo*, and such modified dendritic cells are contemplated by the present invention. As an alternative to dendritic cells, secreted vesicles antigen-loaded dendritic cells (called exosomes) may be used within a vaccine (*see* Zitvogel et al., *Nature Med.* 4:594-600, 1998).

20 Dendritic cells and progenitors may be obtained from peripheral blood, bone marrow, lymph nodes, spleen, skin, umbilical cord blood or any other suitable tissue or fluid. For example, dendritic cells may be differentiated *ex vivo* by adding a combination of cytokines such as GM-CSF, IL-4, IL-13 and/or TNF α to cultures of monocytes harvested from peripheral blood. Alternatively, CD34 positive cells
25 harvested from peripheral blood, umbilical cord blood or bone marrow may be differentiated into dendritic cells by adding to the culture medium combinations of GM-CSF, IL-3, TNF α , CD40 ligand, LPS, flt3 ligand and/or other compound(s) that induce differentiation, maturation and proliferation of dendritic cells.

Dendritic cells are conveniently categorized as "immature" and "mature"
30 cells, which allows a simple way to discriminate between two well characterized phenotypes. However, this nomenclature should not be construed to exclude all possible intermediate stages of differentiation. Immature dendritic cells are

characterized as APC with a high capacity for antigen uptake and processing, which correlates with the high expression of Fcγ receptor and mannose receptor. The mature phenotype is typically characterized by a lower expression of these markers, but a high expression of cell surface molecules responsible for T cell activation such as class I and class II MHC, adhesion molecules (*e.g.*, CD54 and CD11) and costimulatory molecules (*e.g.*, CD40, CD80, CD86 and 4-1BB).

APCs may generally be transfected with a polynucleotide encoding a *Chlamydial* protein (or portion or other variant thereof) such that the *Chlamydial* polypeptide, or an immunogenic portion thereof, is expressed on the cell surface. Such transfection may take place *ex vivo*, and a composition or vaccine comprising such transfected cells may then be used for therapeutic purposes, as described herein. Alternatively, a gene delivery vehicle that targets a dendritic or other antigen presenting cell may be administered to a patient, resulting in transfection that occurs *in vivo*. *In vivo* and *ex vivo* transfection of dendritic cells, for example, may generally be performed using any methods known in the art, such as those described in WO 97/24447, or the gene gun approach described by Mahvi et al., *Immunology and cell Biology* 75:456-460, 1997. Antigen loading of dendritic cells may be achieved by incubating dendritic cells or progenitor cells with the *Chlamydial* polypeptide, DNA (naked or within a plasmid vector) or RNA; or with antigen-expressing recombinant bacterium or viruses (*e.g.*, vaccinia, fowlpox, adenovirus or lentivirus vectors). Prior to loading, the polypeptide may be covalently conjugated to an immunological partner that provides T cell help (*e.g.*, a carrier molecule). Alternatively, a dendritic cell may be pulsed with a non-conjugated immunological partner, separately or in the presence of the polypeptide.

Routes and frequency of administration of pharmaceutical compositions and vaccines, as well as dosage, will vary from individual to individual. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.*, intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.*, by aspiration) or orally. Between 1 and 3 doses may be administered for a 1-36 week period. Preferably, 3 doses are administered, at intervals of 3-4 months, and booster vaccinations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of polypeptide or DNA that, when administered as described above, is capable of raising an immune

response in an immunized patient sufficient to protect the patient from *Chlamydial* infection for at least 1-2 years. In general, the amount of polypeptide present in a dose (or produced *in situ* by the DNA in a dose) ranges from about 1 pg to about 100 mg per kg of host, typically from about 10 pg to about 1 mg, and preferably from about 100 pg to about 1 µg. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 mL to about 5 mL.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactic galactide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

In general, an appropriate dosage and treatment regimen provides the active compound(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit. Such a response can be monitored by establishing an improved clinical outcome in treated patients as compared to non-treated patients. Increases in preexisting immune responses to a *Chlamydial* protein generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or cytokine assays, which may be performed using samples obtained from a patient before and after treatment.

25

DETECTION AND DIAGNOSIS

In another aspect, the present invention provides methods for using the polypeptides described above to diagnose *Chlamydial* infection. In this aspect, methods are provided for detecting *Chlamydial* infection in a biological sample, using one or more of the above polypeptides, either alone or in combination. For clarity, the term “polypeptide” will be used when describing specific embodiments of the inventive

30

diagnostic methods. However, it will be clear to one of skill in the art that the fusion proteins of the present invention may also be employed in such methods.

As used herein, a "biological sample" is any antibody-containing sample obtained from a patient. Preferably, the sample is whole blood, sputum, serum, plasma, saliva, cerebrospinal fluid or urine. More preferably, the sample is a blood, serum or plasma sample obtained from a patient. The polypeptides are used in an assay, as described below, to determine the presence or absence of antibodies to the polypeptide(s) in the sample, relative to a predetermined cut-off value. The presence of such antibodies indicates previous sensitization to *Chlamydia* antigens which may be indicative of *Chlamydia*-infection.

In embodiments in which more than one polypeptide is employed, the polypeptides used are preferably complementary (*i.e.*, one component polypeptide will tend to detect infection in samples where the infection would not be detected by another component polypeptide). Complementary polypeptides may generally be identified by using each polypeptide individually to evaluate serum samples obtained from a series of patients known to be infected with *Chlamydia*. After determining which samples test positive (as described below) with each polypeptide, combinations of two or more polypeptides may be formulated that are capable of detecting infection in most, or all, of the samples tested.

A variety of assay formats are known to those of ordinary skill in the art for using one or more polypeptides to detect antibodies in a sample. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988, which is incorporated herein by reference. In a preferred embodiment, the assay involves the use of polypeptide immobilized on a solid support to bind to and remove the antibody from the sample. The bound antibody may then be detected using a detection reagent that contains a reporter group. Suitable detection reagents include antibodies that bind to the antibody/polypeptide complex and free polypeptide labeled with a reporter group (*e.g.*, in a semi-competitive assay). Alternatively, a competitive assay may be utilized, in which an antibody that binds to the polypeptide is labeled with a reporter group and allowed to bind to the immobilized antigen after incubation of the antigen with the sample. The extent to which components of the sample inhibit the

binding of the labeled antibody to the polypeptide is indicative of the reactivity of the sample with the immobilized polypeptide.

The solid support may be any solid material known to those of ordinary skill in the art to which the antigen may be attached. For example, the solid support
5 may be a test well in a microtiter plate, or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681.

10 The polypeptides may be bound to the solid support using a variety of techniques known to those of ordinary skill in the art. In the context of the present invention, the term "bound" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the antigen and functional groups on the support or may be a linkage by way of a cross-linking agent).
15 Binding by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the polypeptide, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride)
20 with an amount of polypeptide ranging from about 10 ng to about 1 μ g, and preferably about 100 ng, is sufficient to bind an adequate amount of antigen.

Covalent attachment of polypeptide to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the
25 polypeptide. For example, the polypeptide may be bound to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the polypeptide (*see, e.g.,* Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is an enzyme linked immunosorbent
30 assay (ELISA). This assay may be performed by first contacting a polypeptide antigen that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that antibodies to the polypeptide within the sample are allowed

to bind to the immobilized polypeptide. Unbound sample is then removed from the immobilized polypeptide and a detection reagent capable of binding to the immobilized antibody-polypeptide complex is added. The amount of detection reagent that remains bound to the solid support is then determined using a method appropriate for the specific detection reagent.

More specifically, once the polypeptide is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin (BSA) or Tween 20™ (Sigma Chemical Co., St. Louis, MO) may be employed. The immobilized polypeptide is then incubated with the sample, and antibody is allowed to bind to the antigen. The sample may be diluted with a suitable diluent, such as phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is that period of time that is sufficient to detect the presence of antibody within an HGE-infected sample. Preferably, the contact time is sufficient to achieve a level of binding that is at least 95% of that achieved at equilibrium between bound and unbound antibody. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. Detection reagent may then be added to the solid support. An appropriate detection reagent is any compound that binds to the immobilized antibody-polypeptide complex and that can be detected by any of a variety of means known to those in the art. Preferably, the detection reagent contains a binding agent (such as, for example, Protein A, Protein G, immunoglobulin, lectin or free antigen) conjugated to a reporter group. Preferred reporter groups include enzymes (such as horseradish peroxidase), substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. The conjugation of binding agent to reporter group may be achieved using standard methods known to those of ordinary skill in the art. Common binding agents may also be purchased conjugated to a variety of reporter groups from many commercial sources (*e.g.*, Zymed Laboratories, San Francisco, CA, and Pierce, Rockford, IL).

The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound antibody. An appropriate amount of time may generally be determined from the manufacturer's instructions or by assaying the level of binding that occurs over a period of time.

5 Unbound detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin

10 may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally 'for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of anti-*Chlamydia* antibodies in

15 the sample, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value is the average mean signal obtained when the immobilized antigen is incubated with samples from an uninfected patient. In general, a sample generating a signal that is three standard deviations above

20 the predetermined cut-off value is considered positive for *Chlamydia*-infection. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, pp. 106-107. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true

25 positive rates (*i.e.*, sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (*i.e.*, the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered

30 positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In

general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for Chlamydial infection.

In a related embodiment, the assay is performed in a rapid flow-through or strip test format, wherein the antigen is immobilized on a membrane, such as nitrocellulose. In the flow-through test, antibodies within the sample bind to the immobilized polypeptide as the sample passes through the membrane. A detection reagent (*e.g.*, protein A-colloidal gold) then binds to the antibody-polypeptide complex as the solution containing the detection reagent flows through the membrane. The detection of bound detection reagent may then be performed as described above. In the strip test format, one end of the membrane to which polypeptide is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing detection reagent and to the area of immobilized polypeptide. Concentration of detection reagent at the polypeptide indicates the presence of anti-*Chlamydia* antibodies in the sample. Typically, the concentration of detection reagent at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of polypeptide immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of antibodies that would be sufficient to generate a positive signal in an ELISA, as discussed above. Preferably, the amount of polypeptide immobilized on the membrane ranges from about 25 ng to about 1 μ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount (*e.g.*, one drop) of patient serum or blood.

Of course, numerous other assay protocols exist that are suitable for use with the polypeptides of the present invention. The above descriptions are intended to be exemplary only. One example of an alternative assay protocol which may be usefully employed in such methods is a Western blot, wherein the proteins present in a biological sample are separated on a gel, prior to exposure to a binding agent. Such techniques are well known to those of skill in the art.

BINDING AGENTS AND THEIR USES

The present invention further provides agents, such as antibodies and antigen-binding fragments thereof, that specifically bind to a *Chlamydial* protein. As

used herein, an antibody, or antigen-binding fragment thereof, is said to "specifically bind" to a *Chlamydial* protein if it reacts at a detectable level (within, for example, an ELISA) with a *Chlamydial* protein, and does not react detectably with unrelated proteins under similar conditions. As used herein, "binding" refers to a noncovalent association between two separate molecules such that a complex is formed. The ability to bind may be evaluated by, for example, determining a binding constant for the formation of the complex. The binding constant is the value obtained when the concentration of the complex is divided by the product of the component concentrations. In general, two compounds are said to "bind," in the context of the present invention, when the binding constant for complex formation exceeds about 10^3 L/mol. The binding constant may be determined using methods well known in the art.

Binding agents may be further capable of differentiating between patients with and without a *Chlamydial* infection using the representative assays provided herein. In other words, antibodies or other binding agents that bind to a *Chlamydial* protein will generate a signal indicating the presence of a *Chlamydial* infection in at least about 20% of patients with the disease, and will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without infection. To determine whether a binding agent satisfies this requirement, biological samples (e.g., blood, sera, sputum urine and/or tissue biopsies) from patients with and without *Chlamydial* infection (as determined using standard clinical tests) may be assayed as described herein for the presence of polypeptides that bind to the binding agent. It will be apparent that a statistically significant number of samples with and without the disease should be assayed. Each binding agent should satisfy the above criteria; however, those of ordinary skill in the art will recognize that binding agents may be used in combination to improve sensitivity.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome, with or without a peptide component, an RNA molecule or a polypeptide. In a preferred embodiment, a binding agent is an antibody or an antigen-binding fragment thereof. Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, antibodies can be produced by cell culture techniques, including the generation

of monoclonal antibodies as described herein, or via transfection of antibody genes into suitable bacterial or mammalian cell hosts, in order to allow for the production of recombinant antibodies. In one technique, an immunogen comprising the polypeptide is initially injected into any of a wide variety of mammals (*e.g.*, mice, rats, rabbits, sheep
5 or goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule
10 incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for an antigenic polypeptide of interest
15 may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (*i.e.*, reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as
20 described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid
25 cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and their culture supernatants tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

30 Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable

vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process
5 in, for example, an affinity chromatography step.

Within certain embodiments, the use of antigen-binding fragments of antibodies may be preferred. Such fragments include Fab fragments, which may be prepared using standard techniques. Briefly, immunoglobulins may be purified from rabbit serum by affinity chromatography on Protein A bead columns (Harlow and Lane,
10 *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988) and digested by papain to yield Fab and Fc fragments. The Fab and Fc fragments may be separated by affinity chromatography on protein A bead columns.

Monoclonal antibodies of the present invention may be coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides,
15 differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed
20 antiviral protein.

A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such
25 as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an
30 antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical

reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, 5 sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spitler), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of 15 derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers which provide multiple sites for 25 attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.). A carrier may 30 also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating

compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

A variety of routes of administration for the antibodies and immunoconjugates may be used. Typically, administration will be intravenous, intramuscular, subcutaneous or in site-specific regions by appropriate methods. It will be evident that the precise dose of the antibody/immunoconjugate will vary depending upon the antibody used, the antigen density, and the rate of clearance of the antibody.

Antibodies may be used in diagnostic tests to detect the presence of *Chlamydia* antigens using assays similar to those detailed above and other techniques well known to those of skill in the art, thereby providing a method for detecting Chlamydial infection in a patient.

Diagnostic reagents of the present invention may also comprise DNA sequences encoding one or more of the above polypeptides, or one or more portions thereof. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify *Chlamydia*-specific cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for a DNA molecule encoding a polypeptide of the present invention. The presence of the amplified cDNA is then detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes specific for a DNA molecule encoding a polypeptide of the present invention may be used in a hybridization assay to detect the presence of an inventive polypeptide in a biological sample.

The following Examples are offered by way of illustration and not by way of limitation.

EXAMPLE 1

CD4 T CELL EXPRESSION CLONING FOR THE IDENTIFICATION
OF T CELL STIMULATING ANTIGENS FROM CHLAMYDIA TRACHOMATIS
SEROVAR E

5

In this example, a CD4⁺ T cell expression cloning strategy was used to identify Chlamydia trachomatis antigens recognized by patients enrolled in Corixa Corporation's blood donor program. A genomic library of Chlamydia trachomatis serovar E was constructed and screened with Chlamydia specific T cell lines generated by stimulating PBMCs from these donors. Donor CT1 is a 27 yr. old male whose clinical manifestation was non-gonococcal urethritis and his urine was tested positive for Chlamydia by ligase chain reaction. Donor CT3 is a 43 yr. old male who is asymptomatic and infected with serovar J. Donor CT10 is a 24 yr. old female who is asymptomatic and was exposed to Chlamydia through her partner but did not develop the disease. Donor CT11 is a 24 yr. old female with multiple infections (serovar J, F and E).

Chlamydia specific T-cell lines were generated from donors with chlamydial genital tract infection or donors exposed to chlamydia who did not develop the disease. T cell lines from donor CT-1, CT-3 and CT-10 were generated by stimulating PBMC's with reticulate bodies of C. trachomatis serovar E. T-cell lines from donor CT-11 were generated by stimulating PBMC's with either reticulate bodies or elementary bodies of C. trachomatis serovar E. A randomly sheared genomic library of C. trachomatis serovar E was constructed in lambda Zap II vector and an amplified library plated out in 96 well microtiter plates at a density of 25 clones/well. Bacteria were induced to express the recombinant protein in the presence of 2mM IPTG for 2hr, then pelleted and resuspended in 200ul RPMI/10% FBS. 10 ul of the induced bacterial suspension was transferred to 96 well plates containing autologous monocyte-derived dendritic cells. After a 2 hour incubation, dendritic cells were washed to remove E. coli and the T cells were added. Positive E. coli pools were identified by determining IFN gamma production and proliferation of T cells in the pools. The number of pools identified by each T-cell line is as follows: CT1 line : 30/480 pools; CT3 line : 91/960

pools; CT10 line : 40/480 pools; CT11 line : 51/480 pools. The clones identified using this approach are set forth in SEQ ID NO:1-14.

In another example using substantially the same approach described
 5 above, we identified 12 additional T-cell reactive clones from *Chlamydia trachomatis* serovar E expression screening. Clone E5-E9-3 (CT1 positive) contains a 636 bp insert that encodes partially the ORF for dnaK like gene. Part of this sequence was also identified in clone E1-A5-53. Clone E4-H3-56 (CT1 positive, 463 bp insert) contains a partial ORF for the TSA gene (CT603) on the complementary strand. The insert for
 10 clone E2-G12-52 (1265 bp) was identified with the CT11 line. It contains a partial ORF for clpB, a protease ATPase. Another clone identified with the CT11 line, E1-F9-79 (167 bp), contains a partial ORF for the gene CT133 on the complementary strand. CT133 is a predicted rRNA methylase. Clone E4-D2-79 (CT3 positive) contains a 1181 bp insert that is a partial ORF for nrdA gene. The ORF for this gene was also identified
 15 in clone E2-B10-52 (CT10 positive). Clone E6-C8-95 contains a 731 bp insert that was identified using the donor lines CT3, CT1, and CT12. This insert has a carboxy terminal half for the gene for the 60 kDa ORF. Clone E7-H11-61 (CT3 positive-1135 bp) has partial inserts for fliA (CT061), tyrS (CT062), TSA (CT603) and a hypothetical protein (CT602). The insert for clone E5-A11-8 (CT10 positive-1736 bp) contains the complete
 20 ORF for groES (CT111) and a majority of the ORF for groEL (CT110). Clone E3-F2-37 (CT10, CT3, CT11, and CT12 positive-1377bp insert) contains a partial ORF for gene tRNA-Trp (CT322) and a complete ORF for the gene secE (CT321). E4-G9-75 is another CT10 clone that contains a partial ORF (723 bp insert) for the amino terminal region of the pmpH gene (CT872). Clone E2-D5-89 (516bp) is also a CT10 positive
 25 clone that contains a partial ORF for pmpD gene (12). The insert for clone E5-E2-10 (CT10 positive) is 427 bp and contains a partial ORF for the major outer membrane protein omp1.

Example 2

30 ADDITIONAL CD4 T CELL EXPRESSION CLONING FOR THE
IDENTIFICATION OF T CELL STIMULATING ANTIGENS FROM CHLAMYDIA
TRACHOMATIS SEROVAR E

Twenty sequences were isolated from single clones using a *Chlamydia trachomatis* serovar E (Ct E) library expression screening method. Descriptions of how the clones and lines were generated are provided in Example 1.

Clone E5-A8-85 (identified using the CT1 patient line) was found to
5 contain a 1433 bp insert. This insert contains a large region of the C-terminal half of the CT875, a *Chlamydia trachomatis* hypothetical specific gene that is disclosed in SEQ ID NO:34. Also present in the clone is a partial open reading frame (ORF) of a hypothetical protein CT001 which is on the complementary strand.

The clone E9-G2-93 (identified using the C10 patient line) was shown to
10 contain a 554 bp insert, the sequence of which is disclosed in SEQ ID NO:33. This sequence encodes a partial ORF for CT178, a hypothetical CT protein.

Clone E7-B1-16 (identified using the patient lines CT10, CT3, CT5, CT11, CT13, and CHH037) has a 2577 bp insert, the sequence of which is disclosed in SEQ ID NO:32. This clone was found to contain three ORFs. The first ORF contains
15 almost the entire ORF for CT694, a *Chlamydia trachomatis* (CT) specific hypothetical protein. The second ORF is a full length ORF for CT695, another hypothetical CT protein. The third ORF is the N-terminal portion of CT696.

Clone E9-D5-8 (identified using the patient lines CT10, CT1, CT4, and CT11) contains a 393 bp insert, which is disclosed in SEQ ID NO:31. It was found to
20 encode a partial ORF for CT680, the S2 ribosomal protein.

Clone E9-E10-51 (identified using the patient line CT10) contains an 883 bp insert, the sequence of which is disclosed in SEQ ID NO:30. This clone contains two partial ORF. The first of these is for the C-terminal half of CT680, which may show some overlap with the insert present in clone E9-D5-8. The second ORF is
25 the N-terminal partial ORF for CT679, which is the elongation factor TS.

Clone E3-B4-18 (identified using the CT1 patient line) contains a 1224 bp insert, the sequence of which is disclosed in SEQ ID NO:29. This clone contains 4 ORFs. At the N-terminal end of the clone is the complete ORF for CT772, coding for inorganic pyrophosphatase. The second ORF is a small portion of the C-terminal end of
30 CT771, on the complementary frame. The third is a partial ORF of the hypothetical protein, CT191 and the fourth is a partial ORF for CT190, DNA gyrase-B.

Clone E10-B2-57 (identified using the CT10 patient line) contains an 822 bp insert, the sequence of which is disclosed in SEQ ID NO:42. This clone contains the complete ORF for CT066, a hypothetical protein, on the complementary strand.

5 Clone E3-F3-18 (identified using the CT1 patient line) contains an 1141 bp insert, the sequence of which is disclosed in SEQ ID NO:41. It contains a partial ORF for pmpG (CT871) in frame with the β -gal gene.

Clone E4-D6-21 (identified using the CT3 patient line) contains a 1297 bp insert, the sequence of which is disclosed in SEQ ID NO:40. This clone contains a
10 very small portion of xseA (CT329), the entire ORF for tpiS (CT328) on the complementary strand, and a partial amino terminal ORF for trpC (CT327) on the top frame.

Clone E1-G9-23 (identified using the CT3 patient line) contains an 1180 bp insert, the sequence of which is disclosed in SEQ ID NO:39. This clone contains
15 almost the entire ORF for glycogen synthase (CT798).

Clone E3-A3-31 (identified using the CT1 patient line) contains an 1834 bp insert, the sequence of which is disclosed in SEQ ID NO:38. This clone contains a large region of the hypothetical gene CT622.

Clone E2-F7-11 (identified using both the CT3 and CT10 patient lines)
20 contains a 2093 bp insert, the sequence of which is disclosed in SEQ ID NO:37. This clone contains a large region of the rpoN gene (CT609) in frame with β -gal and the complete ORF for the hypothetical gene CT610 on the complementary strand. In addition, it also contains the carboxy-terminal end of CT611, another hypothetical gene.

Clone E7-H11-10 (identified using the CT3 patient line) contains a 1990
25 bp insert, the sequence of which is disclosed in SEQ ID NO:36. This clone contains the amino terminal partial ORF for CT610, a complete ORF for CT611, another complete ORF for CT612, and a carboxy-terminal portion of CT613. All of these genes are hypothetical and all are present on the complementary strand.

Clone E10-C6-45 (identified using the CT3 patient line) contains a 196
30 bp insert, the sequence of which is disclosed in SEQ ID NO:35. This clone contains a partial ORF for nrdA (CT827) in frame with β -gal. This clone contains a relatively

small insert and has particular utility in determining the epitope of this gene that contributes to the immunogenicity of Serovar E.

Clone E3-H6-10 (identified using the CT12 patient line) contains a 3734 bp insert, the sequence of which is disclosed in SEQ ID NO:48. This clone contains
5 ORFs for a series of hypothetical proteins. It contains the partial ORFs for CT223 and CT229 and the complete ORFs for CT224, CT225, CT226, CT227, and CT228.

Clone E4-C3-40 (identified using the CT10 patient line) contains a 2044 bp insert, the sequence of which is disclosed in SEQ ID NO:47. This clone contains a partial ORF for *nrdA* (CT827) and the complete ORF for *nrdB* (CT828).

10 Clone E2-D8-19 (identified using the CT1 patient line) contains a 2010 bp insert, the sequence of which is disclosed in SEQ ID NO:46. This clone contains ORF from the *Chlamydia trachomatis* plasmid as well as containing partial ORFs for ORF3 and ORF6, and complete ORFs for ORF4 and ORF5.

Clone E3-D10-46 (identified using the patient lines CT1, CT3, CT4,
15 CT11, and CT12) contains a 1666 bp insert, the sequence of which is identified in SEQ ID NO: 45. This clone contains a partial ORF for CT770 (*fab F*), a complete ORF for CT771 (hydrolase/phosphatase homologue), a complete ORF for CT772 (*ppa*, inorganic phosphatase), and a partial ORF for CT773 (*Idh*, Leucine dehydrogenase).

Clone E10-H8-1 (identified using both the CT3 and CT10 patient lines)
20 contains an 1862 bp insert, the sequence of which is disclosed in SEQ ID NO:44. It contains the partial ORFs for CT871 (*pmpG*) as well as CT872 (*pmpH*).

Clone E3-F3-7 (identified using the CT1 patient line) contains a 1643 bp insert, the sequence of which is identified in SEQ ID NO:43. It contains the partial ORFs for both CT869 (*pmpE*) and CT870 (*pmpF*).

25

EXAMPLE 3

ADDITIONAL CD4 T CELL EXPRESSION CLONING FOR THE IDENTIFICATION OF T CELL STIMULATING ANTIGENS FROM CHLAMYDIA TRACHOMATIS SEROVAR E

30

The T cell line CHH037 was generated from a 22 year-old healthy female sero-negative for *Chlamydia*. This line was used to screen the *Chlamydia*

trachomatis serovar E library. Nineteen clones were identified from this screen, as described below.

Clone E7-B12-65, contains an 1179 bp insert, the sequence of which is disclosed in SEQ ID NO:114. It contains the complete ORF of the gene for Malate
5 dehydrogenase (CT376) on the complementary strand.

Clone E4-H9-83 contains a 772 bp insert, the sequence of which is identified in SEQ ID NO:115. It contains the partial ORF for the heat shock protein GroEL (CT110).

Clone E9-B10-52 contains a 487 bp insert, the sequence of which is
10 identified in SEQ ID NO:116. It contains a partial ORF for the gene yscC (CT674), a general secretion pathway protein.

Clone E7-A7-79 contains a 1014 bp insert, the sequence of which is disclosed in SEQ ID NO:117. It contains the complete ORF for the histone like development gene, hctA (CT743) and a partial ORF for the rRNA methyltransferase
15 gene ygcA (CT742).

Clone E2-D11-18 contains a 287 bp insert, the sequence of which is disclosed in SEQ ID NO:118. It contains the partial ORF for hctA (CT743).

Clone E9-H6-15, identified using the CT3 line, contains a 713 bp insert
20 the sequence of which is disclosed in SEQ ID NO:125. It contains the partial ORF of the pmpB gene (CT413).

Clone E3-D10-87, identified using the CT1 line, contains a 780 bp insert, the sequence of which is disclosed in SEQ ID NO:126. It contains the partial ORF for CT388, a hypothetical gene, on the complementary strand, and a partial ORF
25 for CT389, another hypothetical protein.

Clone E9-D6-43, identified using the CT3 line, contains a 433 bp insert, the sequence of which is disclosed in SEQ ID NO:127. It contains a partial ORF for CT858.

Clone E3-D10-4, identified using the CT1 line, contains an 803 bp
30 insert, the sequence of which is disclosed in SEQ ID NO:128. It contains a partial ORF for pGP3-D, an ORF encoded on the plasmid pCHL1.

Clone E3-G8-7, identified using the CT1 line, contains an 842 bp insert, the sequence of which is disclosed in SEQ ID NO:129. It contains partial ORFs for CT557 (Lpda) and CT558 (LipA).

Clone E3-F11-32, identified using the CT1 line, contains an 813 bp insert, the sequence of which is disclosed in SEQ ID NO:130. It contains a partial ORF for pmpD (CT812).

Clone E2-F8-5, identified using the CT12 line, contains a 1947 bp insert, the sequence of which is disclosed in SEQ ID NO:131. It contains a complete ORF for the 15 kDa ORF (CT442) and a partial ORF for the 60 kDa ORF (CT443).

Clone E2-G4-39, identified using the CT12 line, contains a 1278 bp insert, the sequence of which is disclosed in SEQ ID NO:132. It contains the partial ORF of the 60kDa ORF (CT443).

Clone E9-D1-16, identified using the CT10 line, contains a 916 bp insert, the sequence of which is disclosed in SEQ ID NO:133. It contains the partial ORF for the pmpH (CT872).

Clone E3-F3-6, identified using the CT1 line, contains a 751 bp insert, the sequence of which is disclosed in SEQ ID NO:134. It contains the partial ORFs, all on the complementary strand, for genes accB (CT123), L13 ribosomal (CT125), and S9 ribosomal (CT126).

Clone E2-D4-70, identified using the CT12 line, contains a 410 bp insert, the sequence of which is disclosed in SEQ ID NO:135. It contains the partial ORF for the pmpC gene (CT414).

Clone E5-A1-79, identified using the CT1 line, contains a 2719 bp insert, the sequence of which is disclosed in SEQ ID NO:136. It contains a partial ORF for ydhO (CT127), a complete ORF for S9 ribosomal gene (CT126 on the complementary strand), a complete ORF for the L13 ribosomal gene (CT125 on the complementary strand) and a partial ORF for accC (CT124 on the complementary strand).

Clone E1-F7-16, identified using the lines CT12, CT3, and CT11, contains a 2354 bp insert, the sequence of which is disclosed in SEQ ID NO:137. It contains a partial ORF of the ftsH gene (CT841) and the entire ORF for the pnp gene (CT842) on the complementary strand.

Clone E1-D8-62, identified using the CT12 line, contains an 898 bp insert, the sequence of which is disclosed in SEQ ID NO:138. It contains partial ORFs for the *ftsH* gene (CT841) and for the *pnp* gene (CT842).

5

EXAMPLE 4

EXPRESSION OF CHLAMYDIA TRACOMATIS RECOMBINANT PROTEINS

Several *Chlamydia trachomatis* serovar E specific genes were cloned into pET17b. This plasmid incorporates a 6X histidine tag at the N-terminal to allow for expression and purification of recombinant protein.

Two full-length recombinant proteins, CT622 and CT875, were expressed in *E. coli*. Both of these genes were identified using CtLGVII expression screening, but the serovar E homologues were expressed. The primers used to amplify these genes were based on serovar D sequences. The genes were amplified using serovar E genomic DNA as the template. Once amplified, the fragments were cloned in pET-17b with a N-terminal 6X-His Tag. After transforming the recombinant plasmid in XL-I blue cells, the DNA was prepared and the clones fully sequenced. The DNA was then transformed into the expression host BL21-pLysS cells (Novagen) for production of the recombinant proteins. The proteins were induced with IPTG and purified on Ni-NTA agarose using standard methods. The DNA sequences for CTE622 and CTE875 are disclosed in SEQ ID NO:28 and 27 respectively, and their amino acid sequences are disclosed in SEQ ID NO: 140 and 139, respectively

Five additional *Chlamydia trachomatis* genes were cloned. The *Chlamydia trachomatis* specific protein CT694, the protein CT695, and the L1 ribosomal protein, the DNA sequences of which are disclosed in SEQ ID NO:119, 120 and 121 respectively. The protein sequences of these 6X-histidine recombinant proteins are disclosed in SEQ ID NO: 122 (CT694), 123 (CT695), and 124 (L1 ribosomal protein). The genes CT875 and CT622, from serovar E were also cloned using pET17b as 6X-His fusion proteins. These recombinant proteins were expressed and purified and their the amino acid sequences disclosed in SEQ ID NO:139 and 140, respectively.

EXAMPLE 5

RECOMBINANT CHLAMYDIAL ANTIGENS RECOGNIZED BY T CELL LINES

Patient T cell lines were generated from the following donors: CT1, CT2, CT3,
 5 CT4, CT5, CT6, CT7, CT8, CT9, CT10, CT11, CT12, CT13, CT14, CT15, and CT16.
 A summary of their details is included in Table II.

Table II: <i>C. trachomatis</i> patients						
Patients	Gender	Age	Clinical Manifestation	Serovar	IgG titer	Multiple Infections
CT1	M	27	NGU	LCR	Negative	No
CT2	M	24	NGU	D	Negative	E
CT3	M	43	Asymptomatic Shed Eb Dx was HPV	J	Ct 1:512 Cp 1:1024 Cps 1:256	No
CT4	F	25	Asymptomatic Shed Eb	J	Ct 1:1024	Y
CT5	F	27	BV	LCR	Ct 1:256 Cp 1:256	F/F
CT6	M	26	Perinial rash Discharge, dysuria	G	Cp 1:1024	N
CT7	F	29	BV Genital ulcer	E	Ct 1:512 Cp 1:1024	N
CT8	F	24	Not Known	LCR	Not tested	NA
CT9	M	24	asymptomatic	LCR	Ct 1:128 Cp 1:128	N

CT10	F	20	Mild itch vulvar	negative	negative	12/1/98
CT11	F	21	BV Abnormal pap smear	J	Ct 1:512	F/F/J/E/E PID 6/96
CT12	M	20	asymptomatic	LCR	Cp 1:512	N
CT13	F	18	BV, gonorrhea, Ct vaginal discharge, dysuria	G	Ct 1:1024	N
CT14	M	24	NGU	LCR	Ct 1:256 Cp 1:256	N
CT15	F	21	Muco-purulent cervicitis Vaginal discharge	culture	Ct 1:256 Ct IgM 1:320 Cp 1:64	N
CT16	M	26	Asymptomatic/ contact	LCR	NA	N
CL8	M	38	No clinical history of disease	negative	negative	No

NGU=Non-Gonococcal Urethritis; BV=Bacterial Vaginosis; CT=Chlamydia trachomatis; Cp=Chlamydia pneumoniae; Eb=Chlamydia elementary bodies; HPV=human papilloma virus; Dx=diagnosis; PID=pelvic inflammatory disease;

5 LCR=Ligase change reaction.

PBMC were collected from a second series of donors and T cell lines have been generated from a sub-set of these. A summary of the details for three such T cell lines is listed in the table below.

Table III: Normal Donors				
Donor	Gender	Age	CT IgG Titer	CP IgG Titer
CHH011	F	49	1:64	1:16
CHH037	F	22	0	0
CHH042	F	25	0	1:16

Donor CHH011 is a healthy 49 year old female donor sero-negative for
5 *C. trachomatis*. PBMC produced higher quantities of IFN-gamma in response to *C. trachomatis* elementary bodies as compared to *C. pneumoniae* elementary bodies, indicating a *C. trachomatis*-specific response. Donor CHH037 is a 22 year old healthy female donor sero-negative for *C. trachomatis*. PBMC produced higher quantities of IFN-gamma in response to *C. trachomatis* elementary bodies as compared to *C. pneumoniae* elementary bodies, indicating a *C. trachomatis*-specific response. CHH042
10 is a 25 year old healthy female donor with an IgG titer of 1:16 to *C. pneumoniae*. PBMC produced higher quantities of IFN-gamma in response to *C. trachomatis* elementary bodies as compared to *C. pneumoniae* elementary bodies, indicating a *C. trachomatis*-specific response.

15 Recombinant proteins for several *Chlamydia trachomatis* genes were generated as described above. Sequences for MOMP was derived from serovar F. The genes CT875, CT622, pmp-B-2, pmpA, and CT529 were derived from serovar E and sequences for the genes gro-EL, Swib, pmpD, pmpG, TSA, CT610, pmpC, pmpE, S13, lpdA, pmpI, and pmpH-C were derived from LII.

20 Several of the patient and donor lines described above were tested against the recombinant Chlamydia proteins. Table IV summarizes the results of the T cell responses to the recombinant Chlamydia proteins.

Table IV: Recombinant Chlamydia Antigens Recognized By T Cell Lines

Antigen	Serovar	#of hits	CL 8 L2	CT 10 E	CT 1 E	CT 3 E	CT 4 L2	CT 5 E	CT 11 E	CT 12 E	CT 13 E	CHH- 011 E	CHH- 037 E
gro-EL (CT110)	L2	10	-	+	+	+	+	+	+	+	+	+	+
MompF (CT681)	F	10	-	+	+	+	+	+	+	+	+	+	+
CT875	E	8	-	+	+	-	+	+	+	+	+	-	+
SWIB (CT460)	L2	8	+	+	-	+	-	+	-	+	+	+	+
pmpD (CT812)	L2	5	-	+	+	+	+	-	-	+	+	-	-
pmpG (CT871)	L2	6	-	+	+	-	+	+	nt	-	+	+	-
TSA (CT603)	L2	6	-	-	+	+	+	+	-	-	+	-	+
CT622	E	3	-	-	+	-	+	-	-	-	+	-	-
CT610	L2	3	-	+	-	+	-	-	-	+	-	-	-
pmpB-2 (CT413)	E	3	-	-	+	+	+	-	-	-	-	-	-
pmpC (CT414)	L2	4	-	-	-	+	-	+	-	+	-	-	+
pmpE (CT869)	L2	3	-	+	+	-	-	-	-	-	+	-	-
S13 (CT509)	L2	2	+	-	-	-	+	-	-	-	-	-	-
lpdA (CT557)	L2	3	-	-	+	+	-	-	-	-	-	+	-
pmpI (CT874)	L2	2	-	-	+	-	-	-	-	-	-	+	-
pmpH-C (CT872)	L2	1	-	-	-	-	-	-	-	+	-	-	-

pmpA (CT412)	E	0	-	-	-	-	-	-	-	-	-	-	-
CT529	E	0	-	-	-	-	-	-	-	-	-	-	-

Although the present invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, changes and
5 modifications can be carried out without departing from the scope of the invention which is intended to be limited only by the scope of the appended claims.

Claims

What is Claimed:

1. An isolated polynucleotide comprising a sequence selected from the group consisting of:

- (a) sequences provided in SEQ ID NO:1-48, 114-121, and 125-138;
- (b) complements of the sequences provided in SEQ ID NO: 1-48, 114-121, and 125-138;
- (c) sequences consisting of at least 20 contiguous residues of a sequence provided in SEQ ID NO:1-48, 114-121, 125-138;
- (d) sequences that hybridize to a sequence provided in SEQ ID NO:1-48, 114-121, and 125-138, under highly stringent conditions;
- (e) sequences having at least 95% identity to a sequence of SEQ ID NO:1-48, 114-121, and 125-138;
- (f) sequences having at least 99% identity to a sequence of SEQ ID NO: 1-48, 114-121, and 125-138; and
- (g) degenerate variants of a sequence provided in SEQ ID NO: 1-48, 114-121, and 125-138.

2. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- (a) sequences encoded by a polynucleotide of claim 1;
- (b) sequences having at least 95% identity to a sequence encoded by a polynucleotide of claim 1; and
- (c) sequences having at least 99% identity to a sequence encoded by a polynucleotide of claim 1.

3. An isolated polypeptide comprising at least an immunogenic fragment of a polypeptide sequence selected from the group consisting of:

- (a) a polypeptide sequence set forth in SEQ ID NO: 122-124 and 139-140,

(b) a polypeptide sequence having at least 95% identity with a sequence set forth in SEQ ID NO: 122-124 and 139-140, and

(c) a polypeptide sequence having at least 99% identity with a sequence set forth in SEQ ID NO: 122-124 and 139-140.

4. An expression vector comprising a polynucleotide of claim 1 operably linked to an expression control sequence.

5. A host cell transformed or transfected with an expression vector according to claim 4.

6. An isolated antibody, or antigen-binding fragment thereof, that specifically binds to a polypeptide of claim 2 or claim 3.

7. A method for detecting the presence of Chlamydia in a patient, comprising the steps of:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with a binding agent that binds to a polypeptide of claim 2 or claim 3;
- (c) detecting in the sample an amount of polypeptide that binds to the binding agent; and
- (d) comparing the amount of polypeptide to a predetermined cut-off value and therefrom determining the presence of Chlamydia in the patient.

8. A fusion protein comprising at least one polypeptide according to claim 2 or claim 3.

9. An oligonucleotide that hybridizes to a sequence recited in any one of SEQ ID NO: 1-48, 114-121, and 125-138 under highly stringent conditions.

10. A method for stimulating and/or expanding T cells specific for a Chlamydia protein, comprising contacting T cells with at least one component selected from the group consisting of:

- (a) a polypeptide according to claim 2 or claim 3;
- (b) a polynucleotide according to claim 1; and
- (c) an antigen-presenting cell that expresses a polynucleotide according to claim 1,

under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells.

11. An isolated T cell population, comprising T cells prepared according to the method of claim 10.

12. A composition comprising a first component selected from the group consisting of physiologically acceptable carriers and immunostimulants, and a second component selected from the group consisting of:

- (a) a polypeptide according to claim 2 or claim 3;
- (b) a polynucleotide according to claim 1;
- (c) an antibody according to claim 6;
- (d) a fusion protein according to claim 8;
- (e) a T cell population according to claim 11; and
- (f) an antigen presenting cell that expresses a polypeptide according to claim 2 or claim 3.

13. A method for stimulating an immune response in a patient, comprising administering to the patient a composition selected from the group consisting of:

- (a) a composition of claim 12;
- (b) a polynucleotide sequence of any one of SEQ ID NO:80-94; and
- (c) a polypeptide sequence of any one of SEQ ID NO:95-109.

14. A method for the treatment of Chlamydia infection in a patient,

comprising administering to the patient a composition selected from the group consisting of:

- (a) a composition of claim 12;
- (b) a polynucleotide sequence of any one of SEQ ID NO:80-94; and
- (d) a polypeptide sequence of any one of SEQ ID NO:95-109.

15. A method for determining the presence of Chlamydia in a patient, comprising the steps of:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with an oligonucleotide according to claim 9;
- (c) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide; and
- (d) comparing the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, and therefore determining the presence of the cancer in the patient.

16. A diagnostic kit comprising at least one oligonucleotide according to claim 9.

17. A diagnostic kit comprising at least one antibody according to claim 6 and a detection reagent, wherein the detection reagent comprises a reporter group.

18. A method for the treatment of Chlamydia in a patient, comprising the steps of:

- (a) incubating CD4+ and/or CD8+ T cells isolated from a patient with at least one component selected from the group consisting of:
 - (i) a polypeptide according to any one of claims 2 or 3;
 - (ii) a polypeptide sequence of any one of SEQ ID NO: 95-109;
 - (iii) a polynucleotide according to claim 1;
 - (iv) a polynucleotide sequence of any one of SEQ ID NO:80-94;

- (v) an antigen presenting cell that expresses a polypeptide sequence set forth in any one of claims 2 or 3;
- (vi) an antigen presenting cell that expresses a polypeptide sequence of any one of SEQ ID NO:95-109, such that the T cells proliferate; and
- (b) administering to the patient an effective amount of the proliferated T cells.

SEQUENCE LISTING

<110> Corixa Corporation
Bhatia, Ajay
Probst, Peter
Stromberg, Erika Jean

<120> COMPOUNDS AND METHODS FOR TREATMENT AND DIAGNOSIS
OF CHLAMYDIAL INFECTION

<130> 210121.515PC

<140> PCT

<141> 2001-04-23

<160> 140

<210> 1

<211> 1311

<212> DNA

<213> Chlamydia trachomatis

<400> 1

```

taattcgctt ttacctctct tcttgctgaa gacttggcta tgttttttat tttgacgata 60
aacctagtta aggcataaaa gagttgcgaa ggaagagccc taaacttttc ttatcatctt 120
ctttaactag gagtcacca tgagtcaaaa taagaactct gctttcatgc agcctgtgaa 180
cgtatccgct gatttagctg ccatcgttgg tgcaggacct atgcctcgca cagagatcat 240
taagaaaatg tgggattaca ttaagaagaa tagccttcaa gatcctacaa acaaacgtaa 300
tatcaatccc gatgataaat tggctaaaagt ttttggaact gaaaaaccta tcgatatgtt 360
ccaaatgaca aaaatggttt ctcaacacat cattaaataa aatagaaatt gactcacgtg 420
ttcctcgtct ttaagatgag gaactagtct attctttttg ttcgtttctg tgggtattac 480
tgtatcttta acaactatct tagcagcacc tgttttgaca tgggtttggg ccaatcactt 540
agagcctaac ctattgagag taacgcggtt aaattggaat ctgcctaaaa aatttgctca 600
tcttcattggg cttcgacatta tacagatttc ggatttacac ctaaacccact cgacgcctga 660
tgcctttcta aaaaaagtat ctcgtaagat ctcttctctt tctccagata ttcttgtatt 720
tacaggagac tttgtctgtc gcgctaaaagt agaaactcct gaaagattaa aacatttcct 780
atgttctctg catgcgccct taggctgttt tgcttgcta ggaatcatg attacgccac 840
ctacgtatcc cgtgatattc acgggaaaat taataccatc tcagcaatga atagccgtcc 900
tttaaaaaa gcttttacct ctgtttatca aagtctatct gcctcttctc gcaatgaatt 960
tgcagatact ctgaatccac aaattcctaa tccacaccta gtcagtatat tacgcaatac 1020
tccatttcaa ttattgcata atcaaagcgc gacactttcc gatacaatca acatcgtggg 1080
attaggcgat ttttttgcca aacaattcga tcccaaaaaa gcttttactg actataatcc 1140
cacgttacct ggtattatcc tttctcataa tcccagatcg attcaccatc tccaagatta 1200
cccaggtgat gttgtttttt ccgggcactc gcattggcct caaatctctc ttccctggcc 1260
taagtttgcc aatacgataa ccaataaact ttcagggtta gaaaaccag a 1311

```

<210> 2

<211> 1516

<212> DNA

<213> Chlamydia trachomatis

<400> 2

```

tttgagctcg tgcgctcgt gcgggtgct gtgaacgcgt tcttcaaaag cttgtcttaa 60
aagatattgt ctgccttccg gattagttac atgtttaaaa attgctagaa caatattatt 120
cccaaccaag ctctctgcgg tgcgtaaaaa acctaaattc aaaagaatga ctgcgcgctc 180
atcttcagaa agacgatccg acttccataa ttcgatgtct ttcccatg ggatctctgt 240
agggagccag ttatttgcgc agccattcaa ataagtgtcc caagcccat tgtacttaat 300
aggaacaagt tggttgacat cgacctggtt gcagttcact agacgcttgc tatttagatt 360
aacgcgtttc tgttttccat ctaaaatctc tgcttgcata agaaccgtta attttattgt 420
taatttatat gattaattac tgacatgctt cacacccttc ttccaaagaa cagacagggt 480
ctttcttcgc tctttcaaca ataattcctg ccgaagcaga cttattcttc atccaacgag 540

```

```

gctgaattcc tctcttatta atatctacaa aagatttttc aacggtcgtt gctgatgaag 600
atctcagata atacgtagtt ttcaaacctt ttttccaagc cgttaaatac atattcgaca 660
gttttttccc gtctggctgg gcaagataaa ggtagaggga ttgcccata tcaatccatt 720
tttgtcttcg agacgcgcac tgcataatcc attctgggtc aatctcaaaa gctgtoaaga 780
aaatatgttt taagtgtctt ggtatacgct cgatttccaa taaagaccca tcaaaatatt 840
tcaggtcacg taacatatca gcatcccaga tacctaattt cttcaacttc tcaattaaat 900
acacatttgg aatcgtgaat tctccggaca aattagactt cacaacaaa tgtttgtacg 960
ttggctcaat agattgagtt actcotataa tgttggagat cgtcgtgtgc ggagctatag 1020
ccataagctg acaatgtcgc ataccatgct ctttaaccaa actacggata ggttcccaat 1080
cttttcttga tgacgtatcc atctggagat ttgcttctcc tcgatagtgc gctaacaact 1140
gaatcgtatc aatagggagc aaacctctat ccattttcga tcctttataa gagctgtaag 1200
tgccctcgtc tttagcgagc agacaagaag cttgaatcgc atagtaagaa atcaactctg 1260
aactgtagtc agcaaatctt acagcttctt gcgaagcata gcttatatct agcttataca 1320
aggcatcttg gaatcccatc acccctaata caatagcgcg gtgagcaaaag ttgcgctctt 1380
tagcttccct tgttggataa aagttaatat caatcacgtt atccaacata cggactgcta 1440
tagagatcgt cttagagagt ttttctctat caaacccatc ccctacgata tgttgaacta 1500
agttaatcga tcttaa 1516

```

<210> 3

<211> 2397

<212> DNA

<213> *Chlamydia trachomatis*

<400> 3

```

agagtgtgct ggaggagcta tttttgcaaa acgggttctg attgtagata accaagaggc 60
cgttgtattc tcgaacaact tctctgatat ttatggcggc gccattttta caggttctct 120
tcgagaagag gataagttag atgggcaaat ccctgaagtc ttgatctcag gcaatgcagg 180
ggatgttgtt ttttccggaa attcctcgaa gcgtgatgag catcttcctc atacagggtg 240
ggagccattt tgtactcaa atttgacgat ttctcagaat acagggaatg ttctgtttta 300
taacaacgtg gcctgttcgg gaggagctgt tcgtatagag gatcatggta atgttctttt 360
agaagctttt ggaggagata ttgtttttta aggaaattct tctttcagag cacaaggatc 420
cgatgccatc tattttgcag gtaaaagaatc gcatattaca gccctgaatg ctacggaaag 480
acatgctatt gttttccacg acgcattagt ttttgaaaaa ctagaagaaa ggaaatctgc 540
tgaagtattg ttaatcaata tgcgagaaaa tccaggttca aaatttctca agtttgatgc 600
aattgtgcta ttcgctacct tttagtttct atgtccacgg taaagggatc ggaaagatc 660
gcattttatt tcatagtctt tagcttcgat ccctagtgtc tccgcatgga ctcgctctgc 720
aagacttttg gttacgaaaa caacaggctc tcggtgagaa atgatttgga gtactcttag 780
cgtgaggtgt tttttctgtt tctcgtgggt tgaaagattg actagaggag agacttcaat 840
acataactcg ctgccgtttt ttaataaaat ttgaccagag gagggtcttt ccgactgctc 900
tagtaataga cgaatattgc ccaatgctct ggaagcattt ttccctgatt catctcgaaa 960
ctttgcgcag gattccaatt cttcgattac tgtaaaaggg ataatgatgc gagggttaga 1020
aaaagaggaa agggccttag gatcgtaaat caaaacgctg gtatcaataa cagagggttt 1080
tttcattaca aattcctaaa tgactcaagt gtaaggggga gatagtactt tgattgtgta 1140
tcatatccag aaaaattaaa acatgtcttt gttagagaga agtcgggaga gagggttttt 1200
agcaatcaac ctccgcgtgt gctaattctg ttgtcaaaaa tgtacccctt aactacaatg 1260
ccgaggaaaag cgagtccttc tgttggaggt tgttatgaaa gtcaaaatta atgatcagtt 1320
catttgtatt tcccataca tttctgctcg atggaatcag atagctttca tagagtcttg 1380
tgatggaggg acggaagggg gtattacttt gaaactccat ttaattgatg gagagacagt 1440
ctctataccc aatctaggac aagcgattgt tgatgaggtg ttccaagagc acttgctata 1500
tttagagtcc acagctctc agaaaaacaa ggaagaggaa aaaattagct ctttgttagg 1560
agctgttcaa caaatggcta aaggatgcga agtacaggtt ttttctcaaa agggcttggt 1620
ttctatgtta ctaggaggag ctggttcgat taatatgttg ttgcaacatt ctccagaaca 1680
taaggatcat cctgatcttc ctaccgattt actggagagg atagcgcaaa tgatgcgttc 1740
attatctata ggaccaactt ctatttttagc taagccagag cctcattgca actgtttgca 1800
ttgtcaaat ggacgagcta cagtgaaga agaggatgcc ggagtatcgg atgaggatct 1860
cacttttctg tcatgggata tctctcaaag tggagaaaag atgtacactg ttacagatcc 1920
tttgaatcca gaagtatacc ttttgttttt tttatacgag ccagcactcc aatttctgac 1980
tgtgagaata tatcataaat agaccggcct ctacgcgtgc gaatagaaaa agtctttgtc 2040
atagcactat caagccttcc ctttatacgc tcaagcaata gaaacggaga tctacgcaat 2100
ggattttcat tgaatcatt aaacgagcgg aaaatgaaat tactcaaatt ttcttcagcg 2160
ctacacacgc tcaaatcatc gaggaaaacc gtatgagaaa cggatctact cgtgccgaat 2220

```

3

tcggcagcag gtctctaato ttgcagaagg agcacaaatt tttgctgtcc aagggttaaa 2280
 tactgtctga gaaataggat actgccotcc ttgccctcca gatgcgaagc atcgctatta 2340
 cttttatgct tatgcgctcg atgttgtgct ttccgatgaa gaaggagtga ccaaaga 2397

<210> 4

<211> 1094

<212> DNA

<213> Chlamydia trachomatis

<400> 4

tgatgcagaa gacactgtta agaagttaca agaagccggt gctaaggctg ttgctaaagg 60
 gctgtaattg ttatgggaaa gagaatgctt tgggggttgc ttgcaagctt ctcttttctg 120
 ttagctgcac agtagctggg cacagagggg ttcccgggtac gtcttaacag atttgtctgg 180
 acttaacttt tagtggttgg catcgcaaac agaataattc tgttgcaatg gttttttctt 240
 aatggaatca aggtgatagt atttgtcgga tggacaagtg tatagagagt atccagtgtc 300
 tctgtattgg atagactctg ttttgtccta gctggaaaagc atctgtcgta ttcotgttta 360
 gagatcacag agggactaaa tagggaaatg gtatcgccaa aagtcttaaa gtcttaggag 420
 agctcgcatg ttcaagtgcc cggagcgggt cagcgtcaaa aagaaagaag atattttaga 480
 tcttcctaata cttgtcgaaag ttcaaatcaa gtcgtataag cagtttcttc aaatcgggaa 540
 gcttgctgaa gagcgagaaa acattgggtt agaagaagtc ttcagagaaa ttttccctat 600
 caagtcttat aatgaagcta cgattttaga gtacctctct tataacttag gagtgcccaa 660
 atactcccca gaagagtgtt ttctgctggg aatcacctat agtggtactt taaaggttgc 720
 tttccgttta actgatgaaa cggggattaa agaagaagaa gtctatatgg gaaccatccc 780
 catcatgact cataagggaa cctttattat taatggggca gagagagtcg ttgtttctca 840
 agtccacogt tctccaggaa tcaattttga acaagaaaaa cattctaaag ggaatgtttt 900
 attttctttt agaattatto cttatcgagg aagttgggta gaagctgtct tcgacattaa 960
 tgaccttata tatatccata ttgataggaa aaaacgtcgc agaaagattt tagctattga 1020
 cgtttatccg agcttttaga tattcaacag atgcagatat tattgaagag ttcttttctg 1080
 tagaggagcg ttcc 1094

<210> 5

<211> 2129

<212> DNA

<213> Chlamydia trachomatis

<400> 5

gcttctttta gagataagca acaaccgagc aatccactcc tccagacata gcaacaatga 60
 tagtttttacg cacaatgagc ccagaaaacg ctttctgtta ttgaagtttg cacattacaa 120
 agggccatca tgttagcaaa aaaacaggat caaaaaaaccc tattttotcaa gccgcctctt 180
 ttaaatctta attacaaaaa taaaaatcaa ttcaactttt caaaaaaaga atttaaacat 240
 taattgttat aaaaaaataa tttattataa aataataaacc atagttgcgg ggaaatctct 300
 ttcatggttt attttagagc tcatcaacct aggcatacgc ctaaaacatt tcctttggaa 360
 gttcaccatt cgttctccga taagcatcct caaattgcta aagctatgcy gattacgggg 420
 ataaccctcg cagctctatc tctgctcgct gtagtcgcct gcgttattgc cgtctctgcy 480
 ggaggagctg ccattcctct tctgtctatt ggtggaattg ctgcaatgtc tggcctctta 540
 tccgtgcca ccattatctg ttctgcaaaa aaggctctgg ctcaacgaaa acaaaaacaa 600
 ctagaagagt tgcttccgtt agataatgcg accgagcatg tgaattacct gacctcagac 660
 acctcttatt ttaatcaatg ggaatcctta gatgctctaa ataagcagtt gtctcagatt 720
 gacttaacta ttcaagctcc cgaaaaaaaa ctattaaaag aagttcttgg ttccagatac 780
 gattccatta atcaatccat cgaagagatc tccgatcgct ttacgaaaaa gctctctctt 840
 cttcgattaa gagaacattt ttgtcgagga gaagagcggt atgcccccta ttaagccct 900
 cctctactta acaagaatcg tttgtgacc caaatcacat ccaatatgat taggatgcta 960
 ccaaaatccg gtggtgtttt ttccctcaaa gccaatcacac taagtcatgc cagccgcaca 1020
 ctatatagc atttgaaagt cgttttatcc ttaggagtto tgcgtggagt cgtgctctt 1080
 atcatctttc ttcccccctag cctgcctttt atcgctgtta taggagtatc ttcccttagca 1140
 ttggggatgg catcttttct tatgattcgg ggcattaaat atttgcctga acattctctc 1200
 ctgaatagaa agcaattagc taaagatatt caaaaaacca ttatcccaga tgtcttggcc 1260
 tctatggttc attaccagca tcaattacta tcacatctac atgaaactct attagatgaa 1320
 gccatcacag ctagatggag cgagcccttc tttattgaac acgctaactc taaggcaaaa 1380
 attgaagatt tgacaaaaca atatgatata ttgaacgcag cttttaataa atctttacaa 1440
 caagatgagg cgctccgttc tcaattagag aaacgagctt acttattccc aattcctaata 1500

```

aacgcagaaa atgctaaaac taaagaatcg cagcttctag actcagaaaa tgattcaaat 1560
tctgaatttc aggagattat aaataaagga ctagaagctg ccaataaacg acgagctgac 1620
gctaagtcaa aattctatac ggaagacgaa acctctgaca aaagattctc tatatggaaa 1680
cccacaaaaga acttggcatt agaagatttg tggagagtgc atgaagcttg caatgaagag 1740
caacaagctc tcctcttaga agatttatatg agttataaaa cctcagaatg tcaagctgca 1800
ctccaaaaag tgagtcaaga actgaaggcg gcacaaaaat cattcgcagt cctagaaaag 1860
catgctctag acagatctta tgaatccagt gttagccatga tggatttagc tagagcgaat 1920
caagaaacac accggcttct gaacatctc tctgaattac aacaactagc acaataacctg 1980
ttagataatc actaacgggt cttcataaat gacaaaaaga aaaaggagag ctgttgctgt 2040
gctctccttt ttctctaaat attcctgaaa gactaacctt tttatggttg cgttgagcct 2100
cctcctcctg ttcccgagga gcccgcaac 2129

```

<210> 6

<211> 1828

<212> DNA

<213> Chlamydia trachomatis

<400> 6

```

gagggagcag cctaactctc cctctctctc ttaaaaaaga ggggagcctt ttttccttac 60
aaagatacgc tagctttttc ctgaagaatc tcatcaagag atattttgcat tttcccacgg 120
ataaaggcat cccaagggaag ccttggaaac acttcatatt ctcccgttgc tagcatttga 180
caagggaaac caaagattaa atcttccggt aatccatagg gattgtggtc cgaacacact 240
ccggaagaaa accattctcc ttcttttggc tgatatattg atcagagcagc ctctgctaaa 300
gctcgtgctg cagaagctgc cgaagacttc cctcgtgctt cgattactgc actaccacga 360
ctctgtacag aaggcaccat aatattctct aaccaatcac gatccgctat cgtctctgcg 420
ataggacggt cattaatcag agcttgcgta aaatcaggca cttgtttggc ggagtgattt 480
ccccaaacca caacttgtga tacagccgat aaaggtactt ctgctctatg cgataacatg 540
ctatgcatac gattctgggt caatcgtagc atcgcatgaa agttctttct caataatctg 600
ggagcatgat tcattgctat ccagcaattg gtattcacag ggttcccaac aacaaaaatc 660
tttgcattcc gcttggctgt tgtgttcaaa gcttttctct gcgtagcaaa aatctcccca 720
tttttcttta gaagatccct tctctccatt cctgggcctc taggaactga cctataagg 780
aatgccgcat caatgccatc aaaagcatca tgcaatgatg tcgttacctg cacacgctgt 840
aataaaggga aagcaccatc atctagctcc atgcgcacac cagataaagc ctttctgtt 900
ccaggaatat cgtagatacg cagatcgatg ccacaatcaa ggccaaaaac atctccatga 960
gccagagaaa atagaaagct ataggctatt tgccctgttc ctctgtttac tgctacactc 1020
actgtttgag aaaccataag ccacctctc tttactttta caaaacgcac atactctcaa 1080
cactacgttt gcaactaact aattttggtc ccaacatacg tttggatgat aaaagaatca 1140
agtacctaga ttcttagta aaagcttttg gcaaaaaaaa gctcatctat ttttcaatag 1200
atgagccgag tttaactgaa taagaactta gaaaacttta taaaaaatag gcccggtgta 1260
tctaccatc atacttgatc ccgaccgat aacttgttgt ccctttttag cagccaaata 1320
accgtggaca tctaaaaaac caataaacgg tgcgcgaata aagaacataa agcccctaaa 1380
aaaacgattt taagagagaa gtaatagaca gattgttaaca tatttaaaat aaaaactctg 1440
caaacaaaaa aactttgcct ggccgtctcc gtagaaagca ctttatgtta aaacgttaaa 1500
aagtcctaac atacctcgag cttcgggaaa ctctacagga gcattcccg acatgatgcc 1560
tataatttgc gttgccaat ctctccctaa tgaaaccct tcttgatcaa aagaattgat 1620
tccccagcaa aacccttgaa atgcaaatt atgctcataa aaagccaata aactaccagc 1680
aatacgagga gaaagctgtt gcgctaccaa tatogaagaa ggtctgttcc ctttaaacct 1740
cttattcggg ttgcgattat ctctaccctg agctaaagct aaagattgag caacaagggt 1800
tgcaaaagac ttttgagatc togtgccg 1828

```

<210> 7

<211> 861

<212> DNA

<213> Chlamydia trachomatis

<400> 7

```

gggcgcacta ctttaagat tgcgtgcct tttggtacta cgagagaagt tcgtgtgaaa 60
tggcgttatg ttctgaagg tgtaggagat ttggctacca tagctccttc tatcagggtc 120
ccacagttac agaaatcgat gagaagcttt ttccctaaga aagatgatgc gtttcatcgg 180
tctagttcgc tattctactc tccaatggtt ccgcattttt gggcagagct tcgcaatcat 240
tatgcaacga gtggtttgaa aagcgggtac aatattggga gtaccgatgg gtttctccct 300

```

```

gtcattgggc ctgttatatg ggagtcggag ggtcttttcc gcgcttatat ttcttcgggtg 360
actgatgggg atggtaagag ccataaagta ggattttctaa gaattcctac atatagttgg 420
caggacatgg aagattttga tccttcagga ccgcctcctt gggaagaatt tgctaagatt 480
attoaagtat tttcttctaa tacagaagct ttgattatcg accaaacgaa caaccagggt 540
ggtagtgctc tttatcttta tgcactgctt tocatgttga cagaccgtcc tttagaactt 600
cctaaacata gaatgattct gactcaggat gaagtgggtg atgctttaga ttggttaacc 660
ctgttggaaa acgtagacac aaacgtggag tctcgccctg ctctgggaga caacatggaa 720
ggatatactg tggatctaca ggttgccgag tatttaaaaa gctttggacg tcaagtattg 780
aattgttggg gtaaagggga tatcgagtta tcaacgccta ttcctctttt tggttttgag 840
aagattcctc cacatcctcg a                                     861

```

<210> 8

<211> 763

<212> DNA

<213> Chlamydia trachomatis

<400> 8

```

ataacaaaaa catcttgatt atttttgtta aaagaaatac ttaatgagtt ttatttaatt 60
aacgaaacga aaagcttgct aatgaaaatt attcacacag ctatcgaatt tgctccggta 120
atcaaagcgg gaggcctggg agacgcgcta taoggactag caaaagcttt agccgctaatt 180
cacacaacgg aagtgttaat ccttttatac cctaaattat ttactttgcc caaagaacaa 240
gatctttgct cgatccaaaa attatcttat ttttttgctg gagagcaaga agcaactgct 300
ttctcctact tttatgaagg aattaaagta actctattca aactcgacac acagccagag 360
ttattcgaga atgcggaaac aatctacaca agogatgatg ccttccggtt ttgcgctttt 420
tctgctgctg cggcctccta catccaaaaa gaaggagcca atatcggttca ttacacgat 480
tggcatacag gattagttgc tggactactc aaacaacagc cctgctctca attacaaaag 540
attgtttctta ccctacataa ttttggttat cgaggctata caacacgaga aatatttaga 600
gcctcctctt tgaatgaatt ttatatcagc cagtaccaac tatttcgoga tccacaaact 660
tgtgtgttgc taaaaggagc tttatactgt tcagatttgc tgactacggt ttctcctaca 720
tacgccaaaag aaattcttga agattattcc gattacgaaa ttc                                     763

```

<210> 9

<211> 665

<212> DNA

<213> Chlamydia trachomatis

<400> 9

```

ttgaaactaa aaacctaat tattttaaagc tcaaaataaa aaagagtttt aaaatgggaa 60
attctgggtt ttatttgat aacactgaaa actgcgtctt tgctgataat atcaaagttg 120
ggcaaatgac agagccgctc aaggaccagc aaataatcct tgggacaaca tcaacacctg 180
tcgcagccaa aatgacagct tctgatggaa tatctttaac agtctccaat aattcatcaa 240
ccaatgcttc tattacaatt ggtttggatg cggaaaaagc ttaccagctt attctagaaa 300
agttggggaga tcaaattctt gatggaattg ctgatactat tgttgatagt acagtccaag 360
atatttttaga caaaatcaaa acagaccctt ctctaggttt gttgaaagct tttacaact 420
ttccaatcac taataaaatt caatgcaacg ggttattcac toccagtaac attgaaactt 480
tattaggagg aactgaaata ggaaaattca cagtcacacc caaaagctct gggagcatgt 540
tcttagtctc agcagatatt attgcatcaa gaatggaagg cggcgttggt ctagcttttg 600
tacgagaagg tgattctaag ccctgcgcga ttagttatgg atactcatca ggcatttcta 660
attta                                     665

```

<210> 10

<211> 843

<212> DNA

<213> Chlamydia trachomatis

<400> 10

```

tggaatgtc gaagaatacg attacgttct cgtatctata ggacgccgtt tgaatacaga 60
aaatattggc ttggataaag ctggtgttat ttgtgatgaa cgcggagtca tccctaccga 120
tgccacaatg cgcacaaacg tacctaacat ttatgctatt ggagatatca caggaaaatg 180
gcaacttgcc catgtagctt ctcatcaagg aatcattgca gcacggaata tagctggcca 240
taaagaggaa atcgattact ctgccgtccc ttctgtgatc tttaccttcc ctgaagtcgc 300

```

```

ttcagtaggc ctctcccca cagcagctca acaacaaaa atccccgtca aagtaacaaa 360
attcccattt cgagctattg gaaaagcggg cgcaatgggc gaggcgatg gatttgacgc 420
cattatcagc catgagacta ctcagcagat cctaggagct tatgtgattg gccctcatgc 480
ctcatcactg atttccgaaa ttaccctagc agttcgtaat gaactgactc ttccctgtat 540
ttacgaaact atccacgcac atccaacctt agcagaagtt tgggctgaaa gtgcgttgtt 600
agctgctgat accccattac atatgcccc tgctaataaaa tgaccgattc agaatctcct 660
actcctaaaa aatctatacc cgccagattc cctaagtggc tacgccagaa actcccttta 720
gggcgggtat ttgctcaaac tgataatact atcaaaaaata aagggtcttc tacagtctgt 780
gaggaagcct cttgtccgaa tcgcacccat tgttgggtcta gacataacgc gtacctatct 840
agc

```

<210> 11

<211> 1474

<212> DNA

<213> Chlamydia trachomatis

<400> 11

```

acagaaggga cggcagagta atcgatttcc tctttatggc cagctatatt ccgtgctgca 60
atgattcctt gatgagaagc tacatgggca agttgccatt ttcctgtgat atctccaata 120
gcataaatgt taggtacgtt tgtgcgcatt gtggcatcgg tagggatgac tccgcgttca 180
tcacaaataa caccagcttt atccaagcca atattttctg tattcaaacg gcgtcctata 240
gatacgagaa cgtaatcgta ttcttcgaca ttccattga tagttaaccg aacgcgatct 300
cctatatcct caatatttga tacagaggct tctagtacga aacggagtcc ttgtcgggtg 360
aatttatoga acatggtttt tgaaatatct ggattattca aagcaaggat ttgagagcct 420
gcttcgatca cagaaacttc ggagcctaac gtatggaata aggaagcgaa ttcgcaaccg 480
atcacaccac cgccaataat ggccattttt tgagggattt ctttgagggt tagcacgcct 540
gttgagcata aaatccgagg agattctgcg gaaaaaggaa tccccgggaa agctcgttgt 600
tcagagccgg tggctaggat aatggagtgc gctttgatta cagaagggtt ttctcctaag 660
atttttactt ctgttgaaga gatcaaagag cctcttccag agaagacagt gatcttattg 720
ctgcgaatga gaccattaag tccatcgcg gtgctacgga ctacggaatc ctctcttgtt 780
accatagogg gatagttgat gctgaatcct tctacatgaa tcccaaacgt gtcagcatgg 840
cgtatttggg taacgacttc agctcctgct aagagggtt tagaaggaat acaccctcgg 900
tttaaacagg ttccgccagc ctctcgcttt tcgattagcg cagttttgag tcttgcttga 960
gcggcagtga ttgtcgcaac atagcctcct ggccccgctc cgataactac acagtcgaaa 1020
gcttcattca taacatttcc tcttcaatga gtgttttaga ttgcaacgat ccatatgaga 1080
tgattatctg aaggaagagg attctccttc caagcctttc taggaaaggg aaagagaggt 1140
ccttcagaca aatacatttc ccggattgta catctgggtg gataaaatct caatgaggag 1200
aagtggtagc aggagagaaa aaataggaac gtaagagtgt tatttcgaat gctcaggagg 1260
agagcggtag ccacgataag caagcagaat cccgactagt gcatagatgt atgagcgatt 1320
ctttggccag gagagaacga gtccagagcc tgtcgaaaac aagagaatca tgagcgaaaa 1380
ggtaaggaaa ccgcaacca agaagagagc tgcagtcggc caatattgta gccagtccca 1440
ctgggagggg gcaggctctt gaacaggctc ctca

```

<210> 12

<211> 2017

<212> DNA

<213> Chlamydia trachomatis

<400> 12

```

ataagcattc tcatctaccc agaagtagaa gtcaaaacct tcataagtat ctaaaaagac 60
tcgcataata tcttcgatac catccggagg cgctcctgcg atccatattc catggatgat 120
tttctcaaca ggtacacgat ggccctttaa ttctgttttg atggtttcaa gaacaccttc 180
aatcggagtc gtcttaggtt tttcttcggc tttctgttcc ttagcttttg cctgtttagg 240
ctgagcctgc gatgatgctg gaagcttctt ctgaatggca tcgacgtatt ttcttgggtg 300
aatcaaggaa ttctgtcccg cttccgaatt tttatctggc atagagttgt aagcactaat 360
gaccttttct agtttattta ataggtcttt aacagtagat ttctgttcag gagtaattcc 420
tagtttttct tctatgttct tgggagtaag atcgtatttg ctagcatcaa gattttctat 480
ctttccagaa gaagcttctc cttcttctc tctatagca cgcttttttc tcgataaaac 540
agctgctgta ggaggaactg cactagcaga aatcgttttt acccccccct ctctgaacag 600
agtacgtaog aacgttcact ggctgtgtaa taaacttcgt ctttctctta cgaggagagg 660
ttttgtcgtt acttctgtt ctttagagat tgtagtgaac ttattctctg aagtagaagt 720

```



```

ctctgccgtc tcgtgccgaa ttccggcacga gaagccatgt tatcttttgc tagatcaatg 780
ccttcttggt ttttgaattc atcaagcatc cagttgatga tgactccgtc gaagtcgtct 840
cctcccaagt gagtatcccc gttggttgag agaacttcaa aaactccgtc accgatttcc 900
aagatagaaa tatcgaaagt tctctctcct aagtcgaaga cggcgatttt tttatctcct 960
tccttatcaa taccataagc aagagcggcc gctgttggtt caggaataat gcgtttaaca 1020
tctaattcct cgatacgtcc agcatotttt gtagaagctc tttgagaatc gttaaagtaa 1080
gctggtacgg taatgactgc ttccggttact gtttctccga gataagcctc agcagtttcc 1140
ttcatcttca tgaggatctg agcgcggatt tcttctggag tgtacagttt ttgttccaca 1200
tcaaagaccg catctccttt cgagtttaga gcaactttgt aggggactgt ttttaatttc 1260
gattcgactt cagagaattt totaccgatg aatcgcttag tagaagccaa tgttttttca 1320
ggattggtta ctgcctgacg ttttgaggga attccaacaa gagtttgcgc accttataaa 1380
gcaacgatag aaggagtagt acgagttcct tcagaagagg caataacttt aggttggcca 1440
ccttccataa cagagacgca agagttggtc gtccctaggc cgataccaat aattttgtta 1500
gactttcttt tttcgtctat attgaacacc taatttctag gataattatt ctttttcttc 1560
gttaccgtct gagtttcttt tagcaggaag ttttgctact ttcaactttg ctacgcgaat 1620
aggacgatct cctatcttat aacctttagt aaattcctcc aagatagtc cttctggaat 1680
tggttggtt tcttcgattt ctacagcttc atgcaggtac ggattaaata gttctccttt 1740
cgaggaatat tcaaccacac ctttctcttc gaagatttgc ttaaattggt gaaggatcat 1800
ttggaatcct atagcccaat tttttacttc ttcagaggtt tgagaagcga atcccaaacg 1860
cttttccata ctttcgatag aaggaaggaa atccataaga gcattttcta cagcactctg 1920
catcatttct gtgcgttctt tctgtagtcg ttttcttgag ttttctgctt cagcgagagc 1980
catcagatat cgatcattct gttcttgcct cgtgccg 2017

```

<210> 13

<211> 1171

<212> DNA

<213> Chlamydia trachomatis

<400> 13

```

ggtaaacgag ttaaaacaag agcatacag gctaacggac tcgcctttag tgaaaaaagc 60
tgaggagcag attagtcaag cacaaaaaga tattcaagag atcaaaccta gtggttcgga 120
tattcctatc gttggtccga gtgggtcagc tgcttccgca ggaagtgcgg caggagcgtt 180
gaaatcctct aacaattcag gaagaatttc cttgttgctt gatgatgtag acaatgaaat 240
ggcagcgatt gcaactgcaag gttttcgatc tatgatcgaa caatttaatg taaacaatcc 300
tgcaacagct aaagagctac aagctatgga ggctcagctg actgcgatgt cagatcaact 360
ggttggtgag gatggcgagc tcccagccga aatacaagca atcaaagatg ctcttgcgca 420
agctttgaaa caaccatcag cagatggttt ggctacagct atgggacaag tggcttttgc 480
agctgccaaag gttggaggag gctccgcagg aacagctggc actgtccaga tgaatgtaaa 540
acagctttac aagacagcgt tttcttcgac tcttccagc tcttatgcag cagcactttc 600
cgatggatat tctgcttaca aaacactgaa ctctttatat tccgaaagca gaagcggcgt 660
gcagtcagct attagtcaaa ctgcaaatcc cgcgctttcc agaagcgttt ctogttctgg 720
catagaaagt caaggacgca gtgcagatgc tagccaaaga gcagcagaaa ctattgtcag 780
agatagccaa acgttaggtg atgtatatag ccgcttacag gttctggatt ctttgatgtc 840
tacgattgtg agcaatccgc aagcaaatca agaagagatt atgcagaagc tcacggcctc 900
tattagcaaa gctccacaat ttgggtatcc tgctgttcag aattctgcgg atagcttgca 960
gaagtttgct gcgcaattgg aaagagagtt tggtgatggg gaacgtagtc tcgcagaatc 1020
tcaagagaat gcgtttagaa aacagcccgc tttcattcaa caggtgttgg taaacattgc 1080
ttctctatct tctggttctt tttcttaacg tgtgattgaa gtttgtgaat gagggggagc 1140
caaaaaagaa tttctttttt ggtctttttt t 1171

```

<210> 14

<211> 877

<212> DNA

<213> Chlamydia trachomatis

<400> 14

```

cagagaattc tcgacatact atctaactcg atatgtaag ctgctttaca tcccttgaa 60
tagaaaataa atggaaataa aaagcccaga acaagagaag ttgttctggg ctgacagaag 120
ctgtcagatc attttaataa gattgatgac aactacgaca agttcctgga tccaaaaaag 180
aatctaaaaa gccatacaaa gattgcgtta cttcttgoga tgctctaac actttatcag 240
cgctcatctt gagaagcatc tcaatgagcg ctttttcttc tctagcatgc cgcacatccg 300

```

```

cttcttcatg ttctgtgaaa tatgcatagt cttcaggatt ggaaaatoca aagtactcag 360
tcaatccacg aatttttctct ctagcgatac gtggaatttg actctcataa gaatacaaaag 420
cagccactcc tgcagctaaa gaatctcctg tacaccaccg cacgaaagta gctactttcg 480
cttttgctgc ttacttaggc tcatgagcct ctaactcttc tggagtaact cctagagcaa 540
acacaaactg cttccacaaa tcaatatgat tagggtaacc gttctcttca tccatcaagt 600
tatctaacaa taacttacgc gcctctaaat catcgcaacg actatgaatc gcagataaat 660
atthaggaag ggcttttgata tgtaaataat agtctttggc atacgcctgt aattgctctt 720
tagtaagctc ccccttogac catttcacat aaaacgtgtg ttctagcata tgcttatttt 780
gaataattaa atctaactga tctaaaaaat tcataaacac ctccatcatt tcttttcttg 840
actccacgta accgcttgca aaaaagggtc gtataag 877

```

<210> 15

<211> 396

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 15

```

tgtaccaa atgagcttag atcaatctgt tgttgaactt tacacagata ctgccttctc 60
ttggagcgtg ggcgctcgag cagctttgtg ggagtgcgga tgtgcgactt taggggcttc 120
tttccaatc gctcaatcta aacctaaagt cgaagaatta aacgttctct gtaacgcagc 180
tgagtttact atcaataagc ctaaaggata tgtagggcaa gaattccctc ttgcactcat 240
agcaggaact gatgcagcga cgggcactaa agatgcctct attgattacc atgagtggca 300
agcaagttta gctctctctt acagattgaa tatgttcaact ccctacattg gagttaaatg 360
gtctcgagca agttttgatg ccgatacgat tcgtat 396

```

<210> 16

<211> 516

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 16

```

ctcaaaat ttt gacgatttct cagaatacag ggaatgttct gttttataac aacgtggcct 60
gttcgggagg agctgttctg atagaggatc atggtaatgt tcttttagaa gcttttgag 120
gagatattgt ttttaaagga aattcttctt tcagagcaca aggatccgat gccatctatt 180
ttgcaggtaa agaatcgcat attacagccc tgaatgctac ggaaggacat gctattgttt 240
tccacgacgc attagttttt gaaaatctag aagaaaggaa atctgctgaa gtattgttaa 300
tcaatagtct agaaaatoca ggttacctg gatctattct atttttagaa gcagaaagta 360
aagttctcca atgtattcat gtacaacaag gaagccttga gttgctaaat ggagctacat 420
tatgtagtta tggtttttaa caagatgctg gagctaagtt ggtattggct tctggatcta 480
aactgaagat tttagattca ggaactcctg tacaag 516

```

<210> 17

<211> 723

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 17

```

ctccttttaa gggggacgat gtttacttga atggagactg cgcttttctc aatgtctatg 60
caggggcaga gaacggctca attatctcag ctaatggcga caatttaacg attaccggac 120
aaaaccatac attatcattt acagattctc aagggccagt tcttcaaaat tatgccttca 180
tttcagcagg agagacactt actctgaaag atttttcgag tttgatgttc tcgaaaaatg 240
tttcttgagg agaaaaggga atgatctcag ggaanaacct gagtatttcc ggagcaggcg 300
aagtgaattt ttgggataac tctgtggggt attctccttt gtctattgtg ccagcatcga 360
ctccaactoc tccagcacca gcaccagctc ctgctgcttc aagctcttta tctccaacag 420
ttagtgatgc tcggaaaggg tctatttttt ctgtagagac tagtttggag atctcaggcg 480
tcaaaaaagg ggtcatgttc gataataatg ccgggaattt tggaacagtt ttctgaggta 540
atagtaataa taatgctggg agtgggggta gtgggtctgc tacaacacca agttttacag 600
ttaaaaaactg taaaggggaa gtttctttca cagataacgt agcctcctgt ggaggcggag 660
tagtctacaa aggaactgtg cttttcaaaag acaatgaagg aggcataatc ttccgaggga 720
aca 723

```

<210> 18
 <211> 1377
 <212> DNA
 <213> Chlamydia trachomatis serovar E

<400> 18
 aaacagctaa tcgtcactac gctcacgtgg actgccctgg tcacgctgac tatgttaaaa 60
 acatgatcac cgggtgggct caaatggacg gggctattct agtagtttct gcaacagacg 120
 gagctatgcc tcaaaactaaa gagcatattc ttttggcaag acaagttggg gttccttaca 180
 tcgttggttt tctcaataaa attgacatga tttccgaaga agacgctgaa ttggtcgact 240
 tggttgagat ggagttggct gagcttcttg aagagaaaag atacaaaggg tgtccaatca 300
 tcagagggtc tgctctgaaa gctttggaag gggatgctgc atacatagag aaagttcgag 360
 agctaatagca agcgcgcgat gataatatcc ctactccaga aagagaaatt gacaagcctt 420
 tcttaatgcc tattgaggac gtgttctcta tctccggacg aggaactgta gtaactggac 480
 gtattgagcg tgggaattgtt aaagtttccg ataaagtcca gttggtcggg cttagagata 540
 ctaaaagaaac gattgttact ggggttgaaa tgttcagaaa agaactocca gaaggtcgtg 600
 caggagagaa cgttggattg ctctcagag gtattggtaa gaacgatgtg gaaagaggaa 660
 tggttgtttg cttgcaaaac agtggttaaac ctcatacaca gtttaagtgt gctgtttacg 720
 ttctgcaaaa agaagaaggt ggacgcacata agcctttctt cacaggatat agacctcaat 780
 tcttcttccg tacaacagac gttacaggtg tggtaactct gcctgaggga gttgagatgg 840
 tcactcctgg ggataacggt gagtttgaag tgcaattgat tagcctgtg gctttagaag 900
 aaggtatgag atttgcgatt cgtgaaggtg gtcgtacaat cgggtcgtga actatttcta 960
 agatcattgc ataaattaag tgatgtgttg gcgaggctga aaagccttgc ctttgggtgt 1020
 gtagcttaga tggtagagca gtggcctcca aagccgcggg tcgggggttc gaatccctcc 1080
 gcactcgtat taggtaactg aaagaagaat togtttatgg ggcaagatca ccgaagaaaa 1140
 tttcttaaga aagtatcttt tgcaaaaaaa caagcagctt ttgcgggtaa ctttatcgaa 1200
 gaaattaaga agattgagtg ggtaaataag cgaaatctta aaagatacgt caagattgtt 1260
 ttgatgaata tttttggcct tggattttcc atctattgtg tggatttagc tcttcgaaag 1320
 tccctttcat tgttcggtaa agtaacaagc tttttctttg gttgattcat gtttaag 1377

<210> 19
 <211> 1736
 <212> DNA
 <213> Chlamydia trachomatis serovar E

<400> 19
 gtagcggaac aaagccggac cagcaggcct catagaatat aaaaatcacga ggagcttaaa 60
 catgtcagat caagcaacga ccctcaagat taaacctttg ggagatagaa ttttagttaa 120
 aagagaagaa gaagcttcca ctgcaagagg cggatcattt ctctctgaca ctgccaagaa 180
 aaagcaagat agagctgaag ttttagctct aggaacaggc aaaaaagatg ataaaggcca 240
 gcaacttccct tttgaagttc aggttgggtga catcgtttta attgataaat attctggcca 300
 agaactcact gtagaaggtg aagagtacgt catcgttcaa atgagcgaag ttatcgcagt 360
 tctgcaataa aaactaagag agtgaagtaa gatttaaggg agcgcaccaa tgggtcgctaa 420
 aaacattaaa tacaacgaag aagccagaaa gaaaattcaa aaaggagtta agacttttagc 480
 tgaagctgta aaagtcaact tagggcctaa aggacgacat gttgtcatag ataaaagcctt 540
 cggatccctc caagtaacta aagatgggtg taccgttgcg aaagaagttg agcttgccga 600
 caaacatgaa aatatgggag ctcaaatggt caaagaagtc gccagcaaaa ctgctgacaa 660
 agctggagac ggaactacaa cagctactgt tcttgctgaa gctatctata cagaaggatt 720
 acgcaatgta acagctggag caaatccaat ggacctcaaa cgaggatttg ataaagctgt 780
 taaggttgtt gttgatcaaa tcagaaaaat cagcaaacct gttcagcatc ataaagaaat 840
 tgctcaagtt gcaacaattt ctgctaataa tgatgcagaa atcgggaatc tgattgctga 900
 agcaatggag aaagttggta aaaacggctc tatcactgtt gaagaagcaa aaggatttga 960
 aaccgttttg gatgttgggt aaggaatgaa tttcaataga ggttacctct ctagctactt 1020
 cgcaacaaat ccagaaactc aagaatgtgt attagaagac gctttggttc taatctacga 1080
 taagaaaatt tctgggatca aagatttccct tctgtttta caacaagttg ctgaatccgg 1140
 ccgtctctct ctattatag cagaagacat tgaaggcgaa gctttagcta ctttggctgt 1200
 gaacagaatt cgtggaggat tccgggtttg cgcagttaaa gctccaggct ttggagatag 1260
 aagaaaagct atgttggag acatcgcctat cttaactggc ggtcaactca ttagcgaa 1320
 gttgggcctg aaattagaaa acgctaactt agctatgtta ggtaaaagcta aaaaagttat 1380
 cgtttctaag gaagacacga ccactcgtcg aggaatgggt gaaaaagaag ctttagaagc 1440
 tcgttgcgaa agcatcaaaa aacaaattga agacagctct tctgattacg ataaagaaaa 1500

10

```

actccaagag cgtcttgcga agctctctgg tggagtagca gtcattcgcg ttggagctgc 1560
aacagagatt gagatgaaag agaaaaaaga tcgtgtagac gatgctcaac atgctacaat 1620
cgctgctggt gaagaaggaa ttcttcctgg tggaggaaca gcattaatcc gttgtatccc 1680
tactcttgag gccttcttgc caatgttgac taatgaagat gagcaaattg gagctc 1736

```

<210> 20

<211> 1135

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 20

```

ggctcttgat gaaaaagagc ggcagggttat ggctctttat tactatgatg acttgggtatt 60
aaaagaaatt gggaagattt taggagtgag cgagtcccga gtttctcaga tacactocaa 120
agcttttatg aagttacgag gtacattgtc cagtctgctt tagtaactgt ctccagaaga 180
tcctctttgt atttttccta tcaatattct attggagaag cgcgtcggtt ttttgacgag 240
gtgtctgcta tcgcttgccg tgctataaaa agaacaggat agataagatg ttgctagata 300
agttttatag gatagatttt tatgcaacag ttaatcgata accttaagaa acgggggtatt 360
ctagataatt cttctgcagg attagaaact cgtgccgaag tttgtggaga agagaaagaa 420
atctctctag cagactttcg tggtaagtat gtagtgctct tcttttatoc taaagatttc 480
acctatgtgt gtcctacaga attgcatgct tttcaagata gattggtaga ttttgaagag 540
cggggtgcag tcgtgcttgg ttgctccgtt gacgacattg agacacattc tcgttggctc 600
gctgtagcga gaaatgcagg aggaatagag ggaacagaat atcctctggt agcagacctc 660
tcttttaaaa tatcagaagc ttttgggtgt ttgaatcctg aaggatcgct cgctttaaga 720
gcgactttcc ttatcgataa acatgggggt gttcgctcatg cggttatcaa tgatcttcc 780
ttagggcggt ccattgacga ggaattgcgt attttagatt cattgatctt ctttgagaac 840
cacggaatgg tttgtccagc taactggcgt tctggagagc gtggaatggt gccttctgaa 900
gagggattaa aagaatattt ccagacgatg gattaagcat ctttgaagat aagaaagtcg 960
tacagatctt gatctgaaa gagagaagg ctttttaatt ttctgcagag agccagcgag 1020
gcttcaataa tgttgaagtc tccgccacca ggcaatgcta aggcgatgat attagttagt 1080
gaaatctgag tgtttaaggaa ataaaggcca aagaagtagc tatcaataaa gaagc 1135

```

<210> 21

<211> 731

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 21

```

ttgaagacac tctttctccc ggagtcacag ttcttgaagc tgcaggagct caaatttctt 60
gtaataaagt agtttggact gtgaaagaac tgaatcctgg agagtctota cagtataaag 120
ttctagttag agcacaact cctggacaat tcacaaataa tgttgttgtg aagagctgct 180
ctgactgtgg tacttgtact tcttgccgag aagcgacaac ttactggaaa ggagttgctc 240
ctactcatat gtgcgtagta gatacttgtg accctgtttg tgtaggagaa aatactgttt 300
accgtatttg tgtcaccaac agagggttctg cagaagatac aaatgtttct ttaatgctta 360
aattctctaa agaactgcaa cctgtatcct tctctggacc aactaaagga acgattacag 420
gcaatacagt agtattcgat tcgttaccta gattaggttc taaagaaact gtagagtttt 480
ctgtaacatt gaaagcagtt acagctggag atgctcgtgg ggaagcgatt ctttcttccg 540
atacattgac tgttccagtt tctgatacag agaatacaca catctattaa tctttgattt 600
tatcgatgtg taggtgccc ccagggattc ctgggcccgt tttttttgtt atctatatga 660
aaataaaaga gttcattttt ggtctcagag catattctag acgggttttt gaaaaaata 720
agtgtttgtg t 731

```

<210> 22

<211> 1181

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 22

```

ctatcgtctg aatgctgaac tgaacatctt ttttgattta gacgcgttag ccgatgctat 60
ggatctatct cgagatctac agttttctta catgggtatt caaaatctgt atgatcggtta 120
ttttaatcac cacgaagatt gccgtttaga aactcccaa attttttggg tgcgcgttgc 180
tatgggggtg gcattgaatg agcaagacaa gacttcttgg gctattactt tttataattt 240

```

```

gctttcgaca ttccgatata caccagctac gccaaccttg ttcaattcag gtatgcggca 300
ttctcagtta agctcttgot atctttccac tgtacaagat aatttgggtca atatctataa 360
ggtcattgct gataacgcta tgctatctaa gtgggcagga gggataggtat atgattggac 420
ggcgggtcgt gcaacagggg ctttaattaa aggaaccaat ggaagaagtc agggagtaat 480
tccttttatt aaggtgacaa atgatacagc agtcgcagtg aatcaagggtg gtaaacgcaa 540
gggagctgta tgcgtctatt tagaagtttg gcacctcgac tacgaagatt tccttgaatt 600
gagaaagaat acaggggatg agcgtcgacg ggctcatgat gtcaatatag ctagctggat 660
tccagatctt ttcttcaaac gtttacagca aaaagggaca tggactctat tcagcccaga 720
tgatgttccg ggattacacg atgcttatgg ggaagaattt gagcgtttgt acgaagaata 780
tgagcggaaag gttgataccg gagagattcg gttattcaag aaggtagaag ctgaagatct 840
gtggagaaaa atgctcagca tgctttttga aacgggacac ccatggatga cttttaaaga 900
tccatccaac atcgttccg ctcaagatca taaaggcgtg gtgcgttgtt ccaatctgtg 960
tacggagatt ttgttaaact gctcggagac agaaactgct gtttgtaatt taggatcgat 1020
taacttagtt caacatatcg taggggatgg gtttagatgag gaaaaactct ctgagacgat 1080
ctctatagca gtccgtatgt tggataacgt gattgatatt aacttttatc caacaaagga 1140
agctaaagga gcgaactttg ctcaaccgcg tattggatta g 1181

```

<210> 23

<211> 167

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 23

```

ttaaaaagat tttaaactaa aaagaagatt ttttaattata gtttttcaaa atcatttttga 60
tatttttaat gctgagataa acaagaaaag cggaaaactcc ttgcgacaaa gattttctgc 120
tcgagccctc ttccctgagg atttttttagg ggagatccat tcttcca 167

```

<210> 24

<211> 1265

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 24

```

caggttcttt ctagacgaac aaagaataat cctatgttga tagggggagcc cggagttggg 60
aaaacagcaa tcgtgaagg acttgctctt cgcatagtgc aaggggatgt tccagagagt 120
ttaaaggaaa agcatctgta tgtactggat atgggagctt tgattgcagg tgccaagtat 180
cgaggagagt ttgaagagcg gttaaaaagt gtattgaagg gtgtagaagc ttctgaaggc 240
gagtgtatcc tattcattga tgaagtgcac acttttagtag gagcgggagc tacagatgga 300
gctatggatg cagcgaatct attaaagcct gcttttagcac gaggcacttt gcattgtatt 360
ggcgctacga ctttgaatga atacaaaaaa tatatagaga aagacgcggc tttggaacgg 420
cgtttccagc ctatttttgt aacagaaact tctttggaag atgctgtatt cattctccgg 480
gggttaaggg aaaaatatga aatttttcat ggtgtgcgca ttacagaagg ggctttgaat 540
gcagctgtag ttctttctta tcgttacatc acagaccgat ttcttctga taaggcgatt 600
gacctaatgg atgaggctgc gagtttaatc cgtatgcaaa taggaagttt acctctgcct 660
attgatgaaa aggaagaga attatcagct ttaatcgtga aacaagaagc tattaacgc 720
gagcaagcac cagcttatca ggaagaggct gaagacatgc aaaaagcaat tgaccgggtt 780
aaggaagagc tggccgcttt acgcttgccg tgggatgaag aaaaaggatt aattgcagga 840
ttaaaagaaa agaagaatgc tttagaanaa ttaaaatttg ccgaagagga agctgagcgt 900
actgccgatt acaatcgggt agcagaacta cgctatagtt tgattccttc tttggaggaa 960
gaaattcatt tagctgagga agcttttaaat caaagagatg ggcgcctgct tcaagaggaa 1020
gttgatgagc ggttgattgc gcaagttgtt gcgaattgga ctggaatccc tgtgcaaaaa 1080
atgttgaggg gagaatctga aaagttattg gtgttgagga gtcttttaga gaaagggttg 1140
tcggacagcc ttctgcctatt gccgcagtca gtgattcgat tcgagctgct cgagtaggat 1200
tgagtgatcc gcagcgtctc cctcacaagg gaatattagc tggcgcggcg aaccgctggc 1260
gaaac 1265

```

<210> 25

<211> 463

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 25
 atgacgaaca accccatgtt tatcgataag gaaagtcgct cttaaagcga gcgatccttc 60
 aggattcaaa acaccaaaag cttctgatat tttaaaagaa gggctctgta acagaggata 120
 ttctgttccc tctattcctc ctgcatttct cgctacagcg agccaacgag aatgtgtctc 180
 aatgtcgtca acggagcaac caagcacgac tgcaccccg ctttcaaaat ctaccaatct 240
 atcttgaaaa gcatgcaatt ctgtaggaca cacatagggtg aaatcttttag gataaaagaa 300
 gagcactaca tacttaaccac gaaagtctgc tagagagatt tctttctctt ctccacaaac 360
 aacggcttta ccagaaaaat ccggagcctg tcttccaatt agtgatocca taatactcct 420
 octagaaaaga aacaacgcac cagagaggat ttgaacctct gac 463

<210> 26

<211> 636

<212> DNA

<213> Chlamydia trachomatis serovar E

<400> 26
 ggtagaaaat tctctgaagt cgaatctgaa attaaaacag tccctacaa agttgctcct 60
 aactcgaaaag gagatgcggt ctttgatgtg gaacaaaaac tgtacactcc agaagaaatc 120
 ggcgtcaga tcctcatgaa gatgaaggaa actgctgagg cttatctcgg agaaacagta 180
 acggaagcag tcattaccgt accagcttac tttaacgatt ctcaaagagc ttctacaaaa 240
 gatgctggac gtatcgagc attagatgtt aaacgcatta ttctgaacc aacagcggcc 300
 gctcttgctt atggtattga taaggaagga gataaaaaa tcgccgtctt cgacttagga 360
 ggaggaactt tcgatatttc tatcttgaa atcggtagcg gagtttttga agttctctca 420
 accaacgggg atactcactt gggaggagac gacttcgacg gagtcatcat caactggatg 480
 ctgtagtaat tcaaaaaaca agaaggcatt gatctaagca aagataacat ggctttgcaa 540
 agattgaaag atgctgctga aaaagcaaaa atagaattgt ctggtgtatc gtctactgaa 600
 atcaatcagc cattcatcac tatcgacgct aatgga 636

<210> 27

<211> 1797

<212> DNA

<213> Chlamydia trachomatis serE

<400> 27
 atgcatcacc atcaccatca catgagcatc aggggagtag gaggcaacgg gaatagtcga 60
 atcccttctc ataattggga tggatcgaat cgcagaagtc aaaatacgaa gggtaataat 120
 aaagttaga atcaggtttg ttctctatat tcatctcgta gtaacgaaaa tagagaatct 180
 ccttatgcag tagtagacgt cagctctatg atcgagagca ccccaacgag tggagagacg 240
 acaagagctt cgcgtggagt gctcagtcgt ttccaaagag gtttagtacg aatagctgac 300
 aaagtaagac gagctgttca gtgtgcgtgg agttcagtc ctacaagcag atcgtctgca 360
 acaagagccg cagaatccgg atcaagtagt cgtactgctc gtggtgcaag ttctgggtat 420
 agggagtatt ctcttcagc agctagaggg ctgctcttta tgttcacaga tttctggaga 480
 actcgggttt tacgccagac ctctcctatg gctggagttt ttgggaatct tgatgtgaac 540
 gaggtcgtt tgatggctgc gtacacaagt gagtgcgagg atcatttaga agcgaaggag 600
 ttggctggcc ctgacggggt agcggccgcc cgggaaattg ctaaaagatg ggagaaaaga 660
 gttagagatc tacaagataa aggtgctgca cgaataattat taaatgatcc tttaggccga 720
 cgaacacctt attatcagag caaaaatcca ggtgagtata ctgtagggaa ttccatgttt 780
 tacgatggtc ctcaggtagc gaatctccag aacgtcgaca ctggtttttg gctggacatg 840
 agcaatctct cagacgttgt attatccaga gagattcaaa caggacttcg agcaccgact 900
 actttggaag aatccatgcc gatgttagag aatttagaag agcgttttag acgtttgcaa 960
 gaaacttgtg atcgggctcg tactgagata gaagaatogg gatggactcg agagtcgcga 1020
 tcaagaatgg aaggcgatga ggcgcaagga ccttctagag tacaacaagc ttttcagagc 1080
 tttgtaaatg aatgtaacag catcgagttc tcatttggga gctttggaga gcatgtgcga 1140
 gttctctgag cttagagtac acgaggatta gctgccgag gagaggcgat tcgccgttgc 1200
 ttctcttggt gtaaaggatc gacgcacgc tacgctcctc gcgatgacct atctctgaa 1260
 ggtgcacgt tagcagagac tttggctaga ttccagatg atatgggaat agagcgagggt 1320
 gctgatggaa cctacgatat tccttttgta gatgattgga gaagaggggt tcctagtatt 1380
 gaaggagaag gatctgactc gatctatgaa atcatgatgc ctatctatga agttatgaat 1440
 atggatctag aaacacgaag atcttttgcg gtacagcaag ggcactatca ggaccaaga 1500
 gcttcagatt atgacctccc acgtgctagc gactatgatt tgccatagaag cccatatcct 1560

13

```

actccacett tgcctcctag atatcagcta cagaatatgg atgtagaagc agggttccgt 1620
gaggcagttt atgcttcttt tgtagcagga atgtacaatt atgtagtac acagccgcaa 1680
gagcgtattc ccaatagtca gcaggtggaa gggattctgc gtgatatgct taccaacggg 1740
tcacagacat ttagagacct gatgaagcgt tggaaatagag aagtcgatag ggaataa 1797

```

<210> 28

<211> 1983

<212> DNA

<213> *Chlamydia trachomatis* serE

<400> 28

```

atgcatcacc atcaccatca catggaatca ggaccagaat cagtttcttc taatcagagc 60
tcgatgaatc caattattaa tgggcaaadc gcttctaatt cggagaccaa agagtcacag 120
aaggcgctccg aagcgagtc ttcagcatcg tcctctgtaa gcagctggag ttttttatcc 180
tcagcaaaga atgcattaat ctctcttcgt gatgccatct tgaataaaaa ttccagtcga 240
acagactctc tctctcaatt agaggcctct acttctacct ctacggttac acgtgtagcg 300
gcaaaagatt atgatgagcg taaatcgaat tttgatacgg cgaaaagtgg attagagaac 360
gctaagacac ttgctgaata cgaaacgaaa atggctgatt tgatggcagc tctccaagat 420
atggagcggt tagctaattc agatcctagt aacaatcata ccgaagaagt aaataatatt 480
aagaaagcgc tcgaagcaca aaaagatact attgataagc tgaataaact cgttacgctg 540
caaaatcaga ataaatcttt aacagaagtg ttgaaaacaa ctgactctgc agatcagatt 600
ccagcgatta atagtcagtt agagatcaac aaaaattctg cagatcaaat tatcaaagat 660
ctggaaagac aaacataaag ttatgaagct gttctcacta acgcaggaga ggttatcaaa 720
gcttcttctg aagcgggaat taagttagga caagctttgc agtctattgt ggtgctggg 780
gaccaaagtc aggtcgagc tctgcaagca cagcaaaata atagcccaga taatattgca 840
gccacgaagg aattaattga tgctgctgaa acgaaggtaa acgagttaaa acaagagcat 900
acagggctaa cggactcgcc tttagtgaat aaagctgagg agcagattag tcaagcacia 960
aaagatatcc aagagatcaa acctagtggg tcggatattc ctatcgttgg tccgagtggg 1020
tcagctgctt ccgcaggaag tgcggcagga gcgttgaaat cctctaacaa ttcaggaaga 1080
atttccttgt tgcttgatga ttagacaat gaaatggcag cgattgcact gcaagggttt 1140
cgatctatga tcgaacaatt taatgtaaac aatcctgcaa cagctaaaga gctacaagct 1200
atggaggctc agctgactgc gatgtcagat caactggttg gtgcggatgg cgagctccca 1260
gccgaaatcc aagcaatcaa agatgctctt gcgcaagctt tgaaacaacc atcagcagat 1320
ggtttggtta cagctatggg acaagtggct tttgcagctg ccaaggttgg aggaggctcc 1380
gcaggaacag ctggcactgt ccagatgaat gtaaaacagc tttacaagac agcgttttct 1440
tcgacttctt ccagctctta tgcagcagca ctttccgatg gatattctgc ttacaaaaca 1500
ctgaactctt tatattccga aagcagaagc ggcgtgcagt cagctattag tcaaaactgca 1560
aatcccgccg tttccagaag cgtttctcgt tctggcatag aaagtcaagg acgcagtgca 1620
gatgctagcc aaagagcagc agaaactatt gtcagagata gccaaacgtt aggtgatgta 1680
tatagccgct tacaggttct ggattctttg atgtctacga ttgtgagcaa tccgcaagca 1740
aatcaagaag agattatgca gaagctcacg gcattctatta gcaaagctcc acaatttggg 1800
tatcctgctg ttcagaattc tgcggatagc ttgcagaagt ttgctgogca attggaaaga 1860
gagtttgttg atggggaacg tagtctcgca gaatctcaag agaatgcgtt tagaaaacag 1920
cccgttttca ttcaacaggt gttggttaaac attgcttctc tattctctgg ttatctttct 1980
taa 1983

```

<210> 29

<211> 1224

<212> DNA

<213> *Chlamydia trachomatis* serE

<400> 29

```

gtaacttttc aacatttttc acaatgacaa gaataaaaagc aaaaagaaag gctgccgata 60
aaataaaaag tttactgcga gaacagaaga ctaaaactat ctggacgaat aagccggatg 120
cgcaggataa ttgcgcataa aacaotttta tagagagtga tcttatgtct aaaacaccat 180
tatccatagc tcatccttgg catgggcccag tattaacacg cgatgattat gaatctcttt 240
gttgctatat agaaatcact ccagccgact ccgttaaatt cgaactggat aaagaaactg 300
gtatcctaaa agtggatcgg ccacaaaagt tttctaactt ttgtccttgc ttatacgggc 360
tgttacctaa gacttattgt ggagatcttt ctggagaata cagtgggtcaa caaagtaaca 420
gagagaatat caaaggcgat ggcatcctc ttgatattct tgtgttaacg gaaaaaata 480
ttacacaagg gaacatcctc ttgcaagcgc gtcctatcgg agggattcgt atttttagact 540

```

```

cggaagaagc cgatgataaa atcatcgctg ttctagaaga tgatttagtc tatggcaata 600
tagaagatat ttctgaatgc ccaggcacag ttttggacat gatccaacac tatttcttaa 660
cctataaagc tactccagaa agcttaattc aagcaaaacc agctaaaatt gaaattgtag 720
gtttatacgg caaaaaagaa gctcaaaaag tcattcgtct tgctcacgaa gactattgca 780
atctttttat gtaaatacgac agaaaaagaa aaggctgttg tgggagattc cacaacggcc 840
cctcctaacc aagttttttt catcctaggg gactttatga agcaaataga taactttgaa 900
caaattcatc tctcgtgccc aattcggcac gagattaaaa caaagctctc aaaaagagtt 960
ggtatcccg aattcattcag cagttcccgg tgccaaagtt aaagagatac gctttttatt 1020
aggatagtta tggacgcaca agaaaaagaa tacgacgcat cagccatcac cgtttttagaa 1080
ggattgcaag ctggtcgtga gcgtcctgga atgtacattg gtgatacagg agttaccgga 1140
ttgcatcact tggtttatga agtggtggat aacagtatcg atgaggcaat ggcggggttt 1200
tgtaccgagg tcgttggtcg cata 1224

```

<210> 30

<211> 883

<212> DNA

<213> Chlamydia trachomatis serE

<400> 30

```

atgttgacta acatggcgac catcagaaac tctgtgaaga cattgaacag aattgaattg 60
gatcttgaag cttctaattc tggctttacg aaaaaagaga tcgctttatt aacgaaaaga 120
catcgcaagt tgcttaacaa cctggaaggt gttcgtcata tgaactctct cccagggtct 180
ttaattgtaa ttgaccggg ctatgagcgc attgctgtcg cagaagctgg aaaactaggc 240
attcctgtaa tggccttagt tgatacaaac tgcgatccaa caccaatcaa ccacgttatt 300
ccttgcaacg atgattccat taagagtatc cgtctggttg tcaatgtact taaagacgct 360
gttattgatg cgaagaagcg ttcaggcatc gaaattttat ctccagtacg tctgttagaa 420
agacctgcag aagaagctgt ggaagagttg cctcttccaa caggtgaagc tcaagatgaa 480
gcttcttcta aagaaggttt tttacttttg gcagatattg acaattgcgg ggcattgaaa 540
tgagcgactt ctccatggaa acattgaaaa atttaagaca gcagacaggt gtaggcctga 600
ctaaatgtaa agaggctcta gagcatgcta agggcaattt agaagatgct gttgtttatt 660
tacgtaagct tggctctgcc tctgcaggca aaaaagaaca ccgagaaaaca aaagaaggcg 720
taattgctgc actcgttgat gaacgtgggt cggcacttgt tgaagtcaac gttgaaactg 780
atthttgtgc taacaacagt gttttccgag cattcgttac aggtttgtta tccgatcttc 840
ttgaccacaa gcttagcgat gttgaagctt tagctcgcgt aat 883

```

<210> 31

<211> 393

<212> DNA

<213> Chlamydia trachomatis serE

<400> 31

```

agttgaaaaa ggctgtttct tgcattcaaa aaactatcga gcaagagaga tctattttgt 60
ttgttggaac aaaaaaacag gcaaaacaga tcattagaga agctgctatc gaattgtggc 120
aattctttgc ttcagagaga tggttgggtg gcatgttgac taacatggcg accatcagaa 180
actctgtgaa gacattgaac agaattgaat tggatcttga agcttctaatt tctgggtctta 240
cgaaaaaaga gatcgcttta ttaacgaaaa gacatcgcaa gttgcttaac aacctggaag 300
gtgttcgtca tatgaactct ctcccagggc ttttaattgt aattgacccg ggctatgagc 360
gcattgctgt cgcagaagct ggaaaactag gca 393

```

<210> 32

<211> 2577

<212> DNA

<213> Chlamydia trachomatis serE

<400> 32

```

attacggagg ccatacggta tcttctcgag gaggatttca agggatatgc gtacgaatag 60
ccgattttatt ccgtaactgt ttctctcgta atagaggcac tactactacg ccatctcgaa 120
ctgttatcac tcaggcagat atttatcatc cgactatttc tggacaagga gctcaaccta 180
ttgtctctac aggagataag aaattagata gcgcaattat tcaagcagat ttgcgtgcgc 240
agaataaaca gactttggct acacatattc aaagtaagct aggttctatg gagggacaat 300
ctcctcaaga ttataaagct ggtgcgtata gtgcgctaag attgatgctg tttactccag 360

```



```

gcgaaactac tgtgagtagc gagcgggaac gtcaagcgtg cgttacgggt cgggactctct 420
gggaacaggc tgcaggagat cttgctacca atgggaatac agatgggctt atgttaatgg 480
ctaactatc tgtgggaggg aagcatgtgc ctgcggggca ttttaagaga tacatggata 540
ctgtaaaggg tacgtttact gatgagaacg aggctacaga tcctacggta gatgccattt 600
tagatttagc agcaaaaatc gatgcgacgg aattctctag tcctggttca gggcaagtca 660
ttcttaatta tataggaaat tatggacaag tcgtttttag aaacgaggag atgaaccttc 720
ttgtttttag agatcaaaaat gggcaagatc ctcaacgtgt tcaagataac tcaaaagagt 780
tacaaaaact gttagaaaat gctcgaaaaa cagatcctga gttatatttc caaacactaa 840
ctgtcataac ttcttctgtt ttcttagact aaagagaagg tatacggtgt tcggtccttt 900
caactattaa gaggaagtag tggtagtag cataagccct atagggggga attctggggc 960
agagggattt tctagtgcac ctcgaggcga tgagattgat gatgtaccag atagtgaaga 1020
gggagagcta gaagagcggg tttcggatca tgcagagtct atcattaccg agagctcgga 1080
aacgctgttt cgtactactt cttcatcagg ggtcagtgaa gatcttcagc aacacggttag 1140
cttggaggaa tctccacgac aacgaggttt ccttggacgg atccgtgatg cagtagcttc 1200
tatttgggaag cgtcgtgttg cacgaaggaa tgaaaactat gatgtgaaaa aagcagaaga 1260
gcagcaaggg attgtgcaat atctgcagga ttcgaaaatg cctgctttta cgcgtgccta 1320
tcgccatctc cgtgctttca attctgcacg cttacgtacg attcgtgagt ttttcgctac 1380
catttttctg gctttaaggg atgcgtatta tcgacattgt acacgttctg ggatcaactt 1440
ttgtggagct gataaagact ctttagaagt tcttgttgcg gtgggtttgc ttttgcgtat 1500
ggctacctta cgctcttttg aacatgtcgg ttgggaattac gaagatcgat tagtaaataa 1560
tgatgctccg gtgacagggt cggggagaac tcttgttgat gatgctgtag acgatattga 1620
atcgatttta aatacgagaa ccaactggcc tcaacatgtc atgatagggt tttctcgtgg 1680
tctcgttcaa ttatgtgcga ctctttataa tgcgacttct caagaatgtt tcaagtcgat 1740
tgttcgttta gaaaaagaag acccttcttc agattattct caagctttat tattagcagg 1800
gataatagat cgcttggcgg agaagcccc tatggctgca aagtatgttt tggatgcat 1860
gcgtgttcga acttcggagc tcataggaga actcattatt ctcgatttgc ttctcctgt 1920
atggaaggtt ggccgcggag gcgtattccc tctgtgaat gagcagctcg ttgtgcaaat 1980
tgtaaatgca aacgtagaac gattgcattc cactttcgtc catgagccac aagcttattt 2040
gcgtatgatc gaaggttttg taaccaattt ctttttctta cctagcgagg aagatccttc 2100
ttcggttggg aatatctaac aacattttct aatagggaag aggataaata gcgtgaaata 2160
atactgatta tgtgaagaat aggcaaaaag acctaaatcc ttatatgcta ttagattctc 2220
gtttccctac agattattat ttacgtatcc tagaattagt catccgggat gcttcttgta 2280
aattggtata taaccgacgc ctgcatatgt tggaggcgat ccctcttgat caaaaacttt 2340
ctactgatca agagggggaa tcaagtattt tacgagaagt gattagcgag ctacttgcgc 2400
attctgggga aagttatgcg atttcagctc aattacttgc cgtaatogat atttatttaa 2460
aacaagagca accgtcgaat tcatggttcg ctcgaaatct tcggaagaga gagcgggcta 2520
gaaaacgaca aacaattaat aagttgcttt tgttaaaaag tatcctattt tttgaac 2577

```

<210> 33
 <211> 554
 <212> DNA
 <213> Chlamydia trachomatis serE

```

<400> 33
ttctttatta aaaaaaactt tctcttttct ctcagacttc ttatgagtca agaaactcaa 60
cgagtcttgg tgtatggaga aggatttttt agaaaatgtt tatcgtcatt tccgttaccg 120
ttttttttaa ttaagtgtac ttccagctct tctcggactc tggctatttt ttactcctaa 180
tattcttaac tatttggatt cttctgttat tttatcagat aaaatttgog gcgtcctttt 240
aattttatta tcagctttat ctttttataa tctgtttatt ttgcaactag gcatttttat 300
tggtctctgg gtttctttct tttctgttgc ttccgaccta cttccttttag tatttgccta 360
tgattcgcta ctaggttttg ccacactagc tattattttt ctactcccta atcgtcctga 420
agatctagaa gttggtccta ctattccaga aacttgccat tataatcctt cttccggagg 480
gaaaagagct gcggttctta tttttgcttt tgtaggatgg ttacaaagtc gctacttaac 540
ttccgcggca cgag

```

<210> 34
 <211> 1433
 <212> DNA
 <213> Chlamydia trachomatis serE

<400> 34

```

ctgcacgaaa attattaaat gatccttttg gccgacgaac acctaattat cagagcaaaa 60
atccagggtga gtataactgta gggaaattcca tgtttttacga tgggtcctcag gtagcgaatc 120
tccagaacgt cgacactggt ttttggtgtg acatgagcaa tctctcagac gttgtattat 180
ccagagagat tcaaacagga cttcgagcac gagctacttt ggaagaatcc atgccgatgt 240
tagagaattt agaagagcgt tttagacgtt tgcaagaaac ttgtgatgog gctcgtactg 300
agatagaaga atcgggatgg actcgagagt ccgcatacaag aatggaaggc gatgaggcgc 360
aaggaccttc tagagcacia caagcttttc agagctttgt aaatgaatgt aacagcatcg 420
agttctcatt tgggagcttt ggagagcatg tgcgagttct ctgcgctaga gtatcacgag 480
gattagctgc cgcaggagag gcgattcgcc gttgcttctc ttgtttgtaa ggatcgacgc 540
atcgctacgc tctcgcgat gacctatctc ctgaagggtgc atcgttagca gagactttgg 600
ctagattcgc agatgatag ggaatagagc gaggtgctga tggaaacctac gatattcctt 660
tggtagatga ttggagaaga ggggttccta gtattgaagg agaaggatct gactcgatct 720
atgaaatcat gatgcctatc tatgaagtta tgaatatgga tctagaaaca cgaagatctt 780
ttgcggtaca gcaagggcac tatcaggacc caagagcttc agattatgac ctcccacgtg 840
ctagcgacta tgatttgctt agaagcccat atcctactcc acctttgctt cctagatatc 900
agctacagaa tatggatgta gaagcagggt tccgtgaggc agtttatgct tctttttagt 960
caggaatgta caattatgta gtgacacagc cgcaagagcg tattcccaat agtcagcagg 1020
tggaagggat tctgcgtgat atgcttacca acgggtcaca gacatttaga gacctgatga 1080
agcgtttgaa tagagaagtc gatagggaat aaactggtat ctaccatagg tttgtagcaa 1140
aaaactaagc ccaccaagaa gaaattctct ttgggtgggt tcttttttta ttcaaaaaag 1200
aaagccctct tcaagattat accaagatgg gatgtataat ctgaaaggaa ggcgttttat 1260
tctctatcca tatgatggtg gtggtatcct ccttttagagg agcagcagtc tccatgacgt 1320
tttttgaagc agcacttcaa gaagtttagg cagaccataa cccagcgat tcccgttact 1380
acataagctg cttgtgtcca catggttcct tcaccaagca ggtgagtaag tag 1433

```

<210> 35

<211> 196

<212> DNA

<213> Chlamydia trachomatis

<400> 35

```

ctcgtgcoga tgatacagca gtgcagtgta atcaagggtg taaacgcaag ggagctgtat 60
gcgtctattt agaagtttgg cacctcgact acgaagattt ccttgaattg agaaagaata 120
caggggatga gcgtcgacgg gctcatgatg tcaatatagc tagctggatt ccagatcttt 180
tcttcaaacy tttaca 196

```

<210> 36

<211> 1990

<212> DNA

<213> Chlamydia trachomatis

<400> 36

```

ttcaactaggc tcatgagcct ctaactcttc tggagtaact cctagagcaa acacaaactg 60
cttccacaaa tcaatatgat tagggtaacc gttctcttca tccatcaagt tatctaacia 120
taacttacgc gcctctaaat catcgcaacg actatgaatc gcagataaat atttaggaaa 180
ggctttgata tgtaaaataat agtcttttggc atacgcctgt aattgctctt tagtaagctc 240
ccccttcgac catttcacat aaaacgtgtg ttctagcata tgcttatttt gaataattaa 300
atctaactga tctaataaat tcataaacac ctccatcatt tcttttcttg actocacgta 360
accgcttgca aaaaagggtcc gtataagtcc tctgtttcat ctatgcgcaa agaacaatac 420
tcttctcgag aagtaggatg tgaatggtag accatattag gtgcctgctc tatcaccgct 480
aacgggtgtt gctcattccc ctctcccata caaacaacag ccgcaactgc taaggcatct 540
acaagattac tttgcgtcat ctgtaagaga cgaccgaaac aatctagcga tccatatatg 600
ttgtgtaatg gagaaaaatcc ataccaaac acgccgatac ccagtaactcc acgccgcatt 660
ggagtagtat ggctatctgt aatgattacg cctagctctt tcaactogaaa ataatttctt 720
aaccattctc cgatgcgatt acacgatccc aaaatatctt taggatataa aacaaaaggc 780
tgggtccgat tgcattcatc aatccctgca gaaggatca aaataccttc ttttttcgtt 840
agatatatcc cgcttttctc acaaaacaaa taagcatccg cttctttttt tatcagctct 900
gctttgcaca ttcttgcatc agcgacagcg ccttcacata aactcacaat ctttgaagag 960
acaactacca cactccgttc ttgcagagcg ggcaaaagcct cttgcaagat ctcttgaagc 1020
gaatcatgtg caaatacttt acgtgttttg atcggagtta ttttcataat aataaatact 1080

```

```

gaaatcctct gtattacaaa tacattcctt cttccatcct gataatcgcg tgatagggaa 1140
gaaagtatcg ccccaatatt cctttttgat atgtgtgaca aaacaagctt tcagaagggt 1200
ttgttgga aaactttcaa agagctccgc tcccccaatt aaaaacggat gattcaaaga 1260
tagtgtccca tactctgcaa aggaagaaac tcctatgcat tgttggtggat gcatcctgcg 1320
agaaaagaca acgatatccc gcccatgctt atacttgtct ggaagagact cccaagtctt 1380
tcgtcccata atgatgggat gatttcgaat ggtttctgca aaaaaacgta gatcttcggg 1440
ataactccaa gggagcttgc ctaaagctcc catcactcct ctgggatcaa tagcaacgat 1500
acctgttgct tggatcatat aaacatacca gcccaagcag cagcggctaa ggcacgtctg 1560
ttaccttcaa cctgatgcac gcgtagataa tcaactcctc gatcatgaag agatacagaa 1620
cagccgatog tttcccaatc acgatcgtta ctattaaatc ggcccaacat actcaaacac 1680
gattttctag aatggcctat taatacagga cactctaaaa cactgtttaa ctgctttact 1740
ccatccatca ataacatcga ctgaacggga gtcttcccaa atcctattcc tggatogaaa 1800
acaacttgcc aacttgtatc taaacctact tgagcaaat gttctaactg ggactctccc 1860
caacgcaaca tttgtcfaat aggagattct tcataagaaa gtacacaatc tggctttgga 1920
ggcagcgaac acgaatgatt tattaatagc cgtagcccaa actccttcgc caaatgagcc 1980
atttccaaag

```

<210> 37

<211> 2093

<212> DNA

<213> Chlamydia trachomatis

<400> 37

```

cagaaactct atccgcatac cttcttcggc aaattgatac caattttgoc tcttctcagg 60
aacgtactat agctcagtat attgtaggca acctctcccc agaaggactc tttttagaaa 120
atcctagtct tgtggctgca gatttaaacc tttccgaaca ccttttccac aaggtatggc 180
aacgtatcca acaattacat cctttaggag tggagcgcc ttccctacag tctactggg 240
tatcgctact acagacatct ccccataagg aggccttagc tattattcgc aaccatttcc 300
ctagattagc tcgtttgtgat ttactacta tcgctaggaa aatgcatgca accacaacag 360
agattcttac atttcttaga cagcgttttg cttccatccc ttggtgtcca gcagcaggct 420
tttccgagac actgcacccc cctgctccag cgcttctcga tgcctacott tcttctcgc 480
gaaactctta ttgggatgtc tctattaata aagattgtct cccctctatt agactcaacg 540
acaccgtact agatatctat ccttctcttc ctgctgaaga gaaagaccac ctatcgcaac 600
aaatccgagc agcaaaaaca ttgcttcgca atgtaaaaaa acgagaagaa acgttattgg 660
ctatccttgc agttctcatc cctaccaag aagagttcct tcttaaaaaa cgcacctctc 720
ctaaagcttt ttctgtaaaa caaatagctc gogaactctc tcttcatgaa gctaccgttt 780
gtcgcgccat tgataataaa acgttagcaa cccctgttgg attactccct atgcgatcgc 840
tatttccaca agcgtttgga tcttgccccg atcaatctaa agcaactatt ttgcattgga 900
tccaccagtg gatttctaca gaaaaacatc ctctatctga tgcagctatt agccaaaaaa 960
ttatttgaaa gggcatcccc tgcgcacgac gcacagtagc caaatatcgt tcgcaactga 1020
atatcccacc tgcgcaccaa cgcaaacacc tatgctctgt ttttaacaaca acacgcacag 1080
agaattctcg acatactatc taatcggata tgtaaagctg ctttacatcc cttgaactag 1140
aaataaaaatg gaaataaaaa gcccagaaca agagaagttg ttctgggctg acagaagctg 1200
tcagatcatt ttaataagat tgatgacaac tacgacaagt tcctggatcc aaaaaagaat 1260
ctaaaaagcc atacaaagat tgcgttactt cttgcgatgc ctctaactc ttatcagcgt 1320
catctttgag aagcatctca atgagcgctt tttcttctct agcatgcgc acatccgctt 1380
cttcattgtc tgtgaaatat gcatagtctt caggattgga aaatccaaag tactcagtca 1440
atccacgaat tttctctcta gcgatacgtg gaatttgact ctcataagaa tacaaagcag 1500
ccactcctgc agctaaagaa tctcctgtac accaccgcac gaaagtagct actttcgtt 1560
ttgctgcttc actaggctca tgagcctcta actcttctgg agtaactcct agagcaaaaca 1620
caaactgctt ccacaaatca atatgattag ggtaaccgtt ctcttcatcc atcaagttat 1680
ctaacaataa cttacgcgcc tctaaatcat cgcaacgact atgaatcgca gataaatatt 1740
taggaaaggc tttgatatgt aaataatagt ctttgccata cgctgtaat tgctctttag 1800
taagctcccc cttcgaccat ttcacataaa acgtgtgttc tagcatatgc ttattttgaa 1860
taattaaato taactgatct aaaaaattca taaacacctc catcatttct tttcttgact 1920
ccacgtaacc gcttgcaaaa aaggctccgt taagtctctt gtttcatcta tgcgcaaaga 1980
acaatactct tctcgagaag taggatgtga atggtagacc atattaggtg cctgctctat 2040
caccgctaac ggtgtttgct cattccccctc tccatacaaa acaacagccg caa 2093

```

<210> 38

<211> 1834

<212> DNA

<213> Chlamydia trachomatis

<400> 38

```

ctctacttct acctctacgg ttacacgtgt agcggcaaaa gattatgatg aggctaaatc 60
gaattttgat acggcgaaaa gtggattaga gaacgctaag acacttgctg aatacgaaac 120
gaaaatggct gatttgatgg cagctctcca agatatggag cgttttagcta attcagatcc 180
tagtaacaat cataccgaag aagtaaataa tattaagaaa gcgctcgaag cacaaaaaga 240
tactattgat aagctgaata aactcgttac gctgcaaaat cagaataaat ctttaacaga 300
agtgttgaaa acaactgact ctgcagatca gattccagcg attaatagtc agttagagat 360
caacaaaaat tctgcagatc aaattatcaa agatctggaa agacaaaaca taagtatatga 420
agctgttctc actaacgcag gagaggttat acaaaagatc tctgaagcgg gaattaagtt 480
aggacaagct ttgcagtcta ttgtggatgc tggggaccaa agtcaggctg cagttctgca 540
agcacagcaa aataatagcc cagataatat tgcagccacg aaggaattaa ttgatgctgc 600
tgaaacgaag gtaaacgagt taaaacaaga gcatacaggg ctaacggact cgcctttagt 660
gaaaaaaagt gaggagcaga ttagtcaagc acaaaaagat attcaagaga tcaaacctag 720
tggttcggat attcctatcg ttggtccgag tgggtcagct gcttccgcag gaagtgcggc 780
aggagcgttg aaatcctcta acaattcagg aagaatttcc ttgttgcttg atgatgtaga 840
caatgaaatg gcagcgattg cactgcaagg ttttcgatct atgatcgaac aatttaatgt 900
aaacaatcct gcaacagcta aagagctaca agctatggag gctcagctga ctgcgatgtc 960
agatcaactg gttggtgcgg atggcgagct cccagccgaa atacaagcaa tcaaagatgc 1020
tcttgcgcaa gctttgaaac aaccatcagc agatggtttg gctacagcta tgggacaagt 1080
ggctttttgca gctgccaggg ttggaggagg ctccgcagga acagctggca ctgtccagat 1140
gaatgtaaaa cagctttaca agacagcggt ttotctgact tcttccagct cttatgcagc 1200
agcacttttc gatggatatt ctgcttaca aacactgaac tctttatatt ccgaaagcag 1260
aagcggcgtg cagtcagcta ttagtcaaac tgcaaatccc gogctttcca gaagcgtttc 1320
tcgttctggc atagaaagtc aaggacgcag tgcagatgct agccaaagag cagcagaaac 1380
tattgtcaga gatagccaaa cgttagggtg tgtatatagc ogcttacagg ttctggattc 1440
tttgatgtct acgattgtga gcaatccgca agcaaatcaa gaagagatta tgcagaagct 1500
cacggcatct attagcaaaag ctccacaatt tgggtatcct gctgttcaga attctgcgga 1560
tagcttgcaag aagtttgctg cgcaattgga aagagagttt gttgatgggg aacgtagtct 1620
cgcagaatct caagagaatg cgtttagaaa acagcccgct ttcattcaac aggtgttggt 1680
aaacattgct tctctattct ctggttatct ttcttaacgt gtgattgaag tttgtgaatg 1740
agggggagcc aaaaaagaat ttcttttttg gctctttttt cttttcaaaag gaatctcgtg 1800
tctacagaag tcttttcagc acgagcggca cgag 1834

```

<210> 39

<211> 1180

<212> DNA

<213> Chlamydia trachomatis

<400> 39

```

agaaatttct caaaaatcaa agttttttac atttaagggg catcttacca ccacaacaac 60
cttctatgag cagaaaactat ccattaaata aaagtaatta aatataacaa aaacatcttg 120
attatttttg ttaaaaagaaa tacttaatga gttttattta attaacgaaa cgaaaagctt 180
gctaataaaa attattcaca cagctatcga atttgctccg gtaatcaaaag ccggaggcct 240
gggagacgcg ctatacggac tagcaaaaagc ttttagccgct aatcacacaa cgggaagtgtg 300
aatcccttta taccctaaat tatttacttt gcccaaagaa caagatcttt gctcgatcca 360
aaaattatct tatttttttg ctggagagca agaagcaact gctttctcct acttttatga 420
aggaattaaa gtaactctat tcaaaactcg cacacagcca gagttattcg agaattgcga 480
aacaatctac acaagcgatg atgccttccg tttttgogct ttttctgctg ctgcggcctc 540
ctacatccaa aaagaggag ccaatatcgt tcattttacac gattggcata caggattagt 600
tgctggacta ctcaacaac agccctgctc tcaattacaa aagattgttc ttaccctaca 660
taatttttgt tattcaggct atacaacag tgaatatatta gaagctcctt ctttgaatga 720
attttatatc agccagtacc aactatttct cgatccacaa acttgttgtg tgctaaaagg 780
agcttttatac tgttcagatt tcgtgactac ggtttctcct acatacgcca aagaaattct 840
tgaagattat tccgattacg aaattcacga tgccattact gctagacaac atcatctccg 900
cgggatttta aatggaatcg acacgacaat ttgggggcct gaaacggatc ccaatttagc 960
gaaaaactac actaaagagc ttttcgagac cccttcaatt ttttttgaag ctaaaagcga 1020
gaataaaaaa gccttgtagc aaagattagg cctctcttta gaacactctc cttgcgtgtg 1080
cattatttct agaattgctg agcagaaagg tcctcacttt atgaaacagg ccattctcca 1140

```

tgcaactagaa aacgcttaca cgctcattat tataggtacc

1180

<210> 40

<211> 1297

<212> DNA

<213> Chlamydia trachomatis

<400> 40

```

agaaacttct ataggagggg atgtgatcga cataggtacg tgtgagttat gggatatcga 60
tttgttgtat aatggataag aaattctctg aagataaaga ggctcctcca actaaaagac 120
cattaacatc agggcagagg gcaagtgagc gagcattatc ggctttcaca gatcctccgt 180
aaagaatggg ggtgcggtcc gcaatatctt tggaaaagag agaagcaatc gtttttctac 240
agaaagcatg ggtttcctga actagatcag gatgagctac ttttccggtg cctatagccc 300
agactgggtc ataagctaga atgaaagagg cttgctcagg gagtttagat aatcctatag 360
tcagttgatt taaaagaata tcttgagttg ctccagattc ttgttcttct aaagtttctc 420
caatacacag aactggaatc attccactat ggatagctgc agcagctttt tcagcaagta 480
caggattttg ttcatgaaag atatgacgtc tttcggaatg tccgatgaga acaaaatcga 540
ctccgatatc tttgagcatt ggggctgaaa totcaccagt aaaagctcct gagtcagctt 600
catgagtggt ttgggctcca agaaagatgg gggaatcgct tacagcttgt tgacaagctg 660
acagcagtg gaaaggagga atgattcctg taatgatttg gggattagac agaatgtcac 720
tagagatgaa actttttaaa aaggtctgag cttcggtaaag cgtcttggtc attttccaat 780
taccgaaaac aaattgcttt gatggctcag agtggagaag gtgggcccac gttggaaatg 840
gttttctgtg agtttctttg tctgtaaaac tgagatttgc tgaataacct gtgcatgtat 900
tttgtttgta agatagatca aagcgtaata ctcgatttct tgcaaggag gcttattttt 960
atatgattta ttttctattg cttgatata aatctcttgg atatgcta atctcctgtct 1020
tacttttttc tgtgaatttg cttaaatagt tggtttttag ccctttgtta tatgaagggt 1080
aaaatttggt gtattacgca tcctgatgat gctcgggaag ctgccaaagc gggagccgat 1140
tacattggca tgatttttgc taaagattct cgaagatgtg tgagtgaaga aaaagcaaag 1200
tatatcgtag aggcataaca ggaagggaat tcggaacctg ttggagtatt ccagagcat 1260
tcagtagaag aaatttttag tattactgag acgacag 1297

```

<210> 41

<211> 1141

<212> DNA

<213> Chlamydia trachomatis

<400> 41

```

ctttccataa gttcttttct tcaatgattc tagcttatto ttgctgctct ttaagtggg 60
gggggtatgc agcagaaatc atgattcctc aaggaattta cgatggggag acgttaactg 120
tatcattttc ctatactggt ataggagatc cgagtgggac tactgttttt tctgcaggag 180
agttaacggt aaaaaatctt gacaattcta ttgcagcttt gcctttaagt tgttttggga 240
acttattagg gagttttact gttttaggga gaggacactc gttgactttc gagaacatac 300
ggacttctac aaatggagct gcactaagtg acagcgctaa tagcgggtta ttactattg 360
aggggtttta agaattatct tttccaatt gcaactcatt acttgcgta ctgctgtctg 420
caacgactaa taatggtagc cagactccga cgacaacatc tacacgctc aatggtaacta 480
tttattctaa aacagatctt ttgttactca ataagagaa gttctcattc tatagtaatt 540
tagtctctgg agatggggga gctatagatg ctaagagctt aacggttcaa ggaattagca 600
agctttgtgt cttccaagaa aatactgctc aagctgatgg gggagcttgt caagtagtca 660
ccagtttctc tgctatggct aacgaggctc ctattgcctt tatagcgaat gttgcaggag 720
taagaggggg agggattgct gctgttcagg atgggcagca gggagtgtca tcatctactt 780
caacagaaga tccagtagta agtttttcca gaaatactgc ggtagagttt gatgggaacg 840
tagcccgagt aggagagggg atttactcct acgggaacgt tgctttcctg aataatggaa 900
aaaccttggt tctcaacaat gttgcttctc ctggtttacat tgctgctgag caaccaacaa 960
atggacaggg ttctaatacg agtgataatt acggagatgg aggagctatc ttctgtaaga 1020
atgggtgcgc agcagcagga tccaataact ctggatcagt ttcttttgat ggagagggag 1080
tagttttctt tagtagcaat gtagctgctg ggaaaggggg agctatttat gccaaaaagc 1140
t 1141

```

<210> 42

<211> 822

<212> DNA

<213> Chlamydia trachomatis

<400> 42

```

cggcaccaggt gtatgctgaa caagcagaag ggccactga gaacgagcct ctgagaaaaa 60
aagcttttat taaaaatta aaaaaatact ttacaaaact tattctgtag gttgagaaag 120
agcttcaacg taagcattcc aaagctccgt acttacaata ttattgogga tagagcgaat 180
taattctctt ttttagtgatg gaagagggtt tttggggctg aagcgagcca aaagatcttt 240
atcgccaact tgacgagcta actctaacac ccgttcgata tcggtttttg tgaattcac 300
aaagtctctg cgcttttttag aacctcgagg agctcgtggt ttagggctaa tggatctggg 360
agtgatagaa tcgatcacia acgtctttta catttttaac agttgctcag gagcagagtt 420
cttcattttt ttttaagtaa aatgatgcat gtagccgctt gttggccctg ggagataacg 480
acaaagatca ttttctttac ttctccgac tttgctaato gctttagtta tgagctgctc 540
tatttcttct tggatagtaa tctgtgccgt agccatgaat agctccttag tgggtagtct 600
agttctacag atggtagttt ttgctttatt aattgtaata gtcaactaag tctgtttttt 660
tcgatttaat gttcagtcga aataaaaaatc aattagtgtt tatcttttgg tgaattctat 720
agtggttttt gcttttttct caatctcatt ttagagattt ttttgatttg gacaaaagaa 780
aataaagtac ttcagattgt tttctaagtt tgtttgcata aa 822

```

<210> 43

<211> 1634

<212> DNA

<213> Chlamydia trachomatis

<400> 43

```

ataaaaaatt aaattttggc tactccctgc tcctaataga atttcaccag aggagcttgc 60
tactgttatt gcatttcttc taggaggatt agctgacgta ctggtaccat ttgcattagt 120
tacattagtc acaatatattt cattaaaaat aatagcatgg cgtcggcgag aaatttttga 180
gttactgggt ccgtctatat agatagcgcc ccccttatta ttggcgatat tgtttataaa 240
gtaggtaggg ccattatcca ctagggtaac tacaggagcg taaatagctc cgcataaatt 300
ttttgtgata ttgtcactaa aaaagatcct accacgattg cctgtaacat ctaggcgagt 360
agttacttta attgctcctc catcagaagc ttctgaagaa gctgtttcta catttttaaa 420
gcagcgattg ttatagaaaa cgatgtttacc acgatttctt gttagagaac agatagggga 480
gaagatcgct cctcctgcac aacaggcggt attgatgaag aagagatcgc agttattact 540
ctcaaaaagaa ttgctcgttc cagcatagat agcgccacct tttcctgctg tattagtgtg 600
aatacagatg ttgtccataa agagaaaaaca agactgattc tcgctcacia caaaggattt 660
agcggtaact atggctcctc cttggacata agaaaaagttc ttcataaattc cgaccacatc 720
atgalltatga tttatgtaaa gatttttgagc atgaatggct ccgccttctc ttattttatc 780
agcagcataa ggattttctc atgtaaatag tctgcaacaa gtattatttt caaagattac 840
aggacctatt gtatcacgaa tctccacggt aggagaattg ggactcgcat aaccaatcgc 900
accaccaact tcaggggtga gattttttgc aaaataaata ccttcttttt gtgtatcaaa 960
aaagcttagg taatctgtta ttgtgacagc agctccttca ttgggagttt tttgtagaat 1020
agccagtatg tagcgtaggt tatcgagata gcagttagtg agattgtgag tgtctcctgt 1080
caaaactaatt ttatttgata gcgactcttt cgtaggatct ggaactgagt tgggcataag 1140
aaagattcta gaaggaacct ctctagctag tcctgatagg gagtttccga taaggaaaaa 1200
gaaaaacgct tttttcataa ttaaaagacc agagctcctc ctgcattgat gtagtgtgag 1260
acagtggag, tagccacttc tgcttgatag ttagcaaata gtttcagatg agaaaatttg 1320
agggagtggag aacctctccc ataaaaggaa tgtttagcta atgggggtatt tgtggtgacc 1380
caagaaccgt tattttggat taatagtgtg ttgagtagag gacgtttcca gtagagggtg 1440
ggttggtgaag ctagtcccat ttccaagag agtgttggcc atgtatcaga agaataagct 1500
cctttgattc ctattggaga gacaacggca gtatgggctt gctctaattg aaataatcta 1560
gctagatcac cgctttctcg gatagaagct ggttctgttc gagagaataa agcctgagca 1620
aatggggtga gcat 1634

```

<210> 44

<211> 1862

<212> DNA

<213> Chlamydia trachomatis

<400> 44

```

gttagctttc cctccaggga tttgcaattt aatgatttta atgatttttt tattcgacat 60
attcaaccct ttcatttggc aagacatgga gtcatagtta gaggggtctat gtatgcttct 120

```

```

ctaacaagca atatagaagt atatggccat ggaagatatg agtatcgaga tactttctcga 180
ggttatgggt tgagtgcagg aagtaaaagtc cggtttctaaa aatatgggtt agatagttaa 240
gtgttagcga tgcctttttc tttgagatct acatcatttt gtttttttagc ttgtttgtgt 300
tcctattcgt atggattcgc gagctctcct caagtgttaa cacctaagt aaccactcct 360
tttaaggggg acgatgttta cttgaatgga gactgcgctt ttgtcaatgt ctatgcaggg 420
gcagagaacg gctcaattat ctcagctaatt ggcgacaatt taacgattac cggacaaaac 480
catacattat catttacaga ttctcaaggg ccagttcttc aaaattatgc cttcatttca 540
gcaggagaga cacttactct gaaagatttt tcgagtttga tgtttctcga aaatgtttct 600
tgcgagagaa aggggaatgat ctcaaggaaa accgtgagta tttccggagc aggcgaagt 660
attttttggg ataactctgt ggggtattct cctttgtcta ttgtgccagc atcgactcca 720
actcctccag caccagcacc agctcctgct gcttcaagct ctttatctcc aacagttagt 780
gatgctcgga aagggtctat tttttctgta gagactagt ttggagatctc aggcgtcaaa 840
aaaggggtca tgttcgataa taatgccggg aatttttgaa cagtttttcg aggtaatagt 900
aataataatg ctggtagtgg gggtagtggg tctgtctaaa caccaagttt tacagttaaa 960
aactgtaaa ggaagttttc tttcacagat aacgtagcct cctgtggagg cggagtagtc 1020
tacaaggaa ctgtgctttt caaagacaat gaaggaggca tattcttccg agggaacaca 1080
gcatacagat atttagggat tcttgcctgt actagtcggg atcagaatac ggagacagga 1140
ggcggtggag gagttatttg ctctccagat gattctgtaa agtttgaagg caataaagg 1200
tctattgttt ttgattacaa ctttgcaaaa ggcagaggcg gaagcatcct aacgaaagaa 1260
ttctctcttg tagcagatga ttcggttgtc tttagtaaca atacagcaga aaaaggcggt 1320
ggagctattt atgctcctac tatcgatata agcacgaatg gaggatcgat tctgtttgaa 1380
agaaaccgag ctgcagaagg aggcgccatc tgcgtgagtg aagcaagctc tggttcaact 1440
ggaaatctta ctttaagcgc ttctgatggg gatattgttt tttctgggaa tatgacgagt 1500
gatcgtcctg gagagcgcag cgcagcaaga atottaagt atggaacgac tgtttcttta 1560
aatgcttccg gactatcgaa cctgatcttt tatgatcctg tagtacaata taattcagca 1620
gcgggtgcat cgacaccatc accatcttct tcttctatgc ctggtgctgt cacgattaat 1680
cagtcgcgta atggatctgt gatttttacc gcagagtcac tgactccttc agaaaaactt 1740
caagttctta actctacttc taacttccca ggagctctga ctgtgtcagg aggggagttg 1800
gttgtgacgg aaggagctac ctttaactact gggaccatta cagccacctc tggctcgtgc 1860
cg

```

<210> 45

<211> 1668

<212> DNA

<213> Chlamydia trachomatis

<400> 45

```

agaaaatccg atagcagaaa tagaagaatt cgatgtgggt gcaacaaag ctcaagattg 60
ggatgtcgat gtagctatgt caaattcttt tgggtttggc ggacacaatt caacgatatt 120
attttcgagg tatgaacctt cattatgatg aaaactaagc acgaatatcc ttttggcggt 180
attcctatca gatttttttg tactccggat agaaglacct taaaggcctg ttttatctgc 240
catacagatg ggaacatttg gggtttccct aaggggcatg ctgaggaaaa agaaggccct 300
caggaagctg ctgagagaga acttgtagaa gaaactggtt tggggattgt taattttttc 360
ccaaaaatat ttgtgaaaaa ttattccttt aatgacaaag aagaaatctt tgtacgtaaa 420
gaggtaactt attttcttgc agaggttaaa ggcgaagtac atgctgatcc tgatgagatc 480
tgtgatgtgc agtggctaag ctttcaagaa ggtttacgcc ttttaaatcc occagaaatt 540
cgtaatatgg ttacggaagc agatgaattt gttcaaaagt atctatttgc ttcataaagt 600
cccctaggat gaaaaaaact tgggttaggag gggccgttgt ggaatctccc acaacagcct 660
tttctttttc tgcgatttta cataaaaaaga ttgcaatagt cttcgtgagc aagacgaatg 720
actttttgag cttctttttt gccgtataaa cctacaattt caatttttagc tgggttttgt 780
tgaattaaag tttctggagt agcttttatag gtttaagaaat agtgttggat catgtccaaa 840
actgtgcctg ggcattcaga aatatcttct atattgccat agactaaatc atcttctaga 900
acagcgatga ttttatcatc ggcttcttcc gagtctaaaa tacgaatccc tccgatagga 960
cgcgcttgca agaggatgtt cccttgtgta atattttttt ccgttaacac acagatatca 1020
agaggatcgc catcgccctt gatattctct ctgttacttt gttgaccact gtattctcca 1080
gaaagatctc cacaaatagt cttaggtaac agcccgata agcaaggaca aaagttagaa 1140
aacttttgtg gccatccac ttttaggata ccagtttctt tatccagttc gaatttaacg 1200
gagtcggctg gagtgatttc tatatagcaa caaagagatt cataatcatc gcgtgttaat 1260
actggcccat gccaaaggat agctatggat aatgtgtgtt tagacataag atcactctct 1320
attaaagtgt tttatgcgca attatcctgc gcatccggct tattcgtcca gatagtttta 1380
gtcttctgtt tccgcagtaa aacttttatt ttatcggcag cttttctttt tgctttttat 1440
cttgtcattg tgaaaaatgt tgaaaagtta ctcgtggcaa ctttccagac aggtttttttg 1500

```

taogaaagac	gagagtgatt	gtactgcaaa	ataatatgag	ccggacgtag	gatatgaaat	1560
actcttttgca	aatagaagac	ctacatatgt	aaggatatga	acagggtttg	aaagttactt	1620
gcgagtctgt	acagtttagt	gctgtaattg	ctattcatca	gacaaaag		1668

<210> 46

<211> 2010

<212> DNA

<213> Chlamydia trachomatis

<400> 46

atatcaaaagt	tgggcaaatg	acagagccgc	tcaaggacca	gcaaataatc	cttggggacaa	60
catcaaacacc	tgtcgcagcc	aaaatgacag	cttctgatgg	aatatcttta	acagtcctcca	120
ataattcatc	aaccaatgct	tctattacaa	ttggtttgga	tgcggaaaaa	gcttaccagc	180
ttattctaga	aaagttagga	gatcaaatcc	ttgatggaa	tgctgatact	attgttgata	240
gtacagtcca	agatatttta	gacaaaatca	aaacagaccc	ttctctaggt	ttgttgaaag	300
cttttaacaa	ctttccaatc	actaataaaa	ttcaatgcaa	cgggttattc	actccagta	360
acattgaaac	tttattagga	ggaactgaaa	taggaaaatt	cacagtcaca	cccaaaagct	420
ctgggagcat	gttcttagtc	tcagcagata	ttattgcatc	aagaatggaa	ggcggcggtg	480
ttctagcttt	ggtacgagaa	ggtgattcta	agccctgcgc	gattagttaa	ggatactcat	540
caggcatttc	taatttatgt	agtctaagaa	ccagtattac	taatacagga	ttgactccga	600
caacgtattc	attacgtgta	ggcggtttag	aaagcgggtg	ggtatgggtt	aatgcccttt	660
ctaattggcaa	tgatatttta	ggaataacaa	atacttctaa	tgtatctttt	ttagaggtaa	720
taacctcaaac	aaacgcttaa	acaattttta	ttggattttt	cttatagggt	ttatatattag	780
agaaaacagt	tcgaattacg	gggtttgtta	tgcaaaaata	aagaaaagtg	agggacgatt	840
ttattaaaaat	tggtaaaagat	gtgaaaaaag	atttccccga	attagaccta	aaaaacagag	900
taaacaaagga	aaaagtaact	ttcttaaaat	ctcccttaga	actctaccat	aaaagtgtct	960
cactaattct	aggactgctt	caacaaatag	aaaactcttt	aggattattc	ccagactctc	1020
ctgttcttga	aaaatttagag	gataacagtt	taaagctaaa	aaaggctttg	attatgctta	1080
tcttgtctag	aaaagacatg	ttttccaagg	ctgaatagac	aacttactct	aacgttggag	1140
ttgatttgca	caccttagtt	ttttgctctt	ttaaggggag	aactggaaaa	acaacacttt	1200
ctctaaacgt	gggatgcaac	ttggcccaat	tttttagggaa	aaaagtgtta	cttgotgacc	1260
tagaccogca	atccaattta	tcttctggat	tggggggctag	tgtcagaagt	gaccaaaaag	1320
gcttgacga	catagtatac	acatcaaacg	atttaaaatc	aatcatttgc	gaaacaaaaa	1380
aagatagtgt	ggacctaat	cctgcatcat	tttcatccga	acagtttaga	gaattggata	1440
ttcatagagg	acctagtaac	aacttaagat	tatttctgaa	tgagtactgc	gctccttttt	1500
atgacatctg	cataatagac	actccacctc	gcctaggagg	gttaacgaaa	gaagcttttg	1560
ttgcaggaga	caaattaatt	gcttgtttta	ctccagaacc	tttttctatt	ctaggggttac	1620
aaaagatacg	tgaattctta	agttcggctg	gaaaacctga	agaagaacac	attcttggaa	1680
tagctttgtc	tttttgggat	gacgtaact	cgactaacca	aatgtatata	gacattatcg	1740
agtcatttta	caaaaacaag	cttttttcaa	caaaaattcg	tcgagatatt	tctctcagcc	1800
gttctcttct	taaagaagat	tctgtagcta	atgtctatcc	aaattctagg	gccgcagaag	1860
atattctgaa	gttaacgcat	gaaatagcaa	atattttgca	tatcgaatat	gaacgagatt	1920
actctcagag	gacaacgtga	acaaactaaa	aaaagaagcg	gatgtctttt	ttaaaaaaa	1980
tcaaaactgcc	gcttctctag	attttaagaa				2010

<210> 47

<211> 2044

<212> DNA

<213> Chlamydia trachomatis

<400> 47

gtcatcaaga	aaagattggg	aacctatccg	tagtttgggt	aaagagcatg	gtatgcgaca	60
ttgtcagctt	atggctatag	ctccgacagc	gacgatctcc	aacattatag	gagtaactca	120
atctattgag	ccaacgtaca	aacatttggt	tgtgaagtct	aatttgcctg	gagaattcac	180
gattccaaat	gtgtatttta	ttgagaagtt	gaagaaatta	ggtatctggg	atgctgatat	240
gttagatgac	ctgaaatatt	ttgatgggtc	tttattggaa	atcgagcgta	taccagatca	300
cttaaaacat	attttcttga	cagcttttga	gattgaacca	gaatggatta	tcgaatgcgc	360
gtctcgaaga	caaaaatgga	ttgatattgg	gcaatccctc	aacctttatc	ttgcccagcc	420
agacgggaaa	aaactgtcga	atatgtattt	aacggcttgg	aaaaaagggt	tgaaaactac	480
gtattatctg	agatcttcat	cagcaacgac	cgttgaaaaa	tctttttagt	atattaataa	540
gagaggaaat	cagctctgtt	ggatgaagaa	taagtctgct	tcggcaggaa	ttattgttga	600
aagagcgaag	aaagcacctg	tctgttcttt	ggaagaaggg	tgtgaagcat	gtcagtaatt	660


```

aatcatataa attaacaata aaattaacgg ttottatgca agcagatatt ttagatggaa 720
aacagaaaacg cgttaaatcta aatagcaagc gtctagttaa ctgcaaccag gtogatgtca 780
accaacttgt tctattaag tacaatggg cttgggaaca ttatttgaat ggctgcgcaa 840
ataactggct ccctacagag atcccatgg ggaagacat cgaattatgg aagtcggatc 900
gtctttctga agatgagcgg cgagtcattc ttttgaattt aggttttttc agcaccgcag 960
agagcttggt tgggaataat attgttctag caatttttaa acatgtaact aatccggaag 1020
cgagacaata tcttttaaga caagcttttg aagaagcggg tcacacgcac acatttttgt 1080
atatttgtga gtcactcgga ttagacgaga aagaaatttt caatgcctat aacgagcgtg 1140
ctgcgattaa ggccaaagat gatttccaga tggaaatcac tggcaaggta ttggatccta 1200
attttcgcac ggactctgtt gagggtctac aggagtttgt taaaaactta gtaggatact 1260
acatcattat ggaagggtt ttctctata gtgggtttgt gatgatcctt tccttccaca 1320
gacaaaataa gatgattggg attggagaac aatatcaata catcttaaga gatgagacaa 1380
tccacttgaa ctttgggtatt gatttgatca acgggataaa agaagagAAC ccggggattt 1440
ggactccaga gttacagcaa gaaattgtcg aattaattaa gcgagctgtc gatttagaaa 1500
ttgagtatgc gcaagactgt ctccctagag ggattttggg attgagagct tcgatgttca 1560
tcgattatgt gcagcatatt gcagaccgtc gtttgaaag aatcggatta aaacctattt 1620
atcatacgaa aaacccattc cttggatga gcgaacaat agacctaat aaagagaaaa 1680
acttctttga aacaagggtt atagaatatc aacatgcagc aagcttaact tggtagtcct 1740
gatatacaaaa taggagaaaag cctcaaccat agagttgagg cttttttttg tcatacggta 1800
acctgataag aattttttaga ttttcagggtt tttattaaaa tggaaaaggc tagtcacctc 1860
tttaattttc tcataatatc ttgtagccct tgggtgagatt ccttttcgcac 1920
tctatgact actgttagag tgggtgagatt tgggggttga gcaggtgtag ccttttcgcac 1980
acgaagtatt ttctgtgaa accacaagat ttgaaacttc cctatttttg ggaagaacgt 2040
tctc 2044

```

<210> 48

<211> 3734

<212> DNA

<213> Chlamydia trachomatis

<400> 48

```

gttattcgct tctactocat tagaagtccc taatgctaaa ctcaccattt ttctctcttt 60
ccgttaaaac aggaagaaa ttgtacagaa acattttttt aaagaaatca aaaagccatt 120
tgcaggcaga tatcaggcca ttatatcaa aaacagaaag aatgattagg ataaaacttt 180
gtcttgccat cgttccagag agcattgaga agcgtttttt attataaata cattgacta 240
agaatcttaa aatcgaaacag acaacacaat ggctcgaaca gactgatcca cagcactaa 300
ttcaaatgca aaaaacttct aaaatgaaca cagcaagctt gataaaaaa taaaaagaa 360
ttggatcata gagcttttac agaaggggag cactgcaatc tgtctcgacc aaatagcaat 420
gcaaacagat aaatacccct aatcattggg aaaaattgag tgtagaatag cctctttctc 480
ttctctctat tgttgcctag ctaacgcgat ttctctttta gagatatctg caagtctctg 540
cttatccaaa aagccttgtc tctcattttc caatacaaat ctgtccagag aaactttttt 600
tggctctcca ccatagctag aaattctagt aagaacagca cctagcatca cagatccaaa 660
aacaaccagg gtaaccacta cgtcaatcat aggaagcgta gtccaacctg cccaataaaa 720
taaggctgct cctgtaacta tgaataaaat actaagaata ccgagcgcaa gcacagcaat 780
acgttcgcta caacaagaaa ctctcgcttt agaagcgcta tccaccaaag gagcctctgg 840
catataactt ctaagaggta cactatctcc aacaaaactc atggcatccc ccttaaggta 900
aaagagaagc ttctctctaa atagaaaagc gtatcgtoaa ctcttttata gatctaaaaa 960
gtcttgcttt ccttaatccc acccatgaaa tttagcataa aaacctcca acatattcac 1020
acgtctctct aaaaaggcta tttccctatt tttctgagtc tctaaaaacc tataatggct 1080
ggaaattttc cgcgcacttt ccttggcttc ttgtaatagc tgatctgaat tgcgtatcac 1140
agataacagg taagaaacta atccaaaagc tcctatacaa gaaccaataa ttgcagctct 1200
cccactactc ctaaaactaa ggaagaatag actcccccac gacaaagaaa aactcctcct 1260
aaagctgcaa gcaacttgt tagaacaact acaaaataact ggtatgtttt agaacggtga 1320
ataaaggagt tgttagccac attttctactg tacctcagtt tttgctgaac aacaattccc 1380
taaaaaattg gtaggacgcc aaacgttcat aattactcta cttggaaacc attaatatt 1440
atatcagact ttcttccaat acacatttca acccactttg aagctgttct atttttttt 1500
gagcaagctc taaatctttg ctcttttgag caagcaatcc ttcaacttct ttcaaatctt 1560
cttctgcttc atatagaagt tottgataag ataacaacta tccaggagtc acggcctctg 1620
gagctaactc agatgactct gaagggtgct tctgtcggtt taagaaaac ccatacatat 1680
aaactagact tctctctata caggcagaac ccagtgctat tgctaataag ctaagaatag 1740
gagcaaaaag agagaccaca cttcctgaaa aaagaagcag aagagcacca cctaaaactg 1800
ctagtacccc taataccaag gcacctattg ccaacaattg ctctttacg cttgtagtag 1860

```

```

tctgagcacc gatagtttca gtatgatcgg cagcgaatgg tttgctggaa ttacaacaaa 1920
aagaaaatatt aaacatggcg cctctatttcc gcaaaaaaaa ggccaacatg ctacaggaaa 1980
gctaattaaa gtaaaaatatt ttatatattt caatggtagt taaataccta atctacccaa 2040
ccaaaagatg tctaaatgac aaaaaaataa togtatttat attatcatga gacacttata 2100
gtcacgtctg cttcattcag ctcaaattct aatgaaaaat cggatttaga agaaaataga 2160
ctogaagagt cagaactagc caaaatgttt gttctaattc tattttgcaa tccccgacta 2220
caagaccaat agagaaacgt taacctact cctaaagcca cagaaccaat cataatcgct 2280
ccaataccta aacoggcaaa cacaagcgac gatccccgc aaagcaaaca aagcaaggct 2340
acacaactta aaatagcaaa aattcctaag gaaacggcaa attctatatt tctcttctgt 2400
ttgcaataaa tatgctgttt atacagacac aactctgcgg ggctctccag agttggagcg 2460
caagaggaaac aaaaaagata agacattgtc gactccggac caaaaaaagg cgagataata 2520
cgcgagatgg taaaaataca gaaatatttt tgacatagaa aaccttaacc ctcttttcat 2580
cgcgtagac tagagtgtaa aacaagatgc gaaagcaagg ttcgctatgt ttggaaacaa 2640
acctccacac ggtcccggtat tatcaaaaaca agtcttccag ggatatgtta gagaacgtcc 2700
tatccatacc aaagcaacat atagacgtct tttgtgaaaa gactgaatag aggaatctaa 2760
gaagcttgggt tagcgtctat agatgcttta agagcagctt tttccttttc agcactatcc 2820
aaccatcttg tgtagctaga taaaactaag cgcacatcgg acaataaagc ttgctcattt 2880
ttctctaatac tgtccaaaaca atcaatctca acttctattg ccttagcttc caaagcttgg 2940
agatcgctcg taagacctcg cagaaacatc ttattaatga aagagacgga gaccaaagcg 3000
tcttctcttt ctgaaagatt acgcaaacgt tgctcagcca aaacattttt tgcttctaag 3060
ctagcataag aggatcgaca cataagacga gatattcccg caccacaca agcagatcca 3120
ataattaatg cagcaatacc tattgcagta aatatgacat tgctagcgca caaaacccaa 3180
gctaataccc cagcgacaac aactaaagcg cctacgatag ctaaaagctat atccaaaatt 3240
ttggaacaag tattcccttt tgttgaagac gaagtagatt ttatctctac gcaggaagct 3300
gttggcaatg tcaagaaga agcgtctccg ctaaatagtag tactcatttt tccacatttt 3360
tattttttaa acggaaaaac tgtatcagaa cggcgcttta ttcgcaaatc attataaatc 3420
cgcaacatgc agaactaaag cgcgtaagc aaaaggaacc cctaactctc agatgcaata 3480
tctgaggagt ctttaattat tttttacgac gggatgcctg cacctgcagc cgctctgata 3540
atgtcttatt ctacagatctc aatttacaca actctgctgt taattgactg caagtgttct 3600
gactttgttg caaccgctgt ttaaaccctt ctgtctgatg acgaatttct tgttcagcat 3660
cctcctcaat ggagcaaac gtttcggcat aacgcttaca caaatctaat atttgttctt 3720
ccaactcttg gcaa 3734

```

<210> 49

<211> 2937

<212> DNA

<213> Chlamydia pneumoniae

<400> 49

```

atgcctcttt ctttcaaatac ttcatctttt tgtctacttg cctgtttatg tagtgcaagt 60
tgcgcggtttg ctgagactag actcggaggg aactttgttc ctccaattac gaatcagggt 120
gaagagatct tactcaacttc agatttttgt tgttcaaact tcttgggggc gattttttca 180
agttccttta tcaatagtttc cagcaatctc tccttattag ggaagggcct ttcttaacg 240
tttacctctt gtcaagctcc tacaaatagt aactatgcgc tactttctgc cgcagagact 300
ctgaccttca agaatttttc ttctataaac tttacaggga accaatcgac aggacttggc 360
ggcctcatct acggaaaaga tattgttttc caatctatca aagatttgat cttcactacg 420
aaccgtgttg cctattctcc agcatctgta actacgtcgg caactccgc aatcactaca 480
gtaactacag gagcctctgc tctccaacct acagactcac tcaactgtcg aaacatatcc 540
caatcgatca agtttttttg gaaccttgcc aacttcggct ctgcaattag cagtctctcc 600
acggcagctg ttaaattcat caataacacc gctaccatga gcttctccca taactttact 660
tcgtcaggag gcggcgtgat ttatggagga agctctctcc tttttgaaaa caattctgga 720
tgcacatctc tcaacgcaa ctctgtgtg aacagcttaa aaggcgtcac ccttctca 780
ggaacctatg ctttaggaag tggcggagcc atctgcatcc ctacgggaac tttcgaatta 840
aaaaacaatc aggggaagtg caocttctct tataatggta caccaaatga tgcgggtg 900
atctacgcgg aaacctgcaa catcgtaggg aaccagggtg ccttgctcct agatagcaac 960
actcagcgga gaaatggcg agccatctgt gctaaagtgc tcaatattca aggacgggt 1020
cctattgaat tctctagaaa ccgcgcggag aagggtggag ctattttcat aggccctct 1080
gttgagagac ctgcgaagca aacatcgaca cttacgattt tggcttccga aggtgatatt 1140
gcgttccaag gaacatgct caatacaaaa cctggaatcc gcaatgccat cactgtagaa 1200
gcagggggag agattgtgtc tctatctgca caaggaggct cagctcttgt attttatgat 1260
cccattacac atagcctccc aaccacaagt ccgtctaata aagacattac aatcaacgct 1320

```

```

aatggcgctt caggatctgt agtctttaca agtaaggagc tctcctctac agaactcctg 1380
ttgcctgcca acacgacaac tatacttcta ggaacagtca agatcgctag tggagaactg 1440
aagattactg acaatgcggt tgtcaatggt cttgggttcg ctactcaggg ctcaggtcag 1500
cttaccctgg gctctggagg aaccttaggg ctggcaacac ccacgggagc acctgccgct 1560
gtagacttta cgattggaaa gttagcattc gatccttttt ccttcctaaa aagagatttt 1620
gtttcagcat cagtaaatgc aggcacaaaa aacgtcactt taacaggagc tctggttctt 1680
gatgaacatg acgttacaga tctttatgat atggtgtcat tacaatctcc agtagcaatt 1740
cctatcgctg ttttcaaagg agcaaccgtt actaagacag gatttcctga tggggagatt 1800
gcgactccaa gccactacgg ctaccaagga aagtggtcct acacatggtc ccgtcccctg 1860
ttaattccag ctctgatgg aggatttcct ggaggtccct ctctagcgc aaatactctc 1920
tatgctgtat ggaattcaga cactctcgtg cgttctacct atatcttaga tcccagagct 1980
tacggagaaa ttgtcagcaa cagcttatgg atttcttct taggaaatca ggcatctct 2040
gatattctcc aagatgttct tttgatagat catcccggtt tgtccataac cgcgaaagct 2100
ttaggagcct atgtcgaaca cacaccaaga caaggacatg agggcttttc aggtcgctat 2160
ggaggctacc aagctgcgct atctatgaac tacacggacc acactacgtt aggactttct 2220
ttcgggcagc tttatggaaa aactaacgcc aaccctacg attcacgttg ctcagaacaa 2280
atgtatttac tctcgttctt tgggtcaattc cctatcgtga ctcaaaagag cgaggcctta 2340
atttcttgga aagcagctta tggttattcc aaaaatcacc taaataccac ctacctcaga 2400
cctgacaaag ctccaaaatc tcaagggcaa tggcataaca atagttacta tgttcttatt 2460
tctgcagaac atcctttcct aaactggtgt cttcttacia gacctctggc tcaagcttgg 2520
gatctttcag gttttatttc cgcagaattc ctagggtggt ggcaaaagtaa gttcacagaa 2580
actggagatc tgcaacgtag ctttagtaga ggtaaagggt acaatgtttc cctaccgata 2640
ggatgttctt ctcaatggtt cacaccattt aagaaggctc cttctacact gaccatcaaa 2700
cttgccctaca agcctgatat ctatcgtgtc aaccctcaca atattgtgac tgtcgtctca 2760
aaccaagaga gcacttcgat ctcaggagca aatctacgcc gccacggttt gtttgtacaa 2820
atccatgatg tagtagatct caccgaggac actcaggcct ttctaaacta tacctttgac 2880
gggaaaaatg gatttacaaa ccaccgagtg tctacaggac taaaatccac atttttaa 2937

```

<210> 50
 <211> 801
 <212> DNA
 <213> *Chlamydia pneumoniae*

```

<400> 50
atgcattcaa aatttctttc tcgaagaaaa aaaaatagtt ctcataagga ggaaacctct 60
tgggattgta tagcctcaag ttacaataag atagtccaag ataaagggca ctactatcat 120
agagaaacta tcttccccc actcctgcct tcactcacct taggttcaaa aagttctgta 180
ttggatattg gctgcggtca aggtttttta gaaagggcc ttcttaagga atgtogttat 240
ctaggcatag atatctcttc tagattgatt gctctagcaa agaaaatgcg atcggtaaac 300
tctcatcagt ttaagggtgc agatcttagc aaacgcctag agttcgtaga accgacatta 360
ttctctcatg cagtagcaat cctctccctt caaaatatgg aattccccgg agaggctata 420
cgtaatacac ctacgctcct cgaaccactc gggcaatttt ttatagtttt aaaccatcct 480
tgttttcgta ttctagggc atcatcctgg cactatgatg aaaataaaaa agctatctct 540
cgtcatatag atcgttatct ctccccaatg aaaatcccaa tcatggetca cccaggacaa 600
aaagattcgc cttctaccct ctctttcac tttcctctaa gctattgggt taaagaactg 660
tcttctcatg gattcttagt ttcagggtctt gaggaatgga catcttcaaa aacctcaaca 720
ggaaaacgag ctaaggcaga aaacctttgt cgaaaggaat ttccattatt ccttatgatt 780
tcatgcatta agataaaaata a
801

```

<210> 51
 <211> 252
 <212> DNA
 <213> *Chlamydia pneumoniae*

```

<400> 51
atgaaacaac aacacaatcg taaggcttta tctcgcaaga ttggcacagt gaaaaacaa 60
gccaaatttg caggaagctt tttagatgag attaaaaaaa ttgaatgggt aagcaagcac 120
gatcttaaga aatacataaa agtagttctt atcagtattt ttggttttgg atttgctatt 180

```

tatttcgtag atcttggtgtt gcgtaagtca atcacatggt tagatggtat aacaaccttt 240
 ttgttcggtt aa 252

<210> 52

<211> 1185

<212> DNA

<213> Chlamydia pneumoniae

<400> 52

atgtcaaaag aaacttttca acgtaataag ccccatatca atattgggac gatcggggcac 60
 gttgaccatg gtaaaactac gctaacagcg gcaattacac gcgcgctatc aggggatgga 120
 ttggcctctt tccgtgacta tagttcaatt gacaatactc cagaagaaaa ggctcgtgga 180
 attactatca acgcttctca cgttgaatac gaaaccccaa atcgtcacta cgctcacgta 240
 gactgccctg gtcacgctga ctatgttaaa aatatgatta caggcgccgc tcaaatggac 300
 ggagctatcc tagtcgtttc agctacagac ggagctatgc cacaaactaa agaacatata 360
 ttgctagctc gccaggttgg agttccttat atcgttggtt tcttgaataa agtagatatg 420
 atctctcaag aagatgctga acttattgac cttgttgaga tggaaacttag tgagcttctt 480
 gaagaaaaag gctacaaagg atgcctatt atcgttggtt ctgctttgaa agctcttgaa 540
 ggtgatgcaa attatatcga aaaagtctga gaacttatgc aagctgtgga tgacaacatc 600
 cctacaccag aaagagaaat tgataagcct ttcttaatgc ctatcgaaga cgtattctca 660
 atctctggtc gtggtactgt gttacagga agaatcgagc gtggaatcgt taaagtctct 720
 gataaagttc agctcgtggg attaggagag actaaagaaa caatcgttac tggagtgcga 780
 atgttcagga aagaacttcc tgaaggtcgt gcaggagaaa acgttggttt actcctcaga 840
 ggtattggaa agaacgatgt tgaagaggt atggttggtt gtcagcctaa cagcgtgaag 900
 cctcatcaga aatttaagtc agctgtttac gttcttcaga aagaagaagg cggacgtcat 960
 aagcctttct tcagcgata cagacctcag ttcttcttcc gtactacaga cgtgacagga 1020
 gtgcgtaactc ttctgaagg aactgaaatg gtaatgcctg gagataacgt tgagcttgat 1080
 gttgagctca ttggaacagt tgctcttgaa gaaggaatga gatttgcaat tcgtgaagggt 1140
 ggtcgtacta tcggcgctgg aacgatttca aagatcaatg cttaa 1185

<210> 53

<211> 1431

<212> DNA

<213> Chlamydia pneumoniae

<400> 53

atgagaatcg tacaagtcgc tgtagaattc actccaatcg ttaaagtagg cggctctaggc 60
 gatgctgtag ctagtctatc taaggagtta gcgaaacaaa atgatgtgga agtacttctc 120
 cctcattatc ctttaatttc caaattctct togtctcaag ttctttccga gcgttctttc 180
 tattatgaat ttttaaggcaa gcagcaagcc totgcaattt cttattctta cgagggtctt 240
 acgcttacta taattacgtt ggattcacaa atagagcttt tctcaaccac gtccgtgtac 300
 tctgagaata atgttgtaacg tttctctgct tttgcagctg cagctgcagc ttatcttcaa 360
 gaagcggatc ctgctgacat tgtgcacttg catgactggc atgtaggttt acttgcgggt 420
 ttattaaaaa accctttaaa cctgtgcat tccaagattg tctttactat ccataatttt 480
 ggttatcgag ggtattgtag tacgcagcta ttagcagcgt cgcaaattga tgattttcat 540
 ttgagtcact accaactatt tcgcgatccg caaacttctg ttctaataaa gggagctctc 600
 tattgttcgg attacattac gacagtgtct cttacttatg tgcaggaaat tataaacgac 660
 tattctgatt acgaacttca tgatgcgatt ctagcaagaa attctgtatt ttctgggac 720
 atcaaatggca ttgatgaaga cgtttggaac cogaagacag atcctgcttt agctgtacag 780
 tacgatgcaa gcctattaag cgaacctgac gttctcttta ctaaaaaaga agagaacaga 840
 gcggtattat atgagaagtt ggggatcagt tcagactatt ttctttgat ttgtgtgac 900
 tcacgcattg ttgaggaaaa ggtcctgaa tttatgaaag agattattct ccatgctatg 960
 gagcacagt atgcctttat cttgattggg acaagtcaaa atgaggttct tottaatgag 1020
 ttccgttaact tacaagattg tttagcgagc tcccccaaca ttctgttgat cttggacttt 1080
 aatgatcctt tagccaggct aacttatgct gotgccgata tgatctgcat cccttcacat 1140
 agggaggctt gtggacttac ccagctgata gcgatgcgtt atggcacagt tcttttagtt 1200
 cgtaaaaactg gagggcttgc tgatacagtg attcctgggg taaatggttt cactttcttt 1260
 gatacaaaac attttaatga atttcgggct atgcttagca acgctgtaac gacgtatcgt 1320
 caggagcctg acgttttggtt gaatttgatt gactcgggaa tgcttcgggc ctctggctta 1380
 gatgccatgg ctaagcatta cgtaaatctt tatcaatctt tactctcatg a 1431

<210> 54
 <211> 1041
 <212> DNA
 <213> *Chlamydia pneumoniae*

<400> 54
 atggaagcag atatttttaga tggaaagctc aaacgggttg aggtaagtaa aaaaggattg 60
 gtgaattgta atcaagtaga tgtcaatcag ctagtccta tcaagtataa atgggcttg 120
 gaacattacc tcaatggatg tgcaaacac tggcttcta ctgaagtcc tatggcaaga 180
 gatatcgagt tgtggaaatc agatgaactg tctgaagacg aacgcagggc ctttttgta 240
 aacctaggat ttttcagtac cgcggaaagc ctagtcggaa ataacatcgt tcttgctatc 300
 ttcaaacata tcacaaaccc tgaagcaaga cagtatttac tgcgtcaagc ttttgaggaa 360
 gccgtacata cacatacatt tctctatatt tgcgaatctt taggacttga tgaaggcgaa 420
 gtattcaatg cctataatga aagagcctca attagggcta aagatgattt tcaaatgaca 480
 ttaacagtcg atgtccttga tcttaatttt tctgtacagt cttcagaagg cttggggcag 540
 ttcattaaaa acttagtagg atactatatac attatggaag gaatcttctt ctatagtgg 600
 tttgtaatga ttctctcttt ccatagacaa aataaaatga cagggaattg agaacagtac 660
 caatacatcc tcagagatga aaccatacat ttaaattttg gaatcgatct tatcaatgga 720
 attaaagaag aaaaccccgga agtttggact acggaactac aagaagaaat cgtcgtctt 780
 attgaaaaag ctgtagagct tgaatttgag tacgctaaag attgcttacc tcgaggaaac 840
 ttgggattaa gatcttcgat gtttatagat tacgttcgtc atattgcaga tcgtcgttta 900
 gagagaattg ggttgaagcc tatctatcac tccagaaatc ctttcccttg gatgagcgaa 960
 accatggatc tgaataaaga aaagaatttc tttgaaaccc gggttaccga ataccaaacc 1020
 gctggtaatt taagttggta a 1041

<210> 55
 <211> 3135
 <212> DNA
 <213> *Chlamydia pneumoniae*

<400> 55
 atgggtcgaag ttgaagaaaa gcattacacc atcgtcaaac gtaatggaat gtttgtccca 60
 tttaatcaag atcggatttt ccaggctttt gaggcagctt ttcgagatac gcgtagctta 120
 gaaactagtt tccactacc taaagactta gaagaatcta ttgcgcaaat tactcataaa 180
 gtcgtgaagg aagtcctcgc taaaatttca gaaggtcagg tagtactgt agagagaatc 240
 caggatcttg tagaaagtca gctctatatt agcgggttgc aggatgtggc tcgcgattat 300
 attgtttaca gggaccaacg caaggcagag cgcggtaact cttcgtccat aattgccatc 360
 atactagatg acgggggaag cgctaaattt aatcctatga agatctctgc agctctcgaa 420
 aaagcattca gagcgacgct ccaaattaat gggatgactc ctctcgcaac actatccgaa 480
 attaatgacc ttacccttag gatcgttgaa gatgtcctaa gccttcatgg tgaagaagct 540
 attaatctgg aagagatoca agatattgtt gaaaagcaac ttatggttgc cggctattat 600
 gatgtggcca agaattatat tttatataga gaagctcgtg cagcagcccg tgctaataaa 660
 gatcaagatg gacaagaaga gtttgtcccc caagaggaaa cgtacgttgt tcaaaaagaa 720
 gacggcacca cctaccttct gagaaaaaca gatttagaaa agaggttttc ttgggcatgc 780
 aaacgcttcc ctaaaactac agattctcaa ctgcttgcag atatggcatt tatgaatttg 840
 tattcaggaa tcaagaaga cgaggtcacc acagcatgca tcatggcggc acgtgccaat 900
 atcgagagag aacctgatta cgcttttata gcagcagaac tctcacgag ttccctgtat 960
 gaagagacct taggatgcag ctctcaagac cccaatttat cagaaatata taaaaaacat 1020
 tttaaagaat acatcctcaa tggagaagag tatcgttga atcctcaatt aaaggattat 1080
 gatctcgatg ctcttagtga agtcttagac ctctctagag accaacagtt ttcttatatg 1140
 ggagtccaaa atctctacga tgcgtatttt aatctgcag aaggacgacg ttagagact 1200
 gcgcagatct tttggatgcg ggtttctatg ggcttagcct taaatgaagg agaacaaaag 1260
 aatttttggg caatcacttt ctataatctg ttatccacat tccgtatata ccagcaact 1320
 cctacattgt ttaactccgg aatgcgtcat tcccaactca gttcatgcta tctttccaca 1380
 gtaaaagatg acctaaagtca cttttataag gtgattcttg ataattgctt gctttctaaa 1440
 tgggcagggg gaattggaaa tgattggaca gatgtccgtg ctacaggagc tgtaatttaag 1500
 ggaaccaatg gaaagagtca aggcgtcatt ccttcatta aggttgccaa tgatactgca 1560
 attgcagtga atcagggggg caaacgtaaa ggtgctatgt gcgtatatat agaaaactgg 1620
 cacttggatt acgaagactt tttagaattg cggaagaata caggagatga gcgtcgtaga 1680
 actcacgata tcaatacagc aagctggatt cctgatctct tctttaagag actagaaaaa 1740
 aaaggcatgt ggacactctt tagccccgat gatgtcccag gtttacacga agcctatggg 1800

```

ttagagtttg aaaagcttta tgaagaatat gaacgtaagg ttgaatctgg ggaaatccgt 1860
ctttataaaa aagtagaagc cgaagtgcgt tggcgtaaaa tgtaagcat gctttacgaa 1920
acagggcatc cttggattac atttaaagat ccttcgaata ttcgctcaaa ccaagatcat 1980
gttggcgtcg tacgctgttc taatctatgt acagagattt tattgaactg ttcggaatca 2040
gagactgcag tttgtaattt aggttccata aacttggtag aacatatccg taatgacaag 2100
ttagatgaag aaaaattaaa agaaactatc tcaatagcca tccgtatttt ggataacgtt 2160
attgacctga acttctaccc tacaccagag gctaaacaag ccaacctaac tcacagagct 2220
gtgggggttg gggttatggg attccaggat gttctttacg agttgaacat tagctatgcc 2280
tcacaagaag ctgtcgaatt ttctgacgag tgctcggaga tcatcgcata ctacgctatt 2340
ctagcctcga gcttactcgc gaaagaacga ggtacatatg cttcttattc aggatctaag 2400
tgggatcgtg ggtatctacc cttagatact atcgagcttc tcaaagaaac tcgcgagag 2460
cataatgttc ttgtagacac atcaagtaaa aaagattgga ctccagttcg tgatactatc 2520
cagaaatcac gaatgagaaa tagccagggtc atggcaattg ctctacagc aacgatctcg 2580
aatatcatag gggtcaccca atctatagag cccatgtata aacatctctt tgtaaagtcc 2640
aacctttccg gagagtttac gatccccaac acctacctga ttaaaaaact taaggaatta 2700
ggactttggg atgcagaaat gttagatgat ctaaaatatt ttgacggatc tctattggaa 2760
attgaaagga tccctaatac cttgaaaaag cttttcctta cggcatttga aatcgaaccc 2820
gagtggatta tagagtgtac ctctagaaga cagaaatgga ttgatatggg agtttctcta 2880
aatctgtatc ttgctgagcc agatggtaaa aaactctcca atatgtatct caccgcttgg 2940
aaaaaaggat taagactac ctattattta agatctcaag ctgcaacatc agtagagaaa 3000
tcatttatag atatcaataa acgcggcatt cagcctcggt ggatgaaaaa taaatcagcg 3060
tccacaagta ttgtggtcga aagaaaaaca acccccgttt gttcaatgga agaaggttgc 3120
gaatcttgtc aataa 3135

```

<210> 56

<211> 1386

<212> DNA

<213> Chlamydia pneumoniae

<400> 56

```

atgatgagct ctaagcgtac ctcgaaaata ggggtgcttt caattttatt aacatttact 60
cactctatag ggttcgcaaa tgcgaattcg tccgtaggtc ttggcacggg ctacattaca 120
tccgaggttg taaagaagcc tcagaaagga tcagaaagga aacaagccaa aaaagaacct 180
cgtgctcgta aaggatactt agtcccttct tcaaggactc tttcagctcg agcccaaaag 240
atgaaaaact cctctcgtaa agagtcttca ggtggttgta acgaaatttc tgcaaatct 300
acaccagat ctgtaaaatt acgaagaaac aaacgtgcag aacaaaaggc agctaaacaa 360
ggattttcag ctttttctaa cctaactttg aaaagcctac ttctaaaact tcttcaaaa 420
caaaaaactt caattcacga gagagaaaaa gcaacctcaa gatttggtta tgagtctcag 480
cttagttccg cagcaaaacg ctactgcaca ccatcttcag ccgctccttc cctattttta 540
gaaacagaaa tcgttcgagc tctgtagaa agaactaaag aacttcaaga taatgaaatt 600
catattcctg tagtgcaagt ccaaacgaac ccaaagaac aaaatacaaa gacaactaaa 660
cagttggcat cccaagcctc gattcaacaa tctgaaggaa ccgagcaatc attgcgagag 720
ctcgcccaag gtgctagcct acctgtctta gtgcgtctta atcctgaagt gtctgtacaa 780
agacaaaaag aagagttatt aaaagaactc gtagctgaac gtagacaatg taaaagaaa 840
tctgtaagac aagctcttga agctcgttct ttaactaaga aagttgctag aggcggttct 900
gtgacctoga ctttacgata cgatccagaa aaagcggcgg aaatcaaaag tagacgcaat 960
tgcaaaagtaa gtctgaagc acgtgaacaa aaatattcat cttgcaaaag agatgctcgc 1020
gctaattggga aacaagacaa gacaactcct agtgaagatg cttctcaaga agaacaacaa 1080
actggggcag gactcgtacg caagactcct aaatctcagg ttgcaagtaa tgctcagaac 1140
ttctaccgaa attotaaaaa tacaaacata gatagctatc ttacagctaa ccaatacagc 1200
tgtagttctg aagaacacaga ttggccatgt tcttctgcg tctctaaacg cagaactcac 1260
aacagtatat ctgtatgtac catggtagtt actgtcattg cgatgatcgt aggggcttgg 1320
attatagcta atgtacaga atctcaaaac acatcagatc caactcctcc aactcctact 1380
ccatag 1386

```

<210> 57

<211> 1731

<212> DNA

<213> Chlamydia pneumoniae

<400> 57

```

atgacagatt ttcctactca cttcaaagga cccaaactta accccattaa agtaaatacca 60
aacttttttg agaggaatcc taaagtcgca aggggtactgc aaattacagc cgtagtctta 120
ggaatcattg cctctttatc cggatatagta ctcatatag gcacccctct cggagctcct 180
ataagtatga tcctcggcgg atgtctttta gcttctggag gcgccttatt tgttgggtgt 240
acgattgcta cgatattgca agctagaaat agttataaga aggcctgtaa ccaaaagaaa 300
ctctcagagc ctttgatgga acgcccgaag ttgaaagcct tagattattc cctagatctg 360
aaagaggatg gggacctaca tcattctgtt gtcaaacatc ttaaaaaatt agacctgaat 420
ctttccaaaa cccaaaggga agttctaaat caaatcaaaa ttgatgatga gggacctctc 480
ctaggggaat gcgcctctat gatttcagaa aactacgacg catgcttaaa gatgctcgcg 540
tatcgtgagg agctcctgaa agaacaacc caataccaag agacacgatt caatcagaac 600
ctcactcata gaaataaagt tttgctctcc atoctctcaa ggatcacgga caatatttct 660
aaagcggggc ggggtcttttc tttgaaattt tccacgctaa gctcgcggat gtcacgaatt 720
cataccacca ccactgtgat tctggcttta agtgccgttg tttctgtcat ggtcgtagca 780
gctctaattc cagggtggcat tttagcacta cctatacttt tggctgttgc tatttctgca 840
ggagtgtattg tcacggact ttcctatcta gttcgtcaga ttttaagtaa caccaagcgt 900
aatcgtcagg atttttataa agattttgta aaaaatgtag atatagagct tcttaaccaa 960
acggtaactt tacagcgatt cctctttgaa atgctcaaag gtgttctgaa agaagaagaa 1020
gaagtctcct tagaaggta agattgggtat acacaatata taaccaatgc acccatagaa 1080
aaaagattga tcgaagagat cagagttacc tacaagaga tcgatgctca gaccaaaaaa 1140
atgaagacag acttggagtt cttagaaaat gaggtgcgtt ccgggagact gtctgtagcg 1200
tcccgcgtcg aagatccaag tgaactcct atttttactc aaggtaagga gtttgcaag 1260
ttacgtcgcc aaacctctca gaatatatcc acgatttatg gtccggacaa tgaaaatatt 1320
gatcccgaaat tttccttaac ctggatgcct aaaaaagaa aagaaataga ccatagctta 1380
gaacctgtta caaagttgga acccggttca agagaagagt tgttgttggg agaggggggc 1440
aaccacaact taagagaact caatatgaga attgcacttc tacaacaaca actatcaagt 1500
gtccgaaaat ggagacaccc tcgaggggaa cattacggga atgttatcta ttcagatata 1560
gaactcgatc gtattcagat gctagaaggc gcattttata atcacctcag ggaagctcaa 1620
gaggaaatca ccagctctct cgagacactt gttgacattc aaaaccgtat tttagggatc 1680
atagttgaag gggactcaga ttcaagaaca gaagaagagc cttaggaata g 1731

```

<210> 58

<211> 1086

<212> DNA

<213> Chlamydia pneumoniae

<400> 58

```

atgcaacaaa ctgtaattgt agcaatgtca ggaggcgtgg attctctctgt cgttcgctat 60
ttattcaaaa aatttaccaa ttataagggtt attggcctct tcatgaagaa ttgggaagag 120
gatagcgaag gcggcctttg ctctgtact aaagattatg aagatgtoga gagggatgt 180
cttcagctcg atatccctta ttacaccgta tcttttgcta aagaatatag agaaagagt 240
ttcgtctgtt tcctcaagga ataactctta ggctacactc ctaacccoga cattctttgt 300
aaccgagaaa tcaaatttga cttctacaa aagaaagtcc aggaacttgg cggagattac 360
ctcgtacag ggcactactg ccgattaaat accgagctcc aagaaaccca actccttaga 420
ggttgcgato ctcaaaaaga tcagagctat tttttatcag gaactcctaa aagtgcctct 480
cacaatgtgc tctttcctct tggggaaatg aataagactg aagttcgtgc gattgcagct 540
caagcagctc ttccacagc agaaaaaaa gatagtacag gcatttgctt tatagggaag 600
cgccctttta aagagttcct agagaagttt cttcccaata aaacaggcaa cgttatcgat 660
tgggatacca aggaattgt agggcaacat caggagctc actattatac tatagggcag 720
cggcaggac ttgatcttgg aggatccgag aaaccctgtt atgttgtggg aaaaaatata 780
gaggaaaata gcatttatat tgtgaggggg gaagaccatc ccagctcta cctacgggaa 840
ttaacagcta gagagctcaa ttggtttacc cctcctaaat ccgatgtca ctgtagcgct 900
aaagtccgct accgttctcc tgatgaagct tgcacgatag attatagctc aggtgacgag 960
gtcaaggtgc gatittcaca acccgtcaag gcggtaaact caggacaaac aatagcgttt 1020
tatcaaggag atacctgcct tggtagtgga gttatcgacg ttctatgat tccaagtga 1080
ggctag 1086

```

<210> 59

<211> 4830

<212> DNA

<213> Chlamydia pneumoniae

<400> 59

```

atggttagcga aaaaaacagt acgatcttat aggtotctcat tttotcattc cgtaatatga 60
gcaatatgtg cagcaggcat tgcttttgaa gcacattcct tacacagctc agaactagat 120
ttagggtgtat tcaataaaca gtttgaggaa cattotgctc atgttgaaga ggctcaaca 180
tctgttttaa agggatcaga tcctgtaaat ccctctcaga aagaatccga gaagggtttg 240
tacctcaag tgctctttac ccaaggaaagc tctggagaga gtttgatct cgcgatgct 300
aatttcttag agcattttca gcatcttttt gaagagacta cagtatttgg tatcgatcaa 360
aagctggttt ggtcagattt agatactagg aatttttccc aaccactca agaacctgat 420
acaagtaatg ctgtaagtga gaaaaatctcc tcagatacca aagagaatag aaaagacct 480
gagactgaag atccttcaaa aaaaagtggc cttaaagaag tttcatcaga tctccctaaa 540
agtctgaaa ctgcagtagc agctatttct gaagatcttg aaatctcaga aaacatttca 600
gcaagagatc ctcttcaggg tttagcattt ttttataaaa atacatcttc tcagtctatc 660
tctgaaaagg atcttctcatt tcaaggaaatt atcttttctg gttcaggagc taattcaggg 720
ctagggtttt aaaatcttaa ggccgcgaaa tctggggctg cagtttattc tgatcgagat 780
attgtttttg aaaatcttgt taaaggattg agttttatat cttgtgaatc tttagaagat 840
ggctctgccc caggtgtaaa cattgtttgtg acccattgtg gtgatgtaac tctcactgat 900
tgtgacctg gtttagacct tgaagcttta cgtctggtta aagatttttc tctgtggagga 960
gctgttttca ctgctcgcaa ccatgaagtg caaaataacc ttgcagtggt aattctatcc 1020
gttgtaggca ataaaggagc tattgttgta gagaaaaata gtgctgagaa gtccaatgga 1080
ggagcttttt cttgcggaag tttgttttac agtaacaacg aaaacaccgc cttgtggaaa 1140
gaaaatcaag cattatcagg aggagccata tctcagcaa gtgatattga tattcaaggg 1200
aactgtagcg ctattgaatt ttcaggaaac cagtctctaa ttgctcttgg agagcatata 1260
gggctttacg attttgtagg tggaggagct ttagctgctc aagggaacct taccttaaga 1320
aataatgcag tagtgcaatg tgttaaaaaa acttctaaaa cacatggtgg agctatttta 1380
gcaggtagtg ttgatctcaa cgaaacaatt agcgaagttg cctttaagca gaatacagca 1440
gctctaactg gaggtgcttt aagtgc aaat gataaggtta taattgcaaa taacttttga 1500
gaaattcttt ttgagcaaaa cgaagtggag aatcacggag gagccattta ttgtggatgt 1560
cgatctaata ctaagttaga acaaaaggat tctggagaga acatcaatat tattggaaac 1620
tccggagcta tcaacttttt aaaaaataag gcttctgttt tagaagtgat gacacaagct 1680
gaagattatg ctgggtggagg cgctttatgg gggcataatg ttcttctaga ttccaatagt 1740
gggaatatct aatttatagg aaatataggt ggaagtacct totggatagg agaatatgtc 1800
gggtgtgttg cgattctctc tactgataga gtgacaattt ctaataaact tggagatgtt 1860
gttttttaag gaaacaaagg ccaatgtctt gctcaaaaat atgtagctoc tcaagaaaca 1920
gctcccgtag aatcagatgc ttcatctaca aataaagacg agaagagcct taatgcttgt 1980
agtcattggg atcattatcc tccataaact gtagaagagg aagtgccacc ttcatgttta 2040
gaagaacatc ctggtgtttc ttcgacagat attcgtgttg gtggggccat tctagctcaa 2100
catatcttta ttacagataa tacaggaaat ctgagattct ctgggaacct tgggtgtgtg 2160
gaagagtctt ctactgtcgg tgatttagct atcgtaggag gaggtgcttt gacttcaaat 2220
aatgaagtta atgtttgcag taaccaaaat gttgtttttt ctgataacgt gacttcaaat 2280
ggttgtgatt cagggggagc tatttttagct aaaaaagtag atatctccgc gaaccactcg 2340
gttgaatttt tctctaattg ttcagggaaa ttcgggtgtg ccgtttgcgc tttaaacgaa 2400
tcagtaaaaca ttacggacaa tggctoggca gtatcattct ctaaaaatag aacacgtctt 2460
ggcgggtgctg gagttgcagc tctcaaggc tctgtaacga tttgtggaaa tcagggaaaac 2520
atagcattta aagagaactt tgtttttggc totgaaaatc aaagatcagg tggaggagct 2580
atcattgcta actctctctg aaatattcag gataacgcag gagatctct atttgtaagt 2640
aactctacgg gatcttatgg aggtgtatatt tttgtaggat ctttggttgc ttctgaaggc 2700
agcaaccacg gaacgcttac aattacaggc aacagtgggg atatcctatt tgctaaaaat 2760
agcacgcaaa cagccgcttc tttatcagaa aaagattcct ttggtggagg ggccatctat 2820
acacaaaacc tcaaaattgt aaagaatgca gggaaacgtt ctttctatgg caacagagct 2880
cctagtgttg ctggtgtcca aattgcagac ggaggaactg tttgtttaga ggcttttga 2940
ggagatatct ttttgaagg gaatatcaat tttgatggga gtttcaatgc gattcactta 3000
tgccgggaatg actcaaaaat cgtagagctt tctgctgttc aagataaaaa tattattttc 3060
caagatgcaa ttacttatga agagaacaca attcgtggct tgccagataa agatgtcagt 3120
cctttaagtg ccccttcatt aatttttaac tccaagccac aagatgacag cgctcaacat 3180
catgaaggga cgtacgggtt ttctcgaggg gtatctaaaa ttcctcagat tgcgtctata 3240
caagagggaa ccttagcttt atcaca aaac gtagagcttt ggttggcagg acttaaacag 3300
gaaacaggaa gttctatcgt attgtctgcg ggatctatct tccgtatttt tgattcccag 3360
gttgatagca gtgcgcctct tctacagaa aataaagagg agactcttgt ttctgccgga 3420
gttcaaatta acatgagctc tctacaccc aataaagata aagctgtaga tactccagta 3480

```



```

cttgacagata tcataagtat tactgttagat ttgtcttcat ttgttcctga gcaagacgga 3540
actcttctctc ttctctcctga aattatcatt cctaagggaa caaaattaca ttctaagcc 3600
atagatctta agattataga tcctaccaat gtgggatatg aaaatcatgc tcttctaagt 3660
tctcataaag atattccatt aatttctctt aagacagcgg aaggaatgac agggacgcct 3720
acagcagatg cttctctatc taatataaaa atagatgtat ctttaccttc gatcacacca 3780
gcaacgtatg gtcacacagg agtttgggtct gaaagtaaaa tggaagatgg aagacttgta 3840
gtcgggttggc aacctacggg atataagtta aatcctgaga agcaaggggc tctagttttg 3900
aataatctct ggagtcatta tacagatctt agagctctta agcaggagat ctttgctcat 3960
catacgatag ctcaaagaat ggagtttagat ttctcgacaa atgtctgggg atcaggatta 4020
gggtgttggg aagattgtca gaacatcgga gagtttgatg ggttcaaaca tcatctcaca 4080
gggtatgccc taggcttgga tacacaacta gttgaagact tcttaattgg aggatgtttc 4140
tcacagttct ttggtaaaac tgaaagccaa tcctacaaag ctaagaacga tgtgaagagt 4200
tatatgggag ctgcttatgc ggggatttta gcaggtcctt ggttaataaa aggagctttt 4260
gtttacggta atataaacia cgatttgact acagattacg gtactttagg tatttcaaca 4320
ggttcatgga taggaaaagg gtttatcgca ggcacaagca ttgattaccg ctatattgta 4380
aatcctcgac ggtttatata ggcaatcgta tccacagtgg ttccttttgt agaagccgag 4440
tatgtccgta tagatcttcc agaaattagc gaacagggtg aagaggttag aacgttccaa 4500
aaaactcgtt ttgagaatgt cgcatttcct tttggatttg ctttagaaca tgcattatcg 4560
cgtggctcac gtgtgaagt gaacagtgtg cagcttgctt acgtctttga tgtatatcgt 4620
aagggacctg tctctttgat tacactcaag gatcttgctt attcttgga gagttatggg 4680
gtagatatct cttgtaaaagc ttggaaggct cgcttgagca ataatacgga atggaattca 4740
tatttaagta cgtatttagc gtttaattat gaatggagag aagatctgat agcttatgac 4800
ttcaatggtg gtatccgat tattttctag 4830

```

<210> 60

<211> 591

<212> DNA

<213> *Chlamydia pneumoniae*

<400> 60

```

atgacactct ccctagttgg aaaggaagcc cctgattttg ttgcgcaagc tgttggttaat 60
ggcgaaaagt gtaccgtatc tttaaaagat ttttaggaa agtatgttgt gcttttcttc 120
tatcctaagc attttactta cgtgtgtcct aocgaattgc acgcatttca agatgcttta 180
ggagaattcc acaccogagg agctgaagtc ataggctgtt ccgtggatga cattgccacc 240
catcaacagt ggttagctac taagaaaaag caaggtggta tcgaagggtat tacctatcct 300
cttctctcag acgaagataa agtcatttca agaagttatc atgtgttaaa acccgaagaa 360
gaattatctt tcagaggagt tttoctgatt gataaagggtg gaatcatccg tcatcttgta 420
gtgaatgatc ttctcttagg ccgttctata gaagaagaac ttagaaccct agatgcttta 480
atcttctttg aaactaatgg cttagtctgt cctgcaaatt ggcatgaagg agagcgagcg 540
atggctccaa atgaagaagg actgcaaaat tatttcggga ctatagacta g 591

```

<210> 61

<211> 1983

<212> DNA

<213> *Chlamydia pneumoniae*

<400> 61

```

atgagtgaac acaaaaaatc aagcaaaatt ataggtatag acttaggcac aacaaaactcc 60
tgcgatatctg ttatggaagg aggacaagct aaagtaatta catcatccga aggaacaaga 120
accacgccat cgatcggtgc cttcaaaggc aatgagaaat tagtggggat tccagcaaaa 180
cgtcaagcag tgacaaatcc agaaaaaact ctcggctcta caaaacgctt tattggccgt 240
aagtactctg aagtagcttc ggaaatccaa accgttcctt atacagtcac ctccggatct 300
aaagggtgatg ccgttttcga agttgatggc aaacaatata ctccagaaga aattggcgca 360
caaatcttaa tgaaatgaa agagacagca gaagcttata taggcgaaac tgtcacagaa 420
gcagtgatca ccgtccccgc atacttcaat gattctcaac gagcatccac aaaagatgct 480
ggacgcattg caggtctaga tgtaaaacgt atcattccag aacctaccgc agcagctctt 540
gcctacggaa tcgataaagt cgggtgataaa aaaatcgctg tcttcgacct tgggtggagga 600
acttttgata tctccatcct agaaatcggt gatggcgtct tcgaagttct atctacaaat 660
ggagatactc tcctcggtgg agacgacttt gatgaagtca ttatcaaatt gatgatcgaa 720

```

```

gaattcaaaa aacaagaagg cattgatctt agcaaagata atatggcctt acaaagactt 780
aaagatgctg ctgagaaagc aaaaatagaa ctttcaggag tctcttccac agaaatcaat 840
cagccattca tcacaatgga tgcacaagga cctaaacacc ttgcattgac actcacacgt 900
gcgcaattcg agaaactcgc agcctctcta atcgaaagaa caaaatctcc atgcatcaaa 960
gcactcagtg acgcaaaact ttccgctaag gatatcgatg atgttctctt agttggaggt 1020
atgtcaagaa tgcccgcagt gcaagaaact gtaaaagaac tcttcggcaa agagcctaatt 1080
aaaggagtca accccgacga agttgttgct attggagccg caattcaagg tgggtgttctt 1140
ggcggagaaag ttaaggatgt tctacttcta gacgttatcc ccctatctct gggtatcgaa 1200
actctaggag gcgtcatgac gactctggta gagagaaata ctacaatccc tacacagaaa 1260
aaacaaatct tctccacagc tgctgataac cagcctgcgg ttaccatcgt agttctccaa 1320
ggagagcgtc ccatggccaa agataacaag gaaatcggaa gattcgatct tacagataatc 1380
cctccggctc ctcgaggcca tcctcaaate gaagtctcct tcgatatoga tgcaaacgga 1440
atthttccatg tctcagctaa agatgttgcc agcggtaaaag aacagaaaat tcgtatcgaa 1500
gcaagctcag gacttcaaga agatgaaatc caaagaatgg ttcgagatgc cgaaattaat 1560
aaggaagaag ataaaaaacg tctgtaagct tcagatgcta aaaatgaagc cgatagcatg 1620
atcttcagag ccgaaaaagc tattaagat tataaggagc aaattcctga aactttagtt 1680
aaagaaatcg aagagcgaat cgaaaacgtg cgcaacgcac tcaaagatga cgctcctatt 1740
gaaaaaattt aagaggttac tgaagacctt agcaagcata tgcaaaaaat tggagagttc 1800
atgcaatcgc agtctgcac agcagcagca tcctcggcag ccaatgctaa aggtggacct 1860
aacatcaata cagaagattt gaaaaaacat agtttcagta cgaagcctcc ttcaaaatac 1920
ggttcttcag aagaccatat cgaagaagct gatgtagaaa ttattgataa cgacgataag 1980
taa
1983

```

<210> 62

<211> 1860

<212> DNA

<213> *Chlamydia pneumoniae*

<400> 62

```

atgaaaaaag ggaaattag agccatagtt tttggccttc tatttacaag tagtgttgct 60
ggtttttcta aggatattgac taaagacaac gcttatcaag attttaaagt catagagcat 120
ttaatatcgt taaaatatgc tcctttacca tggaaaggaa tattatttgg ttgggattta 180
tctcagcaaa cacagcaagc tcgcttgcaa ctgggtcttag aagaaaaacc aacaaccaac 240
tactgccaga aggtactctc taactacgtg agatcattaa acgattatca tgcaggaggt 300
acgttttata gtactgaaag tgcgtatatc ccttacgtat tgaagttaa tgagatggt 360
catgtctttg tagtcgacgt acagactagc caaggggata tttacttagg ggatgaaatc 420
cttgaagtag atggaaatgg gattcgtgag gctatcgaaa gccttcgctt tggacgaggg 480
agtgcacag actattctgc tgcagttcgt tccttgacat cgcgttcgcg cgcttttggg 540
gatgcggttc cttaaggaaat tgccatgttg aaacttcgcc gaccagtggt tttgatccgt 600
tcgacacccg tccgttgccg ttatactcca gagcatatcg gagatttttc tttagttgct 660
cctttgatic ctgaacataa acctcaatta cctacacaaa gttgtgtgct attccgttcc 720
ggggtaaatt cacagttctc tagtagctct ttattcagtt cctacatggt gccttatttc 780
tgggaagaat tgcgggttca aaataagcag cgttttgaca gtaatcacca tataggagc 840
cgtaatggat ttttacctac gtttggtcct attccttggg aacaagacaa ggggccctat 900
cgttcctata tctttaaagc aaaagattct cagggcaatc cccatcgcat aggattttta 960
agaatttctt cttatgtttg gactgattta gaaggacttg aagaggatca taaggatagt 1020
ccttgggagc tctttggaga gatcatcgat catttggaag aagagactga tgccttgatt 1080
attgatcaga ccataatcc tggaggcagt gttttctatc tctattcgtt actatctatg 1140
ttaacagatc atcctttaga tactcctaaa catagaatga ttttcaacta ggatgaagtc 1200
agctcggctt tgcaactggca agatctacta gaagatgtct tcacagatga gcaggcagtt 1260
gccgtgctag ggaaactat ggaaggata tgcattggata tgcatgtgt agcctctctt 1320
caaaacttct ctcagagtggt cctttcttcc tgggtttcag gtgatattaa cctttcaaaa 1380
cctatgcctt tgctaggatt tgcacaggtt cgacctcatc ctaaaccatca atatactaaa 1440
cctttgttta tgttgataga cgaggatgac ttctcttgtg gagatttagc gcctgcaatt 1500
ttgaaggata atggcgcgc tactctcatt ggaaagccaa cagcaggagc tggaggtttt 1560
gtattccaag tcaactttcc taaccgttct ggaattaaag gtctttcttt aacaggatct 1620
ttagctgtta ggaaagatgg tgagtttatt gaaaacttag gagtggctcc tcatattgat 1680
ttaggattta cctccaggga tttgcaaaact tccaggttta ctgattacgt tgaggcagtg 1740
aaaactatag ttttaacttc tttgtctgag aacgctaaga agagtgaaga gcagacttct 1800

```

ccgcaagaga cgcctgaagt tattcgagtc tcttatccca caacgacttc tgcttcgtaa 1860

<210> 63

<211> 1956

<212> DNA

<213> *Chlamydia pneumoniae*

<400> 63

```

atggttaatc ctattggtcc aggtcctata gacgaaacag aacgcacacc tcccgcagat 60
ctttctgctc aaggattgga ggcgagtgca gcaaataaga gtgcggaagc tcaaagaata 120
gcaggtgcgg aagctaagcc taaagaatct aagaccgatt ctgtagagcg atggagcacc 180
ttgcgttctg cagtgaatgc tctcatgagt ctggcagata agctgggtat tgcttctagt 240
aacagctcgt ctctactag cagatctgca gaogtggact caacgacagc gaccgcacct 300
acgcctcctc caccacggtt tgatgattat aagactcaag cgaaaacagc ttacgatact 360
atctttacct caacatcact agctgacata caggctgctt tgggtagcct ccaggatgct 420
gtcactaata taaaggatac agcggctact gatgaggaaa ccgcaatcgc tgcggagtgg 480
gaaactaaga atgcccagtc agttaaagtt ggcgcgcaaa ttacagaatt agcgaatat 540
gcttcggata accaagcgat tcttgactct ttaggtaaac tgacttcctt cgacctctta 600
caggctgctc ttctccaate tgtagcaaac aataacaaag cagctgagct tcttaaagag 660
atgcaagata acccagtagt cccagggaaa acgcctgcaa ttgctcaatc tttagttgat 720
cagacagatg ctacagcgac acagatagag aaagatggaa atgcgattag gtagtcatat 780
tttgccaggac agaaccgctag tggagctgta gaaaatgcta aatctaataa cagtataagc 840
aacatagatt cagctaaagc agcaatcgc actgctaaga cacaatatgc tgaagctcag 900
aaaaagttcc ccgactctcc aattcttcaa gaagcggaac aaatggtaat acaggctgag 960
aaagatctta aaaatatcaa acctgcagat ggttctgatg ttccaaatcc aggaactaca 1020
gttggaggct ccaagcaaca aggaagtagt attggtagta ttctgttttc catgctgtta 1080
gatgatgctg aaaaatgagc cgcttccatt ttgatgtctg ggtttcgtca gatgattcac 1140
atgttcaata cggaaaatcc tgattctcaa gctgcccac aggagctcgc agcacaagct 1200
agagcagcga aagccgctgg agatgacagt gctgctgcag cgctggcaga tgctcagaaa 1260
gcttttagaag cggctctagg taaagctggg caacaacagg gcatactcaa tgcttttaga 1320
cagatcgctt ctgctgctgt tgtgagcgca ggagttcttc ccgctgcagc aagttctata 1380
gggtcatctg taaaacagct ttacaagacc tcaaatcta caggttctga ttataaaaa 1440
cagatatcag caggttatga tgcttacaaa tccatcaatg atgcctatgg tagggcacga 1500
aatgatgcga ctctgatgt gataaacaat gtaagtaccc ccgctctcac acgatccgtt 1560
octagagcac gaacagaagc tcgaggacca gaaaaaacag atcaagccct cgctaggggtg 1620
atctctggca atagcagaac tcttgagat gtctatagtc aagtttcggc actacaatct 1680
gtaattcaga tcatccagtc gaatcctcaa gcgaataatg aggagatcag acaaaagctt 1740
acatcggcag tgacaaagcc tccacagttt ggctatcctt atgtgcaact ttctaatac 1800
totacacaga agttcatagc taaattagaa agtttggttg ctgaaggatc taggacagca 1860
gctgaaataa aagcactttc ctttgaaaag aactccttgt ttattcagca ggtgctggtc 1920
aatatcggct ctctatatcc tggttatctc caataa 1956

```

<210> 64

<211> 264

<212> DNA

<213> *Chlamydia pneumoniae*

<400> 64

```

atgagtcaaa aaaataaaaa ctctgctttt atgcatcccg tgaatatctc cacagattta 60
gcagttatag ttggcaaggg acctatgcc agaacggaaa ttgtaagaa agtttgggaa 120
tacattaaaa aacacaactg tcaggatcaa aaaaaataac gtaatatcct tcccgatgcg 180
aatcttgcca aagtctttgg ctctagtgat cctatcgaca tgttccaaat gaccaaagcc 240
ctttccaaac atattgtaaa ataa 264

```

<210> 65

<211> 978

<212> PRT

<213> *Chlamydia pneumoniae*

Met	Pro	Leu	Ser	Phe	Lys	Ser	Ser	Ser	Phe	Cys	Leu	Leu	Ala	Cys	Leu
				5					10					15	
Cys	Ser	Ala	Ser	Cys	Ala	Phe	Ala	Glu	Thr	Arg	Leu	Gly	Gly	Asn	Phe
			20					25					30		
Val	Pro	Pro	Ile	Thr	Asn	Gln	Gly	Glu	Glu	Ile	Leu	Leu	Thr	Ser	Asp
		35					40					45			
Phe	Val	Cys	Ser	Asn	Phe	Leu	Gly	Ala	Ser	Phe	Ser	Ser	Ser	Phe	Ile
	50					55					60				
Asn	Ser	Ser	Ser	Asn	Leu	Ser	Leu	Leu	Gly	Lys	Gly	Leu	Ser	Leu	Thr
	65				70					75					80
Phe	Thr	Ser	Cys	Gln	Ala	Pro	Thr	Asn	Ser	Asn	Tyr	Ala	Leu	Leu	Ser
				85				90						95	
Ala	Ala	Glu	Thr	Leu	Thr	Phe	Lys	Asn	Phe	Ser	Ser	Ile	Asn	Phe	Thr
			100					105					110		
Gly	Asn	Gln	Ser	Thr	Gly	Leu	Gly	Gly	Leu	Ile	Tyr	Gly	Lys	Asp	Ile
		115					120					125			
Val	Phe	Gln	Ser	Ile	Lys	Asp	Leu	Ile	Phe	Thr	Thr	Asn	Arg	Val	Ala
	130					135					140				
Tyr	Ser	Pro	Ala	Ser	Val	Thr	Thr	Ser	Ala	Thr	Pro	Ala	Ile	Thr	Thr
145					150					155					160
Val	Thr	Thr	Gly	Ala	Ser	Ala	Leu	Gln	Pro	Thr	Asp	Ser	Leu	Thr	Val
				165					170					175	
Glu	Asn	Ile	Ser	Gln	Ser	Ile	Lys	Phe	Phe	Gly	Asn	Leu	Ala	Asn	Phe
			180					185					190		
Gly	Ser	Ala	Ile	Ser	Ser	Ser	Pro	Thr	Ala	Val	Val	Lys	Phe	Ile	Asn
		195					200					205			
Asn	Thr	Ala	Thr	Met	Ser	Phe	Ser	His	Asn	Phe	Thr	Ser	Ser	Gly	Gly
	210					215					220				
Gly	Val	Ile	Tyr	Gly	Gly	Ser	Ser	Leu	Leu	Phe	Glu	Asn	Asn	Ser	Gly
225					230					235					240
Cys	Ile	Ile	Phe	Thr	Ala	Asn	Ser	Cys	Val	Asn	Ser	Leu	Lys	Gly	Val
				245					250					255	
Thr	Pro	Ser	Ser	Gly	Thr	Tyr	Ala	Leu	Gly	Ser	Gly	Gly	Ala	Ile	Cys
			260					265					270		
Ile	Pro	Thr	Gly	Thr	Phe	Glu	Leu	Lys	Asn	Asn	Gln	Gly	Lys	Cys	Thr
		275					280					285			
Phe	Ser	Tyr	Asn	Gly	Thr	Pro	Asn	Asp	Ala	Gly	Ala	Ile	Tyr	Ala	Glu
	290					295					300				
Thr	Cys	Asn	Ile	Val	Gly	Asn	Gln	Gly	Ala	Leu	Leu	Leu	Asp	Ser	Asn
305					310					315					320
Thr	Ala	Ala	Arg	Asn	Gly	Gly	Ala	Ile	Cys	Ala	Lys	Val	Leu	Asn	Ile
				325					330					335	
Gln	Gly	Arg	Gly	Pro	Ile	Glu	Phe	Ser	Arg	Asn	Arg	Ala	Glu	Lys	Gly
			340					345					350		
Gly	Ala	Ile	Phe	Ile	Gly										

465					470					475				480
Lys	Ile	Thr	Asp	Asn	Ala	Val	Val	Asn	Val	Leu	Gly	Phe	Ala	Thr
				485					490					495
Gly	Ser	Gly	Gln	Leu	Thr	Leu	Gly	Ser	Gly	Gly	Thr	Leu	Gly	Leu
			500					505					510	
Thr	Pro	Thr	Gly	Ala	Pro	Ala	Ala	Val	Asp	Phe	Thr	Ile	Gly	Lys
		515					520					525		
Ala	Phe	Asp	Pro	Phe	Ser	Phe	Leu	Lys	Arg	Asp	Phe	Val	Ser	Ala
		530				535					540			
Val	Asn	Ala	Gly	Thr	Lys	Asn	Val	Thr	Leu	Thr	Gly	Ala	Leu	Val
545					550					555				560
Asp	Glu	His	Asp	Val	Thr	Asp	Leu	Tyr	Asp	Met	Val	Ser	Leu	Gln
				565					570					575
Pro	Val	Ala	Ile	Pro	Ile	Ala	Val	Phe	Lys	Gly	Ala	Thr	Val	Thr
			580					585					590	
Thr	Gly	Phe	Pro	Asp	Gly	Glu	Ile	Ala	Thr	Pro	Ser	His	Tyr	Gly
		595					600					605		
Gln	Gly	Lys	Trp	Ser	Tyr	Thr	Trp	Ser	Arg	Pro	Leu	Leu	Ile	Pro
		610				615						620		
Pro	Asp	Gly	Gly	Phe	Pro	Gly	Gly	Pro	Ser	Pro	Ser	Ala	Asn	Thr
625					630					635				640
Tyr	Ala	Val	Trp	Asn	Ser	Asp	Thr	Leu	Val	Arg	Ser	Thr	Tyr	Ile
				645					650					655
Asp	Pro	Glu	Arg	Tyr	Gly	Glu	Ile	Val	Ser	Asn	Ser	Leu	Trp	Ile
			660					665					670	
Phe	Leu	Gly	Asn	Gln	Ala	Phe	Ser	Asp	Ile	Leu	Gln	Asp	Val	Leu
		675				680						685		
Ile	Asp	His	Pro	Gly	Leu	Ser	Ile	Thr	Ala	Lys	Ala	Leu	Gly	Ala
	690					695					700			
Val	Glu	His	Thr	Pro	Arg	Gln	Gly	His	Glu	Gly	Phe	Ser	Gly	Arg
705					710					715				720
Gly	Gly	Tyr	Gln	Ala	Ala	Leu	Ser	Met	Asn	Tyr	Thr	Asp	His	Thr
				725					730					735
Leu	Gly	Leu	Ser	Phe	Gly	Gln	Leu	Tyr	Gly	Lys	Thr	Asn	Ala	Asn
			740					745					750	
Tyr	Asp	Ser	Arg	Cys	Ser	Glu	Gln	Met	Tyr	Leu	Leu	Ser	Phe	Phe
		755				760						765		
Gln	Phe	Pro	Ile	Val	Thr	Gln	Lys	Ser	Glu	Ala	Leu	Ile	Ser	Trp
		770				775						780		
Ala	Ala	Tyr	Gly	Tyr	Ser	Lys	Asn	His	Leu	Asn	Thr	Thr	Tyr	Leu
785					790					795				800
Pro	Asp	Lys	Ala	Pro	Lys	Ser	Gln	Gly	Gln	Trp	His	Asn	Asn	Ser
				805					810					815
Tyr	Val	Leu	Ile	Ser	Ala	Glu	His	Pro	Phe	Leu	Asn	Trp	Cys	Leu
			820					825					830	
Thr	Arg	Pro	Leu	Ala	Gln	Ala	Trp	Asp	Leu	Ser	Gly	Phe	Ile	Ser
		835				840						845		
Glu	Phe	Leu	Gly	Gly	Trp	Gln	Ser	Lys	Phe	Thr	Glu	Thr	Gly	Asp
		850				855					860			
Gln	Arg	Ser	Phe	Ser	Arg	Gly	Lys	Gly	Tyr	Asn	Val	Ser	Leu	Pro
865					870					875				880
Gly	Cys	Ser	Ser	Gln	Trp	Phe	Thr	Pro	Phe	Lys	Lys	Ala	Pro	Ser
				885					890					895
Leu	Thr	Ile	Lys	Leu	Ala	Tyr	Lys	Pro	Asp	Ile	Tyr	Arg	Val	Asn
			900					905					910	
His	Asn	Ile	Val	Thr	Val	Val	Ser	Asn	Gln	Glu	Ser	Thr	Ser	Ile
		915					920					925		
Gly	Ala	Asn	Leu	Arg	Arg	His	Gly	Leu	Phe	Val	Gln	Ile	His	Asp
		930				935					940			
Val	Asp	Leu	Thr	Glu	Asp	Thr	Gln	Ala	Phe	Leu	Asn	Tyr	Thr	Phe
945					950					955				960

Gly Lys Asn Gly Phe Thr Asn His Arg Val Ser Thr Gly Leu Lys Ser
 965 970 975
 Thr Phe

<210> 66
 <211> 266
 <212> PRT
 <213> Chlamydia pneumoniae

<400> 66
 Met His Ser Lys Phe Leu Ser Arg Arg Lys Lys Asn Ser Ser His Lys
 5 10 15
 Glu Glu Thr Ser Trp Asp Cys Ile Ala Ser Ser Tyr Asn Lys Ile Val
 20 25 30
 Gln Asp Lys Gly His Tyr Tyr His Arg Glu Thr Ile Leu Pro Gln Leu
 35 40 45
 Leu Pro Ser Leu Thr Leu Gly Ser Lys Ser Ser Val Leu Asp Ile Gly
 50 55 60
 Cys Gly Gln Gly Phe Leu Glu Arg Ala Leu Pro Lys Glu Cys Arg Tyr
 65 70 75 80
 Leu Gly Ile Asp Ile Ser Ser Arg Leu Ile Ala Leu Ala Lys Lys Met
 85 90 95
 Arg Ser Val Asn Ser His Gln Phe Lys Val Ala Asp Leu Ser Lys Arg
 100 105 110
 Leu Glu Phe Val Glu Pro Thr Leu Phe Ser His Ala Val Ala Ile Leu
 115 120 125
 Ser Leu Gln Asn Met Glu Phe Pro Gly Glu Ala Ile Arg Asn Thr Ala
 130 135 140
 Thr Leu Leu Glu Pro Leu Gly Gln Phe Phe Ile Val Leu Asn His Pro
 145 150 155 160
 Cys Phe Arg Ile Pro Arg Ala Ser Ser Trp His Tyr Asp Glu Asn Lys
 165 170 175
 Lys Ala Ile Ser Arg His Ile Asp Arg Tyr Leu Ser Pro Met Lys Ile
 180 185 190
 Pro Ile Met Ala His Pro Gly Gln Lys Asp Ser Pro Ser Thr Leu Ser
 195 200 205
 Phe His Phe Pro Leu Ser Tyr Trp Phe Lys Glu Leu Ser Ser His Gly
 210 215 220
 Phe Leu Val Ser Gly Leu Glu Glu Trp Thr Ser Ser Lys Thr Ser Thr
 225 230 235 240
 Gly Lys Arg Ala Lys Ala Glu Asn Leu Cys Arg Lys Glu Phe Pro Leu
 245 250 255
 Phe Leu Met Ile Ser Cys Ile Lys Ile Lys
 260 265

<210> 67
 <211> 83
 <212> PRT
 <213> Chlamydia pneumoniae

<400> 67
 Met Lys Gln Gln His Asn Arg Lys Ala Leu Ser Arg Lys Ile Gly Thr
 5 10 15
 Val Lys Lys Gln Ala Lys Phe Ala Gly Ser Phe Leu Asp Glu Ile Lys
 20 25 30
 Lys Ile Glu Trp Val Ser Lys His Asp Leu Lys Lys Tyr Ile Lys Val
 35 40 45
 Val Leu Ile Ser Ile Phe Gly Phe Gly Phe Ala Ile Tyr Phe Val Asp
 50 55 60

Leu Val Leu Arg Lys Ser Ile Thr Cys Leu Asp Gly Ile Thr Thr Phe
 65 70 75 80
 Leu Phe Gly

<210> 68

<211> 394

<212> PRT

<213> Chlamydia pneumoniae

<400> 68

Met Ser Lys Glu Thr Phe Gln Arg Asn Lys Pro His Ile Asn Ile Gly
 5 10 15
 Thr Ile Gly His Val Asp His Gly Lys Thr Thr Leu Thr Ala Ala Ile
 20 25 30
 Thr Arg Ala Leu Ser Gly Asp Gly Leu Ala Ser Phe Arg Asp Tyr Ser
 35 40 45
 Ser Ile Asp Asn Thr Pro Glu Glu Lys Ala Arg Gly Ile Thr Ile Asn
 50 55 60
 Ala Ser His Val Glu Tyr Glu Thr Pro Asn Arg His Tyr Ala His Val
 65 70 75 80
 Asp Cys Pro Gly His Ala Asp Tyr Val Lys Asn Met Ile Thr Gly Ala
 85 90 95
 Ala Gln Met Asp Gly Ala Ile Leu Val Val Ser Ala Thr Asp Gly Ala
 100 105 110
 Met Pro Gln Thr Lys Glu His Ile Leu Leu Ala Arg Gln Val Gly Val
 115 120 125
 Pro Tyr Ile Val Val Phe Leu Asn Lys Val Asp Met Ile Ser Gln Glu
 130 135 140
 Asp Ala Glu Leu Ile Asp Leu Val Glu Met Glu Leu Ser Glu Leu Leu
 145 150 155 160
 Glu Glu Lys Gly Tyr Lys Gly Cys Pro Ile Ile Arg Gly Ser Ala Leu
 165 170 175
 Lys Ala Leu Glu Gly Asp Ala Asn Tyr Ile Glu Lys Val Arg Glu Leu
 180 185 190
 Met Gln Ala Val Asp Asp Asn Ile Pro Thr Pro Glu Arg Glu Ile Asp
 195 200 205
 Lys Pro Phe Leu Met Pro Ile Glu Asp Val Phe Ser Ile Ser Gly Arg
 210 215 220
 Gly Thr Val Val Thr Gly Arg Ile Glu Arg Gly Ile Val Lys Val Ser
 225 230 235 240
 Asp Lys Val Gln Leu Val Gly Leu Gly Glu Thr Lys Glu Thr Ile Val
 245 250 255
 Thr Gly Val Glu Met Phe Arg Lys Glu Leu Pro Glu Gly Arg Ala Gly
 260 265 270
 Glu Asn Val Gly Leu Leu Leu Arg Gly Ile Gly Lys Asn Asp Val Glu
 275 280 285
 Arg Gly Met Val Val Cys Gln Pro Asn Ser Val Lys Pro His Thr Lys
 290 295 300
 Phe Lys Ser Ala Val Tyr Val Leu Gln Lys Glu Glu Gly Gly Arg His
 305 310 315 320
 Lys Pro Phe Phe Ser Gly Tyr Arg Pro Gln Phe Phe Phe Arg Thr Thr
 325 330 335
 Asp Val Thr Gly Val Val Thr Leu Pro Glu Gly Thr Glu Met Val Met
 340 345 350
 Pro Gly Asp Asn Val Glu Leu Asp Val Glu Leu Ile Gly Thr Val Ala
 355 360 365
 Leu Glu Glu Gly Met Arg Phe Ala Ile Arg Glu Gly Gly Arg Thr Ile
 370 375 380
 Gly Ala Gly Thr Ile Ser Lys Ile Asn Ala
 385 390

<210> 69

<211> 476

<212> PRT

<213> Chlamydia pneumoniae

<400> 69

```

Met Arg Ile Val Gln Val Ala Val Glu Phe Thr Pro Ile Val Lys Val
                    5              10              15
Gly Gly Leu Gly Asp Ala Val Ala Ser Leu Ser Lys Glu Leu Ala Lys
                    20              25              30
Gln Asn Asp Val Glu Val Leu Leu Pro His Tyr Pro Leu Ile Ser Lys
                    35              40              45
Phe Ser Ser Ser Gln Val Leu Ser Glu Arg Ser Phe Tyr Tyr Glu Phe
                    50              55              60
Leu Gly Lys Gln Gln Ala Ser Ala Ile Ser Tyr Ser Tyr Glu Gly Leu
                    65              70              75              80
Thr Leu Thr Ile Ile Thr Leu Asp Ser Gln Ile Glu Leu Phe Ser Thr
                    85              90              95
Thr Ser Val Tyr Ser Glu Asn Asn Val Val Arg Phe Ser Ala Phe Ala
                    100             105             110
Ala Ala Ala Ala Ala Tyr Leu Gln Glu Ala Asp Pro Ala Asp Ile Val
                    115             120             125
His Leu His Asp Trp His Val Gly Leu Leu Ala Gly Leu Leu Lys Asn
                    130             135             140
Pro Leu Asn Pro Val His Ser Lys Ile Val Phe Thr Ile His Asn Phe
                    145             150             155             160
Gly Tyr Arg Gly Tyr Cys Ser Thr Gln Leu Leu Ala Ala Ser Gln Ile
                    165             170             175
Asp Asp Phe His Leu Ser His Tyr Gln Leu Phe Arg Asp Pro Gln Thr
                    180             185             190
Ser Val Leu Met Lys Gly Ala Leu Tyr Cys Ser Asp Tyr Ile Thr Thr
                    195             200             205
Val Ser Leu Thr Tyr Val Gln Glu Ile Ile Asn Asp Tyr Ser Asp Tyr
                    210             215             220
Glu Leu His Asp Ala Ile Leu Ala Arg Asn Ser Val Phe Ser Gly Ile
                    225             230             235             240
Ile Asn Gly Ile Asp Glu Asp Val Trp Asn Pro Lys Thr Asp Pro Ala
                    245             250             255
Leu Ala Val Gln Tyr Asp Ala Ser Leu Leu Ser Glu Pro Asp Val Leu
                    260             265             270
Phe Thr Lys Lys Glu Glu Asn Arg Ala Val Leu Tyr Glu Lys Leu Gly
                    275             280             285
Ile Ser Ser Asp Tyr Phe Pro Leu Ile Cys Val Ile Ser Arg Ile Val
                    290             295             300
Glu Glu Lys Gly Pro Glu Phe Met Lys Glu Ile Ile Leu His Ala Met
                    305             310             315             320
Glu His Ser Tyr Ala Phe Ile Leu Ile Gly Thr Ser Gln Asn Glu Val
                    325             330             335
Leu Leu Asn Glu Phe Arg Asn Leu Gln Asp Cys Leu Ala Ser Ser Pro
                    340             345             350
Asn Ile Arg Leu Ile Leu Asp Phe Asn Asp Pro Leu Ala Arg Leu Thr
                    355             360             365
Tyr Ala Ala Ala Asp Met Ile Cys Ile Pro Ser His Arg Glu Ala Cys
                    370             375             380
Gly Leu Thr Gln Leu Ile Ala Met Arg Tyr Gly Thr Val Pro Leu Val
                    385             390             395             400
Arg Lys Thr Gly Gly Leu Ala Asp Thr Val Ile Pro Gly Val Asn Gly
                    405             410             415
Phe Thr Phe Phe Asp Thr Asn Asn Phe Asn Glu Phe Arg Ala Met Leu

```



```

          420          425          430
Ser Asn Ala Val Thr Thr Tyr Arg Gln Glu Pro Asp Val Trp Leu Asn
          435          440          445
Leu Ile Glu Ser Gly Met Leu Arg Ala Ser Gly Leu Asp Ala Met Ala
          450          455          460
Lys His Tyr Val Asn Leu Tyr Gln Ser Leu Leu Ser
465          470          475

```

<210> 70
 <211> 346
 <212> PRT
 <213> Chlamydia pneumoniae

```

<400> 70
Met Glu Ala Asp Ile Leu Asp Gly Lys Leu Lys Arg Val Glu Val Ser
          5          10          15
Lys Lys Gly Leu Val Asn Cys Asn Gln Val Asp Val Asn Gln Leu Val
          20          25          30
Pro Ile Lys Tyr Lys Trp Ala Trp Glu His Tyr Leu Asn Gly Cys Ala
          35          40          45
Asn Asn Trp Leu Pro Thr Glu Val Pro Met Ala Arg Asp Ile Glu Leu
          50          55          60
Trp Lys Ser Asp Glu Leu Ser Glu Asp Glu Arg Val Ile Leu Leu
          65          70          75          80
Asn Leu Gly Phe Phe Ser Thr Ala Glu Ser Leu Val Gly Asn Asn Ile
          85          90          95
Val Leu Ala Ile Phe Lys His Ile Thr Asn Pro Glu Ala Arg Gln Tyr
          100          105          110
Leu Leu Arg Gln Ala Phe Glu Glu Ala Val His Thr His Thr Phe Leu
          115          120          125
Tyr Ile Cys Glu Ser Leu Gly Leu Asp Glu Gly Glu Val Phe Asn Ala
          130          135          140
Tyr Asn Glu Arg Ala Ser Ile Arg Ala Lys Asp Asp Phe Gln Met Thr
145          150          155          160
Leu Thr Val Asp Val Leu Asp Pro Asn Phe Ser Val Gln Ser Ser Glu
          165          170          175
Gly Leu Gly Gln Phe Ile Lys Asn Leu Val Gly Tyr Tyr Ile Ile Met
          180          185          190
Glu Gly Ile Phe Phe Tyr Ser Gly Phe Val Met Ile Leu Ser Phe His
          195          200          205
Arg Gln Asn Lys Met Thr Gly Ile Gly Glu Gln Tyr Gln Tyr Ile Leu
210          215          220
Arg Asp Glu Thr Ile His Leu Asn Phe Gly Ile Asp Leu Ile Asn Gly
225          230          235          240
Ile Lys Glu Glu Asn Pro Glu Val Trp Thr Thr Glu Leu Gln Glu Glu
          245          250          255
Ile Val Ala Leu Ile Glu Lys Ala Val Glu Leu Glu Ile Glu Tyr Ala
          260          265          270
Lys Asp Cys Leu Pro Arg Gly Ile Leu Gly Leu Arg Ser Ser Met Phe
          275          280          285
Ile Asp Tyr Val Arg His Ile Ala Asp Arg Arg Leu Glu Arg Ile Gly
290          295          300
Leu Lys Pro Ile Tyr His Ser Arg Asn Pro Phe Pro Trp Met Ser Glu
305          310          315          320
Thr Met Asp Leu Asn Lys Glu Lys Asn Phe Phe Glu Thr Arg Val Thr
          325          330          335
Glu Tyr Gln Thr Ala Gly Asn Leu Ser Trp
          340          345

```

40

<210> 71
 <211> 1044
 <212> PRT
 <213> Chlamydia pneumoniae

<400> 71
 Met Val Glu Val Glu Glu Lys His Tyr Thr Ile Val Lys Arg Asn Gly
 5 10 15
 Met Phe Val Pro Phe Asn Gln Asp Arg Ile Phe Gln Ala Leu Glu Ala
 20 25 30
 Ala Phe Arg Asp Thr Arg Ser Leu Glu Thr Ser Ser Pro Leu Pro Lys
 35 40 45
 Asp Leu Glu Glu Ser Ile Ala Gln Ile Thr His Lys Val Val Lys Glu
 50 55 60
 Val Leu Ala Lys Ile Ser Glu Gly Gln Val Val Thr Val Glu Arg Ile
 65 70 75 80
 Gln Asp Leu Val Glu Ser Gln Leu Tyr Ile Ser Gly Leu Gln Asp Val
 85 90 95
 Ala Arg Asp Tyr Ile Val Tyr Arg Asp Gln Arg Lys Ala Glu Arg Gly
 100 105 110
 Asn Ser Ser Ser Ile Ile Ala Ile Ile Arg Arg Asp Gly Gly Ser Ala
 115 120 125
 Lys Phe Asn Pro Met Lys Ile Ser Ala Ala Leu Glu Lys Ala Phe Arg
 130 135 140
 Ala Thr Leu Gln Ile Asn Gly Met Thr Pro Pro Ala Thr Leu Ser Glu
 145 150 155 160
 Ile Asn Asp Leu Thr Leu Arg Ile Val Glu Asp Val Leu Ser Leu His
 165 170 175
 Gly Glu Glu Ala Ile Asn Leu Glu Glu Ile Gln Asp Ile Val Glu Lys
 180 185 190
 Gln Leu Met Val Ala Gly Tyr Tyr Asp Val Ala Lys Asn Tyr Ile Leu
 195 200 205
 Tyr Arg Glu Ala Arg Ala Arg Ala Arg Ala Asn Lys Asp Gln Asp Gly
 210 215 220
 Gln Glu Glu Phe Val Pro Gln Glu Glu Thr Tyr Val Val Gln Lys Glu
 225 230 235 240
 Asp Gly Thr Thr Tyr Leu Leu Arg Lys Thr Asp Leu Glu Lys Arg Phe
 245 250 255
 Ser Trp Ala Cys Lys Arg Phe Pro Lys Thr Thr Asp Ser Gln Leu Leu
 260 265 270
 Ala Asp Met Ala Phe Met Asn Leu Tyr Ser Gly Ile Lys Glu Asp Glu
 275 280 285
 Val Thr Thr Ala Cys Ile Met Ala Ala Arg Ala Asn Ile Glu Arg Glu
 290 295 300
 Pro Asp Tyr Ala Phe Ile Ala Ala Glu Leu Leu Thr Ser Ser Leu Tyr
 305 310 315 320
 Glu Glu Thr Leu Gly Cys Ser Ser Gln Asp Pro Asn Leu Ser Glu Ile
 325 330 335
 His Lys Lys His Phe Lys Glu Tyr Ile Leu Asn Gly Glu Glu Tyr Arg
 340 345 350
 Leu Asn Pro Gln Leu Lys Asp Tyr Asp Leu Asp Ala Leu Ser Glu Val
 355 360 365
 Leu Asp Leu Ser Arg Asp Gln Gln Phe Ser Tyr Met Gly Val Gln Asn
 370 375 380
 Leu Tyr Asp Arg Tyr Phe Asn Leu His Glu Gly Arg Arg Leu Glu Thr
 385 390 395 400
 Ala Gln Ile Phe Trp Met Arg Val Ser Met Gly Leu Ala Leu Asn Glu
 405 410 415
 Gly Glu Gln Lys Asn Phe Trp Ala Ile Thr Phe Tyr Asn Leu Leu Ser
 420 425 430
 Thr Phe Arg Tyr Thr Pro Ala Thr Pro Thr Leu Phe Asn Ser Gly Met

			435						440						445					
Arg	His	Ser	Gln	Leu	Ser	Ser	Cys	Tyr	Leu	Ser	Thr	Val	Lys	Asp	Asp					
	450					455					460									
Leu	Ser	His	Ile	Tyr	Lys	Val	Ile	Ser	Asp	Asn	Ala	Leu	Leu	Ser	Lys					
465					470					475					480					
Trp	Ala	Gly	Gly	Ile	Gly	Asn	Asp	Trp	Thr	Asp	Val	Arg	Ala	Thr	Gly					
				485					490					495						
Ala	Val	Ile	Lys	Gly	Thr	Asn	Gly	Lys	Ser	Gln	Gly	Val	Ile	Pro	Phe					
			500					505					510							
Ile	Lys	Val	Ala	Asn	Asp	Thr	Ala	Ile	Ala	Val	Asn	Gln	Gly	Gly	Lys					
		515					520					525								
Arg	Lys	Gly	Ala	Met	Cys	Val	Tyr	Leu	Glu	Asn	Trp	His	Leu	Asp	Tyr					
	530				535					540										
Glu	Asp	Phe	Leu	Glu	Leu	Arg	Lys	Asn	Thr	Gly	Asp	Glu	Arg	Arg	Arg					
545					550					555					560					
Thr	His	Asp	Ile	Asn	Thr	Ala	Ser	Trp	Ile	Pro	Asp	Leu	Phe	Phe	Lys					
				565					570					575						
Arg	Leu	Glu	Lys	Lys	Gly	Met	Trp	Thr	Leu	Phe	Ser	Pro	Asp	Asp	Val					
			580					585					590							
Pro	Gly	Leu	His	Glu	Ala	Tyr	Gly	Leu	Glu	Phe	Glu	Lys	Leu	Tyr	Glu					
	595						600					605								
Glu	Tyr	Glu	Arg	Lys	Val	Glu	Ser	Gly	Glu	Ile	Arg	Leu	Tyr	Lys	Lys					
	610					615					620									
Val	Glu	Ala	Glu	Val	Leu	Trp	Arg	Lys	Met	Leu	Ser	Met	Leu	Tyr	Glu					
625					630					635					640					
Thr	Gly	His	Pro	Trp	Ile	Thr	Phe	Lys	Asp	Pro	Ser	Asn	Ile	Arg	Ser					
				645					650					655						
Asn	Gln	Asp	His	Val	Gly	Val	Val	Arg	Cys	Ser	Asn	Leu	Cys	Thr	Glu					
			660					665					670							
Ile	Leu	Leu	Asn	Cys	Ser	Glu	Ser	Glu	Thr	Ala	Val	Cys	Asn	Leu	Gly					
		675					680					685								
Ser	Ile	Asn	Leu	Val	Glu	His	Ile	Arg	Asn	Asp	Lys	Leu	Asp	Glu	Glu					
	690					695					700									
Lys	Leu	Lys	Glu	Thr	Ile	Ser	Ile	Ala	Ile	Arg	Ile	Leu	Asp	Asn	Val					
705					710					715					720					
Ile	Asp	Leu	Asn	Phe	Tyr	Pro	Thr	Pro	Glu	Ala	Lys	Gln	Ala	Asn	Leu					
				725					730					735						
Thr	His	Arg	Ala	Val	Gly	Leu	Gly	Val	Met	Gly	Phe	Gln	Asp	Val	Leu					
			740					745					750							
Tyr	Glu	Leu	Asn	Ile	Ser	Tyr	Ala	Ser	Gln	Glu	Ala	Val	Glu	Phe	Ser					
	755						760					765								
Asp	Glu	Cys	Ser	Glu	Ile	Ile	Ala	Tyr	Tyr	Ala	Ile	Leu	Ala	Ser	Ser					
	770					775					780	</								

Lys Lys Leu Phe Leu Thr Ala Phe Glu Ile Glu Pro Glu Trp Ile Ile
 930 935 940
 Glu Cys Thr Ser Arg Arg Gln Lys Trp Ile Asp Met Gly Val Ser Leu
 945 950 955 960
 Asn Leu Tyr Leu Ala Glu Pro Asp Gly Lys Lys Leu Ser Asn Met Tyr
 965 970 975
 Leu Thr Ala Trp Lys Lys Gly Leu Lys Thr Thr Tyr Tyr Leu Arg Ser
 980 985 990
 Gln Ala Ala Thr Ser Val Glu Lys Ser Phe Ile Asp Ile Asn Lys Arg
 995 1000 1005
 Gly Ile Gln Pro Arg Trp Met Lys Asn Lys Ser Ala Ser Thr Ser Ile
 1010 1015 1020
 Val Val Glu Arg Lys Thr Thr Pro Val Cys Ser Met Glu Glu Gly Cys
 1025 1030 1035 1040
 Glu Ser Cys Gln

<210> 72

<211> 461

<212> PRT

<213> Chlamydia pneumoniae

<400> 72

Met Met Ser Ser Lys Arg Thr Ser Lys Ile Ala Val Leu Ser Ile Leu
 5 10 15
 Leu Thr Phe Thr His Ser Ile Gly Phe Ala Asn Ala Asn Ser Ser Val
 20 25 30
 Gly Leu Gly Thr Val Tyr Ile Thr Ser Glu Val Val Lys Lys Pro Gln
 35 40 45
 Lys Gly Ser Glu Arg Lys Gln Ala Lys Lys Glu Pro Arg Ala Arg Lys
 50 55 60
 Gly Tyr Leu Val Pro Ser Ser Arg Thr Leu Ser Ala Arg Ala Gln Lys
 65 70 75 80
 Met Lys Asn Ser Ser Arg Lys Glu Ser Ser Gly Gly Cys Asn Glu Ile
 85 90 95
 Ser Ala Asn Ser Thr Pro Arg Ser Val Lys Leu Arg Arg Asn Lys Arg
 100 105 110
 Ala Glu Gln Lys Ala Ala Lys Gln Gly Phe Ser Ala Phe Ser Asn Leu
 115 120 125
 Thr Leu Lys Ser Leu Leu Pro Lys Leu Pro Ser Lys Gln Lys Thr Ser
 130 135 140
 Ile His Glu Arg Glu Lys Ala Thr Ser Arg Phe Val Asn Glu Ser Gln
 145 150 155 160
 Leu Ser Ser Ala Arg Lys Arg Tyr Cys Thr Pro Ser Ser Ala Ala Pro
 165 170 175
 Ser Leu Phe Leu Glu Thr Glu Ile Val Arg Ala Pro Val Glu Arg Thr
 180 185 190
 Lys Glu Leu Gln Asp Asn Glu Ile His Ile Pro Val Val Gln Val Gln
 195 200 205
 Thr Asn Pro Lys Glu Gln Asn Thr Lys Thr Thr Lys Gln Leu Ala Ser
 210 215 220
 Gln Ala Ser Ile Gln Gln Ser Glu Gly Thr Glu Gln Ser Leu Arg Glu
 225 230 235 240
 Leu Ala Gln Gly Ala Ser Leu Pro Val Leu Val Arg Ser Asn Pro Glu
 245 250 255
 Val Ser Val Gln Arg Gln Lys Glu Glu Leu Lys Glu Leu Val Ala
 260 265 270
 Glu Arg Arg Gln Cys Lys Arg Lys Ser Val Arg Gln Ala Leu Glu Ala
 275 280 285
 Arg Ser Leu Thr Lys Lys Val Ala Arg Gly Gly Ser Val Thr Ser Thr

```

      290              295              300
Leu Arg Tyr Asp Pro Glu Lys Ala Ala Glu Ile Lys Ser Arg Arg Asn
305              310              315              320
Cys Lys Val Ser Pro Glu Ala Arg Glu Gln Lys Tyr Ser Ser Cys Lys
      325              330              335
Arg Asp Ala Arg Ala Asn Gly Lys Gln Asp Lys Thr Thr Pro Ser Glu
      340              345              350
Asp Ala Ser Gln Glu Glu Gln Gln Thr Gly Ala Gly Leu Val Arg Lys
      355              360              365
Thr Pro Lys Ser Gln Val Ala Ser Asn Ala Gln Asn Phe Tyr Arg Asn
      370              375              380
Ser Lys Asn Thr Asn Ile Asp Ser Tyr Leu Thr Ala Asn Gln Tyr Ser
385              390              395              400
Cys Ser Ser Glu Glu Thr Asp Trp Pro Cys Ser Ser Cys Val Ser Lys
      405              410              415
Arg Arg Thr His Asn Ser Ile Ser Val Cys Thr Met Val Val Thr Val
      420              425              430
Ile Ala Met Ile Val Gly Ala Leu Ile Ile Ala Asn Ala Thr Glu Ser
      435              440              445
Gln Thr Thr Ser Asp Pro Thr Pro Pro Thr Pro Thr Pro
450              455              460

```

<210> 73
 <211> 576
 <212> PRT
 <213> Chlamydia pneumoniae

```

<400> 73
Met Thr Asp Phe Pro Thr His Phe Lys Gly Pro Lys Leu Asn Pro Ile
      5              10              15
Lys Val Asn Pro Asn Phe Phe Glu Arg Asn Pro Lys Val Ala Arg Val
      20              25              30
Leu Gln Ile Thr Ala Val Val Leu Gly Ile Ile Ala Leu Leu Ser Gly
      35              40              45
Ile Val Leu Ile Ile Gly Thr Pro Leu Gly Ala Pro Ile Ser Met Ile
      50              55              60
Leu Gly Gly Cys Leu Leu Ala Ser Gly Gly Ala Leu Phe Val Gly Gly
      65              70              75              80
Thr Ile Ala Thr Ile Leu Gln Ala Arg Asn Ser Tyr Lys Lys Ala Val
      85              90              95
Asn Gln Lys Lys Leu Ser Glu Pro Leu Met Glu Arg Pro Glu Leu Lys
      100              105              110
Ala Leu Asp Tyr Ser Leu Asp Leu Lys Glu Val Trp Asp Leu His His
      115              120              125
Ser Val Val Lys His Leu Lys Lys Leu Asp Leu Asn Leu Ser Lys Thr
      130              135              140
Gln Arg Glu Val Leu Asn Gln Ile Lys Ile Asp Asp Glu Gly Pro Ser
145              150              155              160
Leu Gly Glu Cys Ala Ala Met Ile Ser Glu Asn Tyr Asp Ala Cys Leu
      165              170              175
Lys Met Leu Ala Tyr Arg Glu Glu Leu Leu Lys Glu Gln Thr Gln Tyr
      180              185              190
Gln Glu Thr Arg Phe Asn Gln Asn Leu Thr His Arg Asn Lys Val Leu
      195              200              205
Leu Ser Ile Leu Ser Arg Ile Thr Asp Asn Ile Ser Lys Ala Gly Gly
      210              215              220
Val Phe Ser Leu Lys Phe Ser Thr Leu Ser Ser Arg Met Ser Arg Ile
225              230              235              240
His Thr Thr Thr Thr Val Ile Leu Ala Leu Ser Ala Val Val Ser Val
      245              250              255

```

```

Met Val Val Ala Ala Leu Ile Pro Gly Gly Ile Leu Ala Leu Pro Ile
      260      265      270
Leu Leu Ala Val Ala Ile Ser Ala Gly Val Ile Val Thr Gly Leu Ser
      275      280      285
Tyr Leu Val Arg Gln Ile Leu Ser Asn Thr Lys Arg Asn Arg Gln Asp
      290      295      300
Phe Tyr Lys Asp Phe Val Lys Asn Val Asp Ile Glu Leu Leu Asn Gln
      305      310      315      320
Thr Val Thr Leu Gln Arg Phe Leu Phe Glu Met Leu Lys Gly Val Leu
      325      330      335
Lys Glu Glu Glu Glu Val Ser Leu Glu Gly Gln Asp Trp Tyr Thr Gln
      340      345      350
Tyr Ile Thr Asn Ala Pro Ile Glu Lys Arg Leu Ile Glu Glu Ile Arg
      355      360      365
Val Thr Tyr Lys Glu Ile Asp Ala Gln Thr Lys Lys Met Lys Thr Asp
      370      375      380
Leu Glu Phe Leu Glu Asn Glu Val Arg Ser Gly Arg Leu Ser Val Ala
      385      390      395      400
Ser Pro Ser Glu Asp Pro Ser Glu Thr Pro Ile Phe Thr Gln Gly Lys
      405      410      415
Glu Phe Ala Lys Leu Arg Arg Gln Thr Ser Gln Asn Ile Ser Thr Ile
      420      425      430
Tyr Gly Pro Asp Asn Glu Asn Ile Asp Pro Glu Phe Ser Leu Pro Trp
      435      440      445
Met Pro Lys Lys Glu Glu Glu Ile Asp His Ser Leu Glu Pro Val Thr
      450      455      460
Lys Leu Glu Pro Gly Ser Arg Glu Glu Leu Leu Leu Val Glu Gly Val
      465      470      475      480
Asn Pro Thr Leu Arg Glu Leu Asn Met Arg Ile Ala Leu Leu Gln Gln
      485      490      495
Gln Leu Ser Ser Val Arg Lys Trp Arg His Pro Arg Gly Glu His Tyr
      500      505      510
Gly Asn Val Ile Tyr Ser Asp Thr Glu Leu Asp Arg Ile Gln Met Leu
      515      520      525
Glu Gly Ala Phe Tyr Asn His Leu Arg Glu Ala Gln Glu Glu Ile Thr
      530      535      540
Gln Ser Leu Gly Asp Leu Val Asp Ile Gln Asn Arg Ile Leu Gly Ile
      545      550      555      560
Ile Val Glu Gly Asp Ser Asp Ser Arg Thr Glu Glu Glu Pro Gln Glu
      565      570      575

```

<210> 74

<211> 361

<212> PRT

<213> Chlamydia pneumoniae

<400> 74

```

Met Gln Gln Thr Val Ile Val Ala Met Ser Gly Gly Val Asp Ser Ser
      5      10      15
Val Val Ala Tyr Leu Phe Lys Lys Phe Thr Asn Tyr Lys Val Ile Gly
      20      25      30
Leu Phe Met Lys Asn Trp Glu Glu Asp Ser Glu Gly Gly Leu Cys Ser
      35      40      45
Ser Thr Lys Asp Tyr Glu Asp Val Glu Arg Val Cys Leu Gln Leu Asp
      50      55      60
Ile Pro Tyr Tyr Thr Val Ser Phe Ala Lys Glu Tyr Arg Glu Arg Val
      65      70      75      80
Phe Ala Arg Phe Leu Lys Glu Tyr Ser Leu Gly Tyr Thr Pro Asn Pro
      85      90      95
Asp Ile Leu Cys Asn Arg Glu Ile Lys Phe Asp Leu Leu Gln Lys Lys

```

```

      100      105      110
Val Gln Glu Leu Gly Gly Asp Tyr Leu Ala Thr Gly His Tyr Cys Arg
      115      120      125
Leu Asn Thr Glu Leu Gln Glu Thr Gln Leu Leu Arg Gly Cys Asp Pro
      130      135      140
Gln Lys Asp Gln Ser Tyr Phe Leu Ser Gly Thr Pro Lys Ser Ala Leu
145      150      155      160
His Asn Val Leu Phe Pro Leu Gly Glu Met Asn Lys Thr Glu Val Arg
      165      170      175
Ala Ile Ala Ala Gln Ala Ala Leu Pro Thr Ala Glu Lys Lys Asp Ser
      180      185      190
Thr Gly Ile Cys Phe Ile Gly Lys Arg Pro Phe Lys Glu Phe Leu Glu
      195      200      205
Lys Phe Leu Pro Asn Lys Thr Gly Asn Val Ile Asp Trp Asp Thr Lys
210      215      220
Glu Ile Val Gly Gln His Gln Gly Ala His Tyr Tyr Thr Ile Gly Gln
225      230      235      240
Arg Arg Gly Leu Asp Leu Gly Gly Ser Glu Lys Pro Cys Tyr Val Val
      245      250      255
Gly Lys Asn Ile Glu Glu Asn Ser Ile Tyr Ile Val Arg Gly Glu Asp
      260      265      270
His Pro Gln Leu Tyr Leu Arg Glu Leu Thr Ala Arg Glu Leu Asn Trp
      275      280      285
Phe Thr Pro Pro Lys Ser Gly Cys His Cys Ser Ala Lys Val Arg Tyr
290      295      300
Arg Ser Pro Asp Glu Ala Cys Thr Ile Asp Tyr Ser Ser Gly Asp Glu
305      310      315      320
Val Lys Val Arg Phe Ser Gln Pro Val Lys Ala Val Thr Pro Gly Gln
      325      330      335
Thr Ile Ala Phe Tyr Gln Gly Asp Thr Cys Leu Gly Ser Gly Val Ile
      340      345      350
Asp Val Pro Met Ile Pro Ser Glu Gly
      355      360

```

<210> 75
 <211> 1609
 <212> PRT
 <213> Chlamydia pneumoniae

```

<400> 75
Met Val Ala Lys Lys Thr Val Arg Ser Tyr Arg Ser Ser Phe Ser His
      5      10      15
Ser Val Ile Val Ala Ile Leu Ser Ala Gly Ile Ala Phe Glu Ala His
      20      25      30
Ser Leu His Ser Ser Glu Leu Asp Leu Gly Val Phe Asn Lys Gln Phe
      35      40      45
Glu Glu His Ser Ala His Val Glu Glu Ala Gln Thr Ser Val Leu Lys
      50      55      60
Gly Ser Asp Pro Val Asn Pro Ser Gln Lys Glu Ser Glu Lys Val Leu
      65      70      75      80
Tyr Thr Gln Val Pro Leu Thr Gln Gly Ser Ser Gly Glu Ser Leu Asp
      85      90      95
Leu Ala Asp Ala Asn Phe Leu Glu His Phe Gln His Leu Phe Glu Glu
      100      105      110
Thr Thr Val Phe Gly Ile Asp Gln Lys Leu Val Trp Ser Asp Leu Asp
      115      120      125
Thr Arg Asn Phe Ser Gln Pro Thr Gln Glu Pro Asp Thr Ser Asn Ala
      130      135      140
Val Ser Glu Lys Ile Ser Ser Asp Thr Lys Glu Asn Arg Lys Asp Leu
145      150      155      160

```

Glu Thr Glu Asp Pro Ser Lys Lys Ser Gly Leu Lys Glu Val Ser Ser
 165 170 175
 Asp Leu Pro Lys Ser Pro Glu Thr Ala Val Ala Ala Ile Ser Glu Asp
 180 185 190
 Leu Glu Ile Ser Glu Asn Ile Ser Ala Arg Asp Pro Leu Gln Gly Leu
 195 200 205
 Ala Phe Phe Tyr Lys Asn Thr Ser Ser Gln Ser Ile Ser Glu Lys Asp
 210 215 220
 Ser Ser Phe Gln Gly Ile Ile Phe Ser Gly Ser Gly Ala Asn Ser Gly
 225 230 235 240
 Leu Gly Phe Glu Asn Leu Lys Ala Pro Lys Ser Gly Ala Ala Val Tyr
 245 250 255
 Ser Asp Arg Asp Ile Val Phe Glu Asn Leu Val Lys Gly Leu Ser Phe
 260 265 270
 Ile Ser Cys Glu Ser Leu Glu Asp Gly Ser Ala Ala Gly Val Asn Ile
 275 280 285
 Val Val Thr His Cys Gly Asp Val Thr Leu Thr Asp Cys Ala Thr Gly
 290 295 300
 Leu Asp Leu Glu Ala Leu Arg Leu Val Lys Asp Phe Ser Arg Gly Gly
 305 310 315 320
 Ala Val Phe Thr Ala Arg Asn His Glu Val Gln Asn Asn Leu Ala Gly
 325 330 335
 Gly Ile Leu Ser Val Val Gly Asn Lys Gly Ala Ile Val Val Glu Lys
 340 345 350
 Asn Ser Ala Glu Lys Ser Asn Gly Gly Ala Phe Ala Cys Gly Ser Phe
 355 360 365
 Val Tyr Ser Asn Asn Glu Asn Thr Ala Leu Trp Lys Glu Asn Gln Ala
 370 375 380
 Leu Ser Gly Gly Ala Ile Ser Ser Ala Ser Asp Ile Asp Ile Gln Gly
 385 390 395 400
 Asn Cys Ser Ala Ile Glu Phe Ser Gly Asn Gln Ser Leu Ile Ala Leu
 405 410 415
 Gly Glu His Ile Gly Leu Thr Asp Phe Val Gly Gly Gly Ala Leu Ala
 420 425 430
 Ala Gln Gly Thr Leu Thr Leu Arg Asn Asn Ala Val Val Gln Cys Val
 435 440 445
 Lys Asn Thr Ser Lys Thr His Gly Gly Ala Ile Leu Ala Gly Thr Val
 450 455 460
 Asp Leu Asn Glu Thr Ile Ser Glu Val Ala Phe Lys Gln Asn Thr Ala
 465 470 475 480
 Ala Leu Thr Gly Gly Ala Leu Ser Ala Asn Asp Lys Val Ile Ile Ala
 485 490 495
 Asn Asn Phe Gly Glu Ile Leu Phe Glu Gln Asn Glu Val Arg Asn His
 500 505 510
 Gly Gly Ala Ile Tyr Cys Gly Cys Arg Ser Asn Pro Lys Leu Glu Gln
 515 520 525
 Lys Asp Ser Gly Glu Asn Ile Asn Ile Ile Gly Asn Ser Gly Ala Ile
 530 535 540
 Thr Phe Leu Lys Asn Lys Ala Ser Val Leu Glu Val Met Thr Gln Ala
 545 550 555 560
 Glu Asp Tyr Ala Gly Gly Gly Ala Leu Trp Gly His Asn Val Leu Leu
 565 570 575
 Asp Ser Asn Ser Gly Asn Ile Gln Phe Ile Gly Asn Ile Gly Gly Ser
 580 585 590
 Thr Phe Trp Ile Gly Glu Tyr Val Gly Gly Gly Ala Ile Leu Ser Thr
 595 600 605
 Asp Arg Val Thr Ile Ser Asn Asn Ser Gly Asp Val Val Phe Lys Gly
 610 615 620
 Asn Lys Gly Gln Cys Leu Ala Gln Lys Tyr Val Ala Pro Gln Glu Thr
 625 630 635 640
 Ala Pro Val Glu Ser Asp Ala Ser Ser Thr Asn Lys Asp Glu Lys Ser

					645						650					655
Leu	Asn	Ala	Cys	Ser	His	Gly	Asp	His	Tyr	Pro	Pro	Lys	Thr	Val	Glu	
			660					665					670			
Glu	Glu	Val	Pro	Pro	Ser	Leu	Leu	Glu	Glu	His	Pro	Val	Val	Ser	Ser	
		675					680					685				
Thr	Asp	Ile	Arg	Gly	Gly	Gly	Ala	Ile	Leu	Ala	Gln	His	Ile	Phe	Ile	
	690					695					700					
Thr	Asp	Asn	Thr	Gly	Asn	Leu	Arg	Phe	Ser	Gly	Asn	Leu	Gly	Gly	Gly	
705					710					715						720
Glu	Glu	Ser	Ser	Thr	Val	Gly	Asp	Leu	Ala	Ile	Val	Gly	Gly	Gly	Gly	Ala
				725					730						735	
Leu	Leu	Ser	Thr	Asn	Glu	Val	Asn	Val	Cys	Ser	Asn	Gln	Asn	Val	Val	
			740					745					750			
Phe	Ser	Asp	Asn	Val	Thr	Ser	Asn	Gly	Cys	Asp	Ser	Gly	Gly	Ala	Ile	
		755					760					765				
Leu	Ala	Lys	Lys	Val	Asp	Ile	Ser	Ala	Asn	His	Ser	Val	Glu	Phe	Val	
	770					775					780					
Ser	Asn	Gly	Ser	Gly	Lys	Phe	Gly	Gly	Ala	Val	Cys	Ala	Leu	Asn	Glu	
785					790					795					800	
Ser	Val	Asn	Ile	Thr	Asp	Asn	Gly	Ser	Ala	Val	Ser	Phe	Ser	Lys	Asn	
				805					810					815		
Arg	Thr	Arg	Leu	Gly	Gly	Ala	Gly	Val	Ala	Ala	Pro	Gln	Gly	Ser	Val	
			820					825					830			
Thr	Ile	Cys	Gly	Asn	Gln	Gly	Asn	Ile	Ala	Phe	Lys	Glu	Asn	Phe	Val	
		835					840					845				
Phe	Gly	Ser	Glu	Asn	Gln	Arg	Ser	Gly	Gly	Gly	Ala	Ile	Ile	Ala	Asn	
	850					855					860					
Ser	Ser	Val	Asn	Ile	Gln	Asp	Asn	Ala	Gly	Asp	Ile	Leu	Phe	Val	Ser	
865					870					875					880	
Asn	Ser	Thr	Gly	Ser	Tyr	Gly	Gly	Ala	Ile	Phe	Val	Gly	Ser	Leu	Val	
				885					890					895		
Ala	Ser	Glu	Gly	Ser	Asn	Pro	Arg	Thr	Leu	Thr	Ile	Thr	Gly	Asn	Ser	
			900					905					910			
Gly	Asp	Ile	Leu	Phe	Ala	Lys	Asn	Ser	Thr	Gln	Thr	Ala	Ala	Ser	Leu	
		915					920					925				
Ser	Glu	Lys	Asp	Ser	Phe	Gly	Gly	Gly	Ala	Ile	Tyr	Thr	Gln	Asn	Leu	
					935						940					
Lys	Ile	Val	Lys	Asn	Ala	Gly	Asn	Val	Ser	Phe	Tyr	Gly	Asn	Arg	Ala	
945					950					955					960	
Pro	Ser	Gly	Ala	Gly	Val	Gln	Ile	Ala	Asp	Gly	Gly	Thr	Val	Cys	Leu	
				965					970					975		
Glu	Ala	Phe	Gly	Gly	Asp	Ile	Leu	Phe	Glu	Gly	Asn	Ile	Asn	Phe	Asp	
			980													

Val Ser Ala Gly Val Gln Ile Asn Met Ser Ser Pro Thr Pro Asn Lys
 1140 1145 1150
 Asp Lys Ala Val Asp Thr Pro Val Leu Ala Asp Ile Ile Ser Ile Thr
 1155 1160 1165
 Val Asp Leu Ser Ser Phe Val Pro Glu Gln Asp Gly Thr Leu Pro Leu
 1170 1175 1180
 Pro Pro Glu Ile Ile Ile Pro Lys Gly Thr Lys Leu His Ser Asn Ala
 1185 1190 1195 1200
 Ile Asp Leu Lys Ile Ile Asp Pro Thr Asn Val Gly Tyr Glu Asn His
 1205 1210 1215
 Ala Leu Leu Ser Ser His Lys Asp Ile Pro Leu Ile Ser Leu Lys Thr
 1220 1225 1230
 Ala Glu Gly Met Thr Gly Thr Pro Thr Ala Asp Ala Ser Leu Ser Asn
 1235 1240 1245
 Ile Lys Ile Asp Val Ser Leu Pro Ser Ile Thr Pro Ala Thr Tyr Gly
 1250 1255 1260
 His Thr Gly Val Trp Ser Glu Ser Lys Met Glu Asp Gly Arg Leu Val
 1265 1270 1275 1280
 Val Gly Trp Gln Pro Thr Gly Tyr Lys Leu Asn Pro Glu Lys Gln Gly
 1285 1290 1295
 Ala Leu Val Leu Asn Asn Leu Trp Ser His Tyr Thr Asp Leu Arg Ala
 1300 1305 1310
 Leu Lys Gln Glu Ile Phe Ala His His Thr Ile Ala Gln Arg Met Glu
 1315 1320 1325
 Leu Asp Phe Ser Thr Asn Val Trp Gly Ser Gly Leu Gly Val Val Glu
 1330 1335 1340
 Asp Cys Gln Asn Ile Gly Glu Phe Asp Gly Phe Lys His His Leu Thr
 1345 1350 1355 1360
 Gly Tyr Ala Leu Gly Leu Asp Thr Gln Leu Val Glu Asp Phe Leu Ile
 1365 1370 1375
 Gly Gly Cys Phe Ser Gln Phe Phe Gly Lys Thr Glu Ser Gln Ser Tyr
 1380 1385 1390
 Lys Ala Lys Asn Asp Val Lys Ser Tyr Met Gly Ala Ala Tyr Ala Gly
 1395 1400 1405
 Ile Leu Ala Gly Pro Trp Leu Ile Lys Gly Ala Phe Val Tyr Gly Asn
 1410 1415 1420
 Ile Asn Asn Asp Leu Thr Thr Asp Tyr Gly Thr Leu Gly Ile Ser Thr
 1425 1430 1435 1440
 Gly Ser Trp Ile Gly Lys Gly Phe Ile Ala Gly Thr Ser Ile Asp Tyr
 1445 1450 1455
 Arg Tyr Ile Val Asn Pro Arg Arg Phe Ile Ser Ala Ile Val Ser Thr
 1460 1465 1470
 Val Val Pro Phe Val Glu Ala Glu Tyr Val Arg Ile Asp Leu Pro Glu
 1475 1480 1485
 Ile Ser Glu Gln Gly Lys Glu Val Arg Thr Phe Gln Lys Thr Arg Phe
 1490 1495 1500
 Glu Asn Val Ala Ile Pro Phe Gly Phe Ala Leu Glu His Ala Tyr Ser
 1505 1510 1515 1520
 Arg Gly Ser Arg Ala Glu Val Asn Ser Val Gln Leu Ala Tyr Val Phe
 1525 1530 1535
 Asp Val Tyr Arg Lys Gly Pro Val Ser Leu Ile Thr Leu Lys Asp Ala
 1540 1545 1550
 Ala Tyr Ser Trp Lys Ser Tyr Gly Val Asp Ile Pro Cys Lys Ala Trp
 1555 1560 1565
 Lys Ala Arg Leu Ser Asn Asn Thr Glu Trp Asn Ser Tyr Leu Ser Thr
 1570 1575 1580
 Tyr Leu Ala Phe Asn Tyr Glu Trp Arg Glu Asp Leu Ile Ala Tyr Asp
 1585 1590 1595 1600
 Phe Asn Gly Gly Ile Arg Ile Ile Phe
 1605

```
<210> 76
<211> 196
<212> PRT
<213> Chlamydia pneumoniae
```

[illegible]

```
<210> 77
<211> 619
<212> PRT
<213> Chlamydia pneumoniae
```

<400>	77															
Met	Lys	Lys	Gly	Lys	Leu	Gly	Ala	Ile	Val	Phe	Gly	Leu	Leu	Phe	Thr	
				5					10					15		
Ser	Ser	Val	Ala	Gly	Phe	Ser	Lys	Asp	Leu	Thr	Lys	Asp	Asn	Ala	Tyr	
			20					25					30			
Gln	Asp	Leu	Asn	Val	Ile	Glu	His	Leu	Ile	Ser	Leu	Lys	Tyr	Ala	Pro	
		35					40					45				
Leu	Pro	Trp	Lys	Glu	Leu	Leu	Phe	Gly	Trp	Asp	Leu	Ser	Gln	Gln	Thr	
	50					55					60					
Gln	Gln	Ala	Arg	Leu	Gln	Leu	Val	Leu	Glu	Glu	Lys	Pro	Thr	Thr	Asn	
	65				70					75					80	
Tyr	Cys	Gln	Lys	Val	Leu	Ser	Asn	Tyr	Val	Arg	Ser	Leu	Asn	Asp	Tyr	
			85					90					95			
His	Ala	Gly	Ile	Thr	Phe	Tyr	Arg	Thr	Glu	Ser	Ala	Tyr	Ile	Pro	Tyr	
		100						105					110			
Val	Leu	Lys	Leu	Ser	Glu	Asp	Gly	His	Val	Phe	Val	Val	Asp	Val	Gln	
		115					120					125				
Thr	Ser	Gln	Gly	Asp	Ile	Tyr	Leu	Gly	Asp	Glu	Ile	Leu	Glu	Val	Asp	
	130					135				140						
Gly	Met	Gly	Ile	Arg	Glu	Ala	Ile	Glu	Ser	Leu	Arg	Phe	Gly	Arg	Gly	
	145				150					155					160	

Ser Ala Thr Asp Tyr Ser Ala Ala Val Arg Ser Leu Thr Ser Arg Ser
 165 170 175
 Ala Ala Phe Gly Asp Ala Val Pro Ser Gly Ile Ala Met Leu Lys Leu
 180 185 190
 Arg Arg Pro Ser Gly Leu Ile Arg Ser Thr Pro Val Arg Trp Arg Tyr
 195 200 205
 Thr Pro Glu His Ile Gly Asp Phe Ser Leu Val Ala Pro Leu Ile Pro
 210 215 220
 Glu His Lys Pro Gln Leu Pro Thr Gln Ser Cys Val Leu Phe Arg Ser
 225 230 235 240
 Gly Val Asn Ser Gln Ser Ser Ser Ser Ser Leu Phe Ser Ser Tyr Met
 245 250 255
 Val Pro Tyr Phe Trp Glu Glu Leu Arg Val Gln Asn Lys Gln Arg Phe
 260 265 270
 Asp Ser Asn His His Ile Gly Ser Arg Asn Gly Phe Leu Pro Thr Phe
 275 280 285
 Gly Pro Ile Leu Trp Glu Gln Asp Lys Gly Pro Tyr Arg Ser Tyr Ile
 290 295 300
 Phe Lys Ala Lys Asp Ser Gln Gly Asn Pro His Arg Ile Gly Phe Leu
 305 310 315 320
 Arg Ile Ser Ser Tyr Val Trp Thr Asp Leu Glu Gly Leu Glu Glu Asp
 325 330 335
 His Lys Asp Ser Pro Trp Glu Leu Phe Gly Glu Ile Ile Asp His Leu
 340 345 350
 Glu Lys Glu Thr Asp Ala Leu Ile Ile Asp Gln Thr His Asn Pro Gly
 355 360 365
 Gly Ser Val Phe Tyr Leu Tyr Ser Leu Leu Ser Met Leu Thr Asp His
 370 375 380
 Pro Leu Asp Thr Pro Lys His Arg Met Ile Phe Thr Gln Asp Glu Val
 385 390 395 400
 Ser Ser Ala Leu His Trp Gln Asp Leu Leu Glu Asp Val Phe Thr Asp
 405 410 415
 Glu Gln Ala Val Ala Val Leu Gly Glu Thr Met Glu Gly Tyr Cys Met
 420 425 430
 Asp Met His Ala Val Ala Ser Leu Gln Asn Phe Ser Gln Ser Val Leu
 435 440 445
 Ser Ser Trp Val Ser Gly Asp Ile Asn Leu Ser Lys Pro Met Pro Leu
 450 455 460
 Leu Gly Phe Ala Gln Val Arg Pro His Pro Lys His Gln Tyr Thr Lys
 465 470 475 480
 Pro Leu Phe Met Leu Ile Asp Glu Asp Asp Phe Ser Cys Gly Asp Leu
 485 490 495
 Ala Pro Ala Ile Leu Lys Asp Asn Gly Arg Ala Thr Leu Ile Gly Lys
 500 505 510
 Pro Thr Ala Gly Ala Gly Gly Phe Val Phe Gln Val Thr Phe Pro Asn
 515 520 525
 Arg Ser Gly Ile Lys Gly Leu Ser Leu Thr Gly Ser Leu Ala Val Arg
 530 535 540
 Lys Asp Gly Glu Phe Ile Glu Asn Leu Gly Val Ala Pro His Ile Asp
 545 550 555 560
 Leu Gly Phe Thr Ser Arg Asp Leu Gln Thr Ser Arg Phe Thr Asp Tyr
 565 570 575
 Val Glu Ala Val Lys Thr Ile Val Leu Thr Ser Leu Ser Glu Asn Ala
 580 585 590
 Lys Lys Ser Glu Glu Gln Thr Ser Pro Gln Glu Thr Pro Glu Val Ile
 595 600 605
 Arg Val Ser Tyr Pro Thr Thr Ser Ala Ser
 610 615

<211> 651

<212> PRT

<213> Chlamydia pneumoniae

<400> 78

```

Met Val Asn Pro Ile Gly Pro Gly Pro Ile Asp Glu Thr Glu Arg Thr
      5      10
Pro Pro Ala Asp Leu Ser Ala Gln Gly Leu Glu Ala Ser Ala Ala Asn
      20      25      30
Lys Ser Ala Glu Ala Gln Arg Ile Ala Gly Ala Glu Ala Lys Pro Lys
      35      40      45
Glu Ser Lys Thr Asp Ser Val Glu Arg Trp Ser Ile Leu Arg Ser Ala
      50      55      60
Val Asn Ala Leu Met Ser Leu Ala Asp Lys Leu Gly Ile Ala Ser Ser
      65      70      75      80
Asn Ser Ser Ser Ser Thr Ser Arg Ser Ala Asp Val Asp Ser Thr Thr
      85      90      95
Ala Thr Ala Pro Thr Pro Pro Pro Thr Phe Asp Asp Tyr Lys Thr
      100      105      110
Gln Ala Gln Thr Ala Tyr Asp Thr Ile Phe Thr Ser Thr Ser Leu Ala
      115      120      125
Asp Ile Gln Ala Ala Leu Val Ser Leu Gln Asp Ala Val Thr Asn Ile
      130      135      140
Lys Asp Thr Ala Ala Thr Asp Glu Glu Thr Ala Ile Ala Ala Glu Trp
      145      150      155      160
Glu Thr Lys Asn Ala Asp Ala Val Lys Val Gly Ala Gln Ile Thr Glu
      165      170      175
Leu Ala Lys Tyr Ala Ser Asp Asn Gln Ala Ile Leu Asp Ser Leu Gly
      180      185      190
Lys Leu Thr Ser Phe Asp Leu Leu Gln Ala Ala Leu Leu Gln Ser Val
      195      200      205
Ala Asn Asn Asn Lys Ala Ala Glu Leu Leu Lys Glu Met Gln Asp Asn
      210      215      220
Pro Val Val Pro Gly Lys Thr Pro Ala Ile Ala Gln Ser Leu Val Asp
      225      230      235      240
Gln Thr Asp Ala Thr Ala Thr Gln Ile Glu Lys Asp Gly Asn Ala Ile
      245      250      255
Arg Asp Ala Tyr Phe Ala Gly Gln Asn Ala Ser Gly Ala Val Glu Asn
      260      265      270
Ala Lys Ser Asn Asn Ser Ile Ser Asn Ile Asp Ser Ala Lys Ala Ala
      275      280      285
Ile Ala Thr Ala Lys Thr Gln Ile Ala Glu Ala Gln Lys Lys Phe Pro
      290      295      300
Asp Ser Pro Ile Leu Gln Glu Ala Glu Gln Met Val Ile Gln Ala Glu
      305      310      315      320
Lys Asp Leu Lys Asn Ile Lys Pro Ala Asp Gly Ser Asp Val Pro Asn
      325      330      335
Pro Gly Thr Thr Val Gly Gly Ser Lys Gln Gln Gly Ser Ser Ile Gly
      340      345      350
Ser Ile Arg Val Ser Met Leu Leu Asp Asp Ala Glu Asn Glu Thr Ala
      355      360      365
Ser Ile Leu Met Ser Gly Phe Arg Gln Met Ile His Met Phe Asn Thr
      370      375      380
Glu Asn Pro Asp Ser Gln Ala Ala Gln Gln Glu Leu Ala Ala Gln Ala
      385      390      395      400
Arg Ala Ala Lys Ala Ala Gly Asp Asp Ser Ala Ala Ala Ala Leu Ala
      405      410      415
Asp Ala Gln Lys Ala Leu Glu Ala Ala Leu Gly Lys Ala Gly Gln Gln
      420      425      430
Gln Gly Ile Leu Asn Ala Leu Gly Gln Ile Ala Ser Ala Ala Val Val
      435      440      445

```

```

Ser Ala Gly Val Pro Pro Ala Ala Ala Ser Ser Ile Gly Ser Ser Val
  450                455                460
Lys Gln Leu Tyr Lys Thr Ser Lys Ser Thr Gly Ser Asp Tyr Lys Thr
  465                470                475                480
Gln Ile Ser Ala Gly Tyr Asp Ala Tyr Lys Ser Ile Asn Asp Ala Tyr
  485                490                495
Gly Arg Ala Arg Asn Asp Ala Thr Arg Asp Val Ile Asn Asn Val Ser
  500                505                510
Thr Pro Ala Leu Thr Arg Ser Val Pro Arg Ala Arg Thr Glu Ala Arg
  515                520                525
Gly Pro Glu Lys Thr Asp Gln Ala Leu Ala Arg Val Ile Ser Gly Asn
  530                535                540
Ser Arg Thr Leu Gly Asp Val Tyr Ser Gln Val Ser Ala Leu Gln Ser
  545                550                555                560
Val Met Gln Ile Ile Gln Ser Asn Pro Gln Ala Asn Asn Glu Glu Ile
  565                570                575
Arg Gln Lys Leu Thr Ser Ala Val Thr Lys Pro Pro Gln Phe Gly Tyr
  580                585                590
Pro Tyr Val Gln Leu Ser Asn Asp Ser Thr Gln Lys Phe Ile Ala Lys
  595                600                605
Leu Glu Ser Leu Phe Ala Glu Gly Ser Arg Thr Ala Ala Glu Ile Lys
  610                615                620
Ala Leu Ser Phe Glu Thr Asn Ser Leu Phe Ile Gln Gln Val Leu Val
  625                630                635                640
Asn Ile Gly Ser Leu Tyr Ser Gly Tyr Leu Gln
  645                650

```

<210> 79
 <211> 87
 <212> PRT
 <213> *Chlamydia pneumoniae*

```

<400> 79
Met Ser Gln Lys Asn Lys Asn Ser Ala Phe Met His Pro Val Asn Ile
      5                10                15
Ser Thr Asp Leu Ala Val Ile Val Gly Lys Gly Pro Met Pro Arg Thr
      20                25                30
Glu Ile Val Lys Lys Val Trp Glu Tyr Ile Lys Lys His Asn Cys Gln
      35                40                45
Asp Gln Lys Asn Lys Arg Asn Ile Leu Pro Asp Ala Asn Leu Ala Lys
      50                55                60
Val Phe Gly Ser Ser Asp Pro Ile Asp Met Phe Gln Met Thr Lys Ala
      65                70                75                80
Leu Ser Lys His Ile Val Lys
      85

```

<210> 80
 <211> 3048
 <212> DNA
 <213> *Chlamydia trachomatis* serovar D

```

<400> 80
atgccttttt ctttgagatc tacatcattt tgttttttag cttgtttgtg ttctatttcg 60
tatggatttc cgagctctcc tcaagtgtta acacctaata taaccactcc ttttaagggg 120
gacgatgttt acttgaatgg agactgcgct tttgtcaatg tctatgcagg ggcagagaa 180
ggctcaatta tctcagctaa tggcgacaat ttaacgatta cgggacaaaa ccatacatta 240
tcatttacag atttcaagg gccagttctt caaaattatg ctttcatttc agcaggagag 300
acacttactc tgaaagattt ttcgagtttg atgttctcga aaaatgtttc ttgaggagaa 360
aagggaatga tctcagggaa aaccgtgagt atttcgggag caggcgaagt gatttttttg 420
gataactctg tggggtattc tcctttgtct attgtgccag catcgactcc aactcctcca 480

```

```

gcaccagcac cagctcctgc tgcttcaagc totttatctc caacagttag tgatgctcgg 540
aaaggggtcta ttttttctgt agagactagt ttggagatct caggcgtcaa aaaaggggtc 600
atgttcgata ataatgccgg gaatttttga acagtttttc gaggtaatag taataataat 660
gctggtagtg ggggtagtgg gtctgtctaca acaccaagtt ttacagttaa aaactgtaaa 720
gggaaagttt ctttcacaga taacgtagcc tcotgtggag goggagtagt ctacaaagga 780
actgtgcttt tcaaagacaa tgaaggagggc atattcttcc gagggaaacac agcatacgt 840
gatttaggga ttcttgcctgc tactagtcgg gatcagaata cggagacagg aggcggtgga 900
ggagttatatt gctctccaga tgattctgta aagtttgaag gcaataaagg ttctattgtt 960
tttgattaca actttgcaaa aggcagagggc ggaagcatcc taacgaaaga attctctctt 1020
gtagcagatg attcggttgt ctttagtaac aatacagcag aaaaaggcgg tggagctatt 1080
tatgctccta ctatcgatat aagcacgaat ggaggatcga ttctatttga aagaaaccga 1140
gctgcagaag gaggcggcat ctgctgtagt gaagcaagct ctggttcaac tggaaatctt 1200
actttaagcg cttctgatgg ggatattgtt ttttctggga atatgacgag tgatcgctct 1260
ggagagcgca gcgcagcaag aatcttaagt gatggaacga ctgtttcttt aaatgcttcc 1320
ggactatcga agctgatctt ttatgatcct cctgggtgctg tcacgattaa tcagtccggt 1440
tcgacaccat caccatcttc ttcttctatg ttgactcctt cagaaaaact tcaagttctt 1500
aatggatctg tgatttttac cgcgagtcga actgtgtcag gaggggagtt ggttgtgacg 1560
aactctactt ctaacttccc aggagctctg actgtgtcag ctggacgagt gactttagga 1620
gaaggagcta ccttaactac tgggaccatt acagccacct gctgcaaata ataattatac ttgtacagta 1680
tccggagctt cgttgtctgc cgttgccagggt ttaactccta actataagac ggccatactg 1740
tctaagttgg ggattgattt agaactcctt ggctctactt tagacctagt gatggagagt 1800
ggtgoggatg gaacagttac tgtaaacagc ggctctactt tgacaattcc ttttgttact 1860
gagggcagagg tatatgataa tccgcttttt gtgggatcgc acaaaaaatt ctgtcactat taatgatgca 1920
ctatcttcta gtagtgctag taacggagtt tggctctgcag attggacgaa accgcctctg 1980
gaogctgocg actatgggta tcaaggctct aataccaata acactctgta tctgacatgg 2040
gctoctgatg ctaaggggat ggtacctcct ctggatcctc agagaaaggg agaactagta 2100
agacctgctt tttgggtagc gggatctgca ttaagaacct ttactaatgg tttgaaagaa 2160
cccaactctc ctagagatgt tggatttgta gcatctctgc atgctctcgg ggattatatt 2220
cactatgttt cgaagatga tcgggatggc ttttttagcta gatatggggg attccaggcg 2280
ttgaattata ccgattatga aaatgggtca atatttggag tggcttttgg acaactctat 2340
accgcagcct agagcagaat gtattactct aaagatgctg ggaacatgac gatgttgtcc 2400
ggtcagacaa gaagttacgt agatattaaa ggaacagaaa ctgttatgta ttgggagacg 2460
tgtttcggaa attctgtgca cagaatgcat acgcagtatt ttaatgacaa aacgcagaag 2520
ttcgatcatt cgaatgtca ttggcacaac aataactatt atgctgttgt ggtgcccag 2580
cataattttc tagagtactg cattcctact cgtcagttcg ctagagatta tgagcttaca 2640
gggtttatgc gttttgaaat ggccggagga tgggtccagtt ctacacgaga aactggctcc 2700
ctaactagat atttcgcctg cgggtcaggg cataatatgt cgcttccaat aggaattgta 2760
gctcatgcag tttctcatgt gcgaagatct cctccttcta aactgacact aaatatggga 2820
tatagaccag acatttggcg tgtcactcca cattgcaata tggaaattat tgctaacgga 2880
gtgaagacac ctatacaagg atctccgctg gcacggcatg ccttcttctt agaagtgcab 2940
gatactttgt atattcatca ttttggaga gcctatatga actattcgct ggatgctcgt 3000
cgctcagacaaa cggcacattt tgtatccatg ggcttgaata gaatcttt 3048

```

<210> 81

<211> 1038

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 81

```

atgcaagcag atattttaga tggaaaacag aaacgcgtta atctaaatag caagcgtcta 60
gtgaactgca accagggtga tgtcaaccaa cttgttccta ttaagtacaa atgggcttgg 120
gaacattatt tgaatggctg cgcataaac tggctcccta cagagatccc catggggaaa 180
gacatcgaaat tatggaagtc ggatcgctct tctgaagatg agcggcagat cattcttttg 240
aatttaggtt ttttcagcac gcgagagagc ttgggttggga ataattattgt tctagcaatt 300
tttaaacatg taactaatcc ggaagcgaga caatatcttt taagacaagc ttttgaagaa 360
gcggttcaca cgcacacatt ttgttatatt tgtgagtcac tccgattaga cgagaaagaa 420
attttcaatg cctataacga gcgtgctgcg attaaggcca aagatgattt ccagatggaa 480
atcactggca aggtattaga tcctaatttt gcacaggact ctgttgaggg tctacaggag 540
tttgttaaaa acttagtagg atactacatc attatggaag ggattttctt ctatagtggt 600
tttgtgatga tcctttcctt ccacagacaa aataagatga ttggtatttg agaacaatat 660

```

caatacatct	taagagatga	gacaatccac	ttgaactttg	gtattgattt	gatcaacggg	720
ataaaagaag	agaaccggga	gatttggact	ccagagttac	agcaagaaat	tgtcgaatta	780
attaagcgag	ctgtcgattt	agaaattgag	tatgcgcaag	actgtctccc	tagagggatt	840
ttgggattga	gagcttcgat	gttcacgat	tatgtgcagc	atattgcaga	ccgtcgtttg	900
gaaagaatcg	gattaaaacc	tatttatcat	acgaaaaacc	cattcccttg	gatgagcgaa	960
acaatagacc	ttaataaaga	gaaaaacttc	tttgaaacaa	gggttataga	atatcaacat	1020
gcagcaagct	taacttgg					1038

<210> 82

<211> 3159

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 82

atgtttacaa	ggatagttat	ggtcgatcta	caagaaaagc	aatgcacaat	tgtaagcgc	60
aatggaatgt	ttgttccttt	cgatcggaac	cgtatttttc	aggctttaga	agcagctttt	120
cgagacactc	gcagaattga	tgatcatatg	cctttgcctg	aagatctgga	aagttccata	180
cgctcgataa	cgcatcaggt	agttaaagaa	gttgtgcaaa	agattacaga	tggacaagtg	240
gttactgtag	agcgtatcca	agatatgggt	gaaagccaac	tatatgtgaa	tggtttgcaa	300
gatgtttgctc	gcgattatata	tgtctatcgc	gatgaccgta	aagcgcatcg	gaaaaaatct	360
tggcaagcc	tatccgttgt	tcgtcgttgt	gggactgttg	tacactttaa	tcctatgaaa	420
atttccgcgc	ctttggaaaa	agctttccga	gctaccgata	agactgaggg	gatgactcca	480
agttctgtgc	gagaggaaat	caatgctttg	acgcaaaaca	ttgtcgcgga	aatagaagaa	540
tggttgcctc	aacaggatag	acgcattgat	atcgagaaga	ttcaagatat	tgttgaacag	600
caactaatgg	ttgttgggca	ttatgctggt	gcaaagaact	atattottta	tcgagaagct	660
cgcgctcgtg	ttcgtgataa	cagagaagag	gacgggagta	cagaaaagac	tatagcagaa	720
gaagctgttg	aggtgctcag	taaagacggt	tctacctata	caatgacgca	ttcgcagttg	780
ttggctcatt	tagcgcgcgc	ttgtagtctg	tttccagaaa	cgacagatgc	ggcgctgctt	840
accgatatgg	cttttcgcaa	tttctatttc	gggtatcaaag	agtctgaagt	agtactggcc	900
tgtattatgg	cggctcgtgc	caatattgaa	aaggagcctg	attatgcctt	tgttgctgca	960
gagctcttac	ttgacgttgt	atataaggaa	gcgttaggga	aatcgaaata	tgotgaggat	1020
ttagaacaag	cacatcgcca	tcattttcaaa	cgctacatcg	cagaagggga	tacctatcgt	1080
ctgaatgctg	aactgaaaca	tctttttgat	ttagacgcgt	tagccgatgc	tatggatcta	1140
tctcgagatc	tacagttttc	ttacatgggt	aattcaaaatc	tgtatgatcg	ttatttttaa	1200
caccacgaag	gttgccgttt	agaaactccc	caaatttttt	ggatgcgcgt	tgctatgggg	1260
ttggcattga	atgagcaaga	caagacttct	tgggctatta	ctttttataa	tttgctttcg	1320
acattccgat	atacaccagc	tacgcccaacc	ttgttcaatt	caggtatgcg	gcattctcag	1380
ttaagctctt	gctatctttc	caactgtacaa	gataatttgg	tcaatatcta	taaggctcatt	1440
gctgataacg	ctatgctatc	taagtgggca	ggagggtag	gtaatgattg	gacggcgagt	1500
cgtgcaacag	gggctttaat	ttaaaggaaac	aatggaagaa	gtcagggagt	aattcctttt	1560
attaaggtga	caaatgatac	agcagtcgca	gtgaatcaag	gtggtaaacg	caagggagct	1620
gtatgcgtct	atttagaagt	ttggcacctc	gactacgaag	atttccttga	attgagaaag	1680
aatacagggg	atgagcgtcg	acgggctcat	gatgtcaata	tagctagctg	gattccagat	1740
cttttcttca	aacgtttaca	gcaaaaaggg	acatggactc	tattcagccc	agatgatgtt	1800
ccgggattac	acgatgctta	tggggaagaa	tttgagcgtt	tgtacgaaga	atatgagcgg	1860
aaggttgata	ccggagagat	tcgggttattc	aagaaggtag	aagctgaaga	tctgtggaga	1920
aaaatgctca	gcatgctttt	tgaacgggga	cacccatgga	tgacttttaa	agatccatcc	1980
aacatccgtt	cggctcaaga	tcataaaggc	gtggtgcgtt	gttccaatct	gtgtacggag	2040
atthttgttaa	actgctcgga	gacagaaact	gctgtttgta	atttaggata	gattaactta	2100
gttcaacata	tcgtagggga	tgggttagat	gaggaaaaac	tctctgagac	gatctctata	2160
gcagtcgcta	tggttgataa	cgtgattgat	attaactttt	atccaacaaa	ggaagctaaa	2220
gaggcgaaat	ttgctcaccg	cgctatttga	ttagggggtga	tgggattcca	agatgccttg	2280
tataagctag	atataagcta	tgcttcgcaa	gaagctgtag	aatttgctga	ctacagttca	2340
gagttgatth	cttactatgc	gattcaagct	tcttgtctgc	tcgctaaaaga	acgaggcact	2400
tacagctctt	ataaaggatc	gaaatgggat	agaggtttgc	tccttattga	tacgattcag	2460
ttgttagcga	actatcgagg	agaagcaaat	ctccagatgg	atacgtcatc	aagaaaagat	2520
tgggaaccta	tcogtagttt	ggttaaagag	catggtatgc	gacattgtca	gcttatggct	2580
atagctccga	cacgagcat	ctccaacatt	ataggagtaa	ctcaatctat	tgagccaacg	2640
tacaaacatt	tgthttgtgaa	gtotaatttg	tccggagaat	tcacgattcc	aaatgtgtat	2700
ttaattgaga	agttgaagaa	attaggtatc	tgggatgctg	atatgttaga	tgacctgaaa	2760
tatttttgatg	ggtctttatt	ggaaatcgag	cgtataccag	atcacttaaa	acatattttc	2820

ttgacagctt	ttgagattga	accagaatgg	attatcgaat	gcgcgtctcg	aagacaaaaa	2880
tggattgata	tggggcaatc	cctcaacctt	tatcttgccc	agccagacgg	gaaaaaactg	2940
tcgaatatgt	atttaacggc	ttggaaaaaa	ggtttgaaaa	ctacgtatta	tctgagatct	3000
tcatacagca	cgaccgttga	aaaatctttt	gtagatatta	ataagagagg	aattcagcct	3060
cgttggatga	agaataagtc	tgcttcggca	ggaattattg	ttgaaagagc	gaagaaagca	3120
cctgtctgtt	ctttggaaga	agggtgtgaa	gcatgtcag			3159

<210> 83

<211> 4593

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 83

atgagttccg	agaaagatat	aaaaagcacc	tgttctaagt	tttctttgtc	tgtagtagca	60
gctatccttg	cctctgttag	cgggttagct	agttgcgtag	atcttcatgc	tggaggacag	120
tctgtaaatg	agctggata	tgtaggccct	caagcggtt	tattgttaga	ccaaattcga	180
gatctattcg	ttgggtctaa	agatagtcag	gctgaaggac	agtatagggt	aattgttaga	240
gatccaagtt	ctttccaaga	gaaagatgcg	gatactcttc	cggggaagg	agagcaaagt	300
actttgttct	cagtaaccaa	tcccggtggt	ttccaagggt	tggaccaaca	ggtcaagtc	360
tcttcccaag	ggttaatttg	tagttttacg	agcagcaacc	ttgattctcc	tcgtgacgga	420
gaatcttttt	taggtattgc	ttttgttggg	gatagtagta	aggctggaat	cacattaact	480
gacgtgaaag	cttctttgtc	tggagcggct	ttatattcta	cagaagatct	tatctttgaa	540
aagattaagg	gtggattgga	atgtgcata	tgttcttctc	tagaacagg	gggagcttgt	600
gcagctcaaa	gtattttgat	tcattgattgt	caaggattgc	agggttaaca	ctgtactaca	660
gccgtgaatg	ctgagggggc	tagtgccaat	gatcatcttg	gatttggagg	aggcgctttc	720
tttgttacgg	gttctcttct	tggagagaaa	agtctctata	tgcctgcagg	agatatggta	780
gttgccaatt	gtgatggggc	tatatctttt	gaaggaaaca	gcgcgaactt	tgctaattga	840
ggagcgattg	ctgcctctgg	gaaagtgcct	tttgtcgcta	atgataaaaa	gacttctttt	900
atagagaacc	gagctttgtc	tggaggagcg	attgcagcct	cttctgatat	tgccttttcaa	960
aaactgcgcag	aaactagtgt	caaaggcaat	tgtgcaattg	gaacagagga	taaagggtct	1020
ttagggtggag	gggctatata	ttctctaggg	accgttcttt	tgcaagggaa	tcacgggata	1080
acttgtgata	agaatgagtc	tgcttcgcaa	ggaggcgcca	tttttggcaa	aaattgtcag	1140
atctctgaca	acgaggggcc	agtggttttc	agagatagta	cagcttgctt	aggaggaggc	1200
gctattgcag	ctcaagaaat	tgtttctatt	cagaacaatc	aggctgggat	ttccttcgag	1260
ggaggtaagg	ctagtctcgg	aggaggatatt	gcgtgtggat	cttttctctc	cgcagggtgt	1320
gcttctgttt	tagggaccat	tgtattttcg	aagaatttag	gcgcgatttc	gttctctcgt	1380
actttatgta	cgacctcaga	tttaggacaa	atggagtacc	agggaggagg	agctctattt	1440
ggtgaaaata	tttctctttc	tgagaatgct	ggtgtgctca	cctttaaaga	caacattgtg	1500
aagacttttg	cttcgaatgg	gaaaattctg	ggaggaggag	cgatttttag	tactggtaag	1560
gtggaataa	ctaataattc	cgaagggaatt	tcttttacag	gaaatgcgag	agctccacaa	1620
gctcttccaa	ctcaagagga	gtttccttta	ttcagcaaaa	aagaaggcg	accactctct	1680
tcaggatatt	ctgggggagg	agcgatttta	ggaagagaag	tagctattct	ccacaacgct	1740
gcagtagtat	ttgagcaaaa	tcgtttgcag	tcagcggaag	aagaagcgac	attattaggt	1800
tgttgtggag	gaggcgctgt	tcattgggatg	gatagcactt	cgattgttgg	caactcttca	1860
gtaagatttg	gtaataatta	cgcaatggga	caaggagtct	caggaggagc	tcttttatct	1920
aaaacagtgc	agtttagctgg	gaatggaagc	gtcgattttt	ctcgaataat	tgctagtgtt	1980
ggaggaggag	ctcttcaagc	ttctgaagga	aattgtgagc	tagtgtataa	cggctatgtg	2040
ctattcagag	ataatcgagg	gagggtttat	gggggtgcta	tttcttgctt	acgtggagat	2100
gtagtcatatt	ctggaaacaa	gggtagagtt	gaattttaa	acaacatagc	aacacgtctt	2160
tatgtggaag	aaactgtaga	aaagggtgaa	gaggtagagc	cagctcctga	gcaaaaagac	2220
aataatgagc	tttctttctt	agggagagca	gaacagagtt	ttattactgc	agotaatcaa	2280
gctcttttct	catctgaaga	tggggattta	tcacctgagt	catccatttc	ttctgaagaa	2340
cttgcgaaaa	gaagagagtg	tgctggagga	gctatttttg	caaaacgggt	tcgtattgta	2400
gtaaaccaag	aggcggttgt	attctcgaat	aacttctctg	atatttatgg	cggcgccatt	2460
tttacagggt	ctcttcgaga	agaggataag	ttagatgggc	aaatccctga	agtottgatc	2520
tcaggcaatg	caggggatgt	tgttttttcc	ggaaattcct	cgaagcgtga	tgagcatctt	2580
cctcatacag	gtgggggagc	catttgtact	caaaatttga	cgattttotca	gaatacaggg	2640
aatgttctgt	tttataacaa	cgtggcctgt	tcgggaggag	ctgttcgtat	agaggatcat	2700
ggtaatgttc	ttttagaagc	ttttggagga	gatattgttt	ttaaaggaaa	ttcttctttc	2760
agagcacaa	gatccgatgc	tatctatttt	gcaggtaaa	aatcgcatat	tacagccctg	2820
aatgctacgg	aaggacatgc	tattgttttc	cacgacgcac	tagtttttga	aaatctagaa	2880

```

gaaaggaaat ctgctgaagt attgttaatc aatagtcgag aaaatccagg ttacactgga 2940
tctattcgat ttttagaagc agaaagttaa gtctctcaat gtattcatgt acaacaagga 3000
agccttgagt tgctaaatgg agccacatta tgtagttagt gttttaaaca agatgctgga 3060
gctaagttgg tattgggtgc tggagctaaa ctgaagattt tagattcagg aactcctgta 3120
caacaagggc atgctatcag taaacctgaa gcagaaatcg agtcacttc tgaaccagag 3180
ggtgcacatt ctctttggat tgcgaagaat gctcaaacaa cagttcctat ggttgatata 3240
catactatct ctgtagattt agcctccttc tcttctagtc aacaggaggg gacagtagaa 3300
gctoctcagg ttattgttcc tggaggaagt tatgttcgat ctggagagct taatttggag 3360
ttagttaaca caacaggtag tggttatgaa aatcatgctt tattgaagaa tgaggctaaa 3420
gttccattga tgtctttcgt tgcctotggg gatgaagctt cagccgaaat cagtaacttg 3480
tcggtttctg atttacagat tcatgtagta actccagaga ttgaagaaga cacatacggc 3540
catatgggag attggtctga ggctaaaatt caagatggaa ctcttgctat tagttggaat 3600
cctactggat atcgattaga tcctcaaaaa gcaggggctt tagtatttaa tgcattatgg 3660
gaagaagggg ctgtcttgct tgcctgaaa aatgcacgct ttgctcataa tctcactgct 3720
cagcgtatgg aattcgatta ttctacaaat gtgtggggat tcgcctttgg tggtttccga 3780
actctatctg cagagaatct ggttgctatt gatggataca aaggagctta tgggtggtct 3840
tctgctggag tcgatattca attgatggaa gattttgttc taggagttag tggagctgct 3900
ttcctaggtt aaatggatag tcagaagttt gatgcggagg tttctcggaa gggagttggt 3960
ggttctgtat atacaggatt tttagctgga tcctgggttct tcaaaggaca atatagcctt 4020
ggagaaacac agaacgatat gaaaacgcgt tatggagtac taggagagtc gagtgttctt 4080
tggacatctc gaggagtact ggcagatgct ttagttgaat accgaagttt agttggtcct 4140
gtgagaccta ctttttatgc tttgcatttc aatccttatg tcgaagtatc ttatgcttct 4200
atgaaattcc ctggctttac agaacaagga agagaagcgc gttcttttga agacgcttcc 4260
cttaccaata tcaccattcc tttagggatg aagtttgaat tggcgttcat aaaaggacag 4320
ttttcagagg tgaactcttt ggaataaagt tatgcatggg aagcttatcg aaaagtagaa 4380
ggaggcgcgg tgcagctttt ggaagctggg tttgattggg agggagctcc aatggatctt 4440
cctagacagg agctgcgtgt cgtctggaa aataatacgg aatggagttc ttacttcagc 4500
acagtcttag gattaacagc tttttgtgga ggatttactt ctacagatag taaactagga 4560
tatgaggcga atactggatt gcgattgac ttt 4593

```

<210> 84

<211> 1422

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 84

```

atgaaaatta ttacacacagc tatcgaattt gctccggtaa tcaaagccgg aggcctggga 60
gacgcgctat acggactagc aaaagcttta gccgctaata acacaacgga agtggttaac 120
cctttatacc ctaaatattt tactttgccc aaagaacaag atctttgctc gatccaaaaa 180
ttatcttatt tttttgctgg agagcaagaa ccaactgctt tctcctactt ttatgaagga 240
attaaagtaa ctctattcaa actcgacaca cagccagagt tattcgagaa tgcggaaaca 300
atctacacaa gcgatgatgc cttccgtttt tgcgcttttt ctgctgctgc gccctoctac 360
atccaaaaag aaggagccaa tatcgttcat ttacacgatt ggcatacagg attagttgct 420
ggactactca aacaacagcc ctgctctcaa ttacaaaaga ttgttcttac cctacataat 480
tttggttatc gaggtatata aacacgagaa atattagaag cctcctcttt gaatgaattt 540
tatatcagcc agtaccaact atttcgcgat ccacaaactt gtgtgttgct aaaaggagct 600
ttatactggt cagatttctg gactacggtt tctcctacat acgccaaaga aattcttgaa 660
gattattccg attacgaaat tcacgatgcc attactgcta gacaacatca tctccgctgg 720
attttaaatg gaatcgacac gacaatttgg gggcctgaaa cggatcccaa ttagcgaaa 780
aactacacta aagagctttt cgagaccctt tcaatttttt ttgaagctaa agccgagaa 840
aaaaaagcct tgtacgaaag attaggcctc tctttagaac actctccttg cgtgtgcatt 900
atctctagaa ttgctgagca gaaaggctct cactttatga aacaggccat tctccatgca 960
ctagaaaaag cttacacgct cattattata ggtacctgct acgggaatca attgcatgaa 1020
gaattttgca atcttcaaga atctccctg aattccctg atgtaaggat tcttttgact 1080
tatagtgatg tgctggcag acaaattttt gccgctgcag atatgatctg cattccttct 1140
atgtttgaa catgtggact cacacaaatg attggaatgc gttacgggac tgtaccgtta 1200
gtaagagcta caggaggact agcagatact gtagcaaatg gaatcaatgg attttccttc 1260
tttaattccg atgacttcta tgaattccga aacatgcttt cggaagcagt gacaacctac 1320
cgtaccaacc acgacaagtg gcaacatatt gtacgtgctt gtctagattt ttcttcagac 1380
ctagaaactg ccgccaataa atatttagaa atttataaac aa 1422

```

<210> 85
 <211> 1179
 <212> DNA
 <213> Chlamydia trachomatis serovar D

<400> 85
 atgaaaaaac tcttgaaatc ggtattagta tttgcccgtt tgagttctgc ttcttccttg 60
 caagctctgc ctgtggggaa tcctgctgaa ccaagcctta tgatcgacgg aattctgtgg 120
 gaagggttctg gcgagatcc ttgcgatcct tgcgccactt ggtgtgacgc tatcagcatg 180
 cgtgttggtt actacggaga ctttgttttc gaccgtgttt tgaaaactga tgtgaataaa 240
 gaatttcaga tgggtgccaa gcctacaact gatacaggca atagtgcagc tccatccact 300
 cttacagcaa gagagaatcc tgcttacggc cgacatatgc aggatgctga gatgtttaca 360
 aatgccgctt gcatggcatt gaatatattg gatogttttg atgtattctg tacattagga 420
 gccaccagtg gatattctta aggaaactct gcttctttca atttagttgg attgtttgga 480
 gataatgaaa atcaaaaaac ggtcaaagcg gagtctgtac caaatatgag ctttgatcaa 540
 tctgttggtt agttgtatag agatactact tttgcgtgga gcgtcggcgc tcgcgcagct 600
 ttgtgggaat gtggatgtgc aacttttagga gcttcattcc aatatgctca atctaaacct 660
 aaagtagaag aattaaacgt tctctgcaat gcagcagagt ttactattaa taaacctaaa 720
 gggtatgtag gtaaggagtt tcctcttgat cttacagcag gaacagatgc tgcgacagga 780
 actaaggatg cctctattga ttaccatgaa tggcaagcaa gtttagctct ctcttacaga 840
 ctgaatatgt tcaactcccta cattggagtt aaatgggtct gagcaagctt tgatgccgat 900
 acgattcgtg tagccagcc aaaatcagct acagctatct ttgatactac cacgcttaac 960
 ccaactattg ctggagctgg cgatgtgaaa actggcgcag agggtcagct cggagacaca 1020
 atgcaaactg tttccttgca attgaacaag atgaaatcta gaaaatcttg cggatttgca 1080
 gtaggaacaa ctattgtgga tgcagacaaa tacgcagtta cagttgagac tcgcttgatc 1140
 gatgagagag cagctcacgt aaatgcacaa ttccgcttc 1179

<210> 86
 <211> 585
 <212> DNA
 <213> Chlamydia trachomatis serovar D

<400> 86
 atgggatcac tagttggaag acaggtccg gatttttctg gtaaagccgt tgtttgtgga 60
 gaagagaaaag aaatctctct agcagacttt cgtggtaagt atgtagtgtt cttcttttat 120
 cctaaagatt ttacctatgt ttgtctaca gaattgcatg cttttcaaga tagattggta 180
 gattttgaaag agcgaggtgc agtcgtgctt ggttgctccg ttgacgacat tgagacacat 240
 ttctcgttggc tcgctgtagc gagaaatgca ggaggaatag agggaaacaga atatcctctg 300
 ttagcagacc cttcttttaa aatatcagaa gcttttggty ttttgaatcc tgaaggatcg 360
 ctgcctttaa gagcgacttt ctttatcgat aaatatgggg ttgttcgtca tgcggttatc 420
 aatgatcttc ctttagggcg ttccattgac gaggaattgc gtattttaga ttcattgatc 480
 ttctttgaga accacggaat ggtttgtcca gctaactggc gttctggaga gcgtggaatg 540
 gtgccttctg aagagggatt aaaagaatat ttccagacga tggat 585

<210> 87
 <211> 258
 <212> DNA
 <213> Chlamydia trachomatis serovar D

<400> 87
 atgagtcaaa ataagaactc tgctttcatg cagcctgtga acgtatocgc tgatttagct 60
 gccatcggtt gtgcaggacc tatgcctcgc acagagatca ttaagaaaaa gtgggattac 120
 attaagaaga atggccttca agatcctaca aacaaacgta atatcaatcc cgatgataaa 180
 ttggctaag tttttggaac tgaaaaacct atcgatatgt tccaaatgac aaaaatgggt 240
 tctcaacaca tcattaaa 258

<210> 88
 <211> 1182
 <212> DNA
 <213> Chlamydia trachomatis serovar D

<400> 88
 atgtcaaaag aaacttttca acgtaataag cctcatatca acatagggac cattggccac 60
 gttgaccatg gtaagactac gttgacagct gctattacgc gtgcgttgct tggagatggg 120
 ttggctgatt ttcgtgatta tagctctatt gacaacactc ctgaagaaaa agctcgcggt 180
 attacaatta acgcttccca cgttgagtac gaaacagcta atcgctacta cgetcacgtg 240
 gactgccctg gtcacgctga ctatgttaaa aacatgatca ccggtgcagc tcaaatggac 300
 ggggctattc tagtagtttc tgcaacagac ggagctatgc ctcaaaactaa agagcatatt 360
 cttttggcaa gacaagttgg ggttcottac atogttgttt ttctcaataa aattgacatg 420
 atttccgaag aagacgctga attggctgcac ttagttgaga tggagttggg tgagcttctt 480
 gaagagaaaag gatacaaagg gtgtccaatc atcagaggtt ctgctctgaa agctttggaa 540
 ggggatgctg catacataga gaaagttcga gagctaattgc aagccgtcga tgataacatc 600
 cctactccag aaagagaaat tgacaagcct ttottaatgc ctattgagga cgtattctct 660
 atctccggac gaggaactgt agtaactgga cgtattgagc gtggaattgt taaagtttcc 720
 gataaagttc agttggtcgg tcttagagat actaaagaaa cgattgttac tgggggtgaa 780
 atgttcagaa aagaactccc agaaggtcgt gcaggagaga acgttggatt gctcctcaga 840
 ggtattggta agaacgatgt ggaaagagga atgggtgttt gcttgccaaa cagtgttaaa 900
 cctcatcacac agttcaagtg tgctgtttac gttttgcaa aagaagaagg tggacgacat 960
 aagcctttct tcacaggata tagacctcaa ttcttcttcc gtacaacaga cgtcacaggt 1020
 gtggtaactc tgctgaggg aattgagatg gtcatgcctg gggataacgt tgagtttgaa 1080
 gtgcaattga ttagccctgt ggcttttaga gaaggtatga gatttgcgat tcgtgaaggt 1140
 ggtcgtacaa tcgggtgctg aactatttct aagatcattg ca 1182

<210> 89

<211> 246

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 89
 atgggggcaag atcaccgaag aaaatttctt aagaaagtat cttttgtaaa aaaacaagca 60
 gcttttgcgg gtaactttat cgaagaaatt aagaagattg agtgggtaaa taagcgagat 120
 cttaaaagat acgtcaagat tgttttgatg aatatttttg gctttggatt ttccatctat 180
 tgtgtggatt tagctcttcg aaagtccctt tcattgttcg gtaaaagtaac aagctttttc 240
 tttggt 246

<210> 90

<211> 1137

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 90
 atgggtgatcc ctaagggtgga tctaggagaa agtgccgtca tgatgggtta caagcttact 60
 tcgcaacttg ctatgctttc gatcttattg actttcacc atactatggg tcatgcaagt 120
 cagatgagcc aaactcttcc tactattata gaagcacaag cggaagaggc attgcaggct 180
 gacaggggag ttgctggaca ggctottaaa aaacttcgta aaaaaagatg tgcttctaga 240
 aaatctgcat gtaaggcttc ttttaagaaa aaggatttct tttcttgat tacaatgga 300
 ttgttctctg gaaatcatga gcagcgttta actgcgaaaa aagagaacaa ggctcgaggt 360
 aaagagcctc gagtagtggt tcaaacgact aaaaaacgac aaataactca gtctgagaaa 420
 gaatttttgc attggctatg taatagtaaa agagaaagaa agcttctcaa gaaaaagcct 480
 gtaaatactt ctcttgctaa gagtgaagaa ttgagtccta aagaagcagc aatagctgct 540
 gctcgagctt ctctttctcc agaagaaaaa cgtcaattga ttctgtagt gttagcagaa 600
 gaaaagactg ctctgaaatc tgggcgtgcg gcttgtgcgg taagtggagaa tottaaaaga 660
 gacggaagta ttacttctac attgcgctat gatgcggaga aagctttgac tacacgtgta 720
 aaacgcaatg aaaattctgt aaatgctaga gcaagacaac gagccgctct tcaaaaagcc 780
 aagaaagcaa agacggagaa acctgaggct gatgagaaag ctgcagaagc tgttgccgca 840
 gctccaacca aacaggcgca taaggagcca gagaattact tcgcagctac agcttctaca 900
 aataatacta atgttatgtc ctatctaaat gctcatcaat accgttgtga ttcttcggag 960
 acggactggc cttgctcttc ttgtgttacg aaacgccgag ctacttctcg tatttctgtg 1020
 tgtactatgg tggttaccgt cattgctatg atcgtaggag ctgttatcat ttctaagtct 1080
 acagactcta ccgttgccgg ctcctcggga acaggaggag gaggctcaac gcaacca 1137

<210> 91

<211> 1689

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 91

```

atggtttatt ttagagctca tcaacctagg catacgccca aaacatttcc tttggaagtt 60
caccattcgt tctccgataa gcatcctcaa attgctaaag ctatgcggat tacggggata 120
gccctcgag ctctatctct gctcgctgta gtgcgctgct ttattgcogt cctctcgagg 180
ggagctgcca ttctctttgc tgctattagt ggaattgctg taatgtctgg cctcttatcc 240
gctgccacca ttatctgttc tgcaaaaaag gctttggctc aacgaaaaca aaaacaacta 300
gaagagtcgc ttccgttaga taatgogacc gagcatgtga gttacctgac ctccagacacc 360
tcttatttta atcaatggga atccttaggt gctctaaata agcagttgtc tcagattgac 420
ttaactattc aagctcccg aaaaaaacta ttaaaagaag ttcttggttc cagatacgat 480
tccattaatc actccatcga agagatctcc gatcgcttta cgaatgct ctctcttctt 540
cgattaagag aacattttta tcgaggagaa gagcgttatg cccctattt aagccctcct 600
ctaactaaca agaatcggtt gctgacccaa atcacatcca atatgattag gatgctacca 660
aaatccggtg gtgttttttc cctcaaagcc aatacactaa gtcattgccag ccgcacacta 720
tatacagtat taaaagtgc tttatcctta ggagttctcg ctggagtgc tgctcttctc 780
atctttcttc cccctagcct gccttttctc gctgttatag gagtatcttc cttagcattg 840
gggatggcat ctttctctat gattcggggc attaagtatt tgctcgaaca ttctctctg 900
aatagaaagc aactagctaa agatattcaa aaaccattg gccagatgt cttggcctct 960
atggttcatt accagcatca attactatca catctacatg aaactctatt agatgaagcc 1020
atcacagcta gatggagcga gcccttcttt attgaacacg ctaactctaa ggcaaaaatt 1080
gaagatttga caaaacaata tgatatattg aacgcagcct ttaataaatc tttacaacaa 1140
gatgaggcgc tccgttctca attagagaaa cgagcttact tattcccaat tcctaataac 1200
gacgaaaatg ctaaaaacta agaatcgcag cttctagact cagaaaatga ttcaaattct 1260
gaatttcagg agattataaa taaaggacta gaagctgcca ataaacgacg agctgacgct 1320
aagtcaaaat tctatacgg aagcgaacc tctgacaaaa tattctctat atggaaaccc 1380
acaaagaact tggcattaga agatttgttg agagtgcatt aagcttgcaa tgaagagcaa 1440
caagctctcc tcttgaaga ttatatgagt tataaaacot cagaatgtca agctgcactc 1500
caaaaagtga gtcaagaact gaaggcggca caaaaatcat tcgcagtcct agaaaagcat 1560
gctctagaca gatcttatga atccagtgt gccacgatgg atttagctag agcgaatcaa 1620
gaaacacacc ggctcttgaa catcctctct gaattacaac aactagcaca atacctgtta 1680
gataatcac                                     1689

```

<210> 92

<211> 1074

<212> DNA

<213> Chlamydia trachomatis serovar D

<400> 92

```

gtgcgtaaaa ctgtcattgt tgctatgtct ggaggagtgg attcctcggg tgttgcttat 60
ctcttaaaaga agcaagggga gtataatgtt gttgggctct tcatgaaaaa ttggggagag 120
caggacgaga atggtgagtg tactgcaacc aaagattttc gcgatgtaga gcggatcgca 180
gaacaattgt ccattccata ttacacagtt tcccttttota aggaatataa agagcgagtg 240
ttttctagat ttctaagaga atatgcgaac ggctacactc ccaatcctga tgtgttatgc 300
aatcgagaaa tcaaatttga tttattacag aagaaggtag gtgagctaaa aggtgatttt 360
ttagccacgg gacattattg tcgaggagg gctgatggaa ctggtttgtc cagagggaata 420
gacccaata aagaccaaag ttatttctta tgtgacactc ctaaggatgc tttatccaat 480
gtacttttcc cctgggagg tatgtataaa acggaggtag gtcgaattgc tcaagaagct 540
ggttttagcta ccgccacaaa aaaagatagc acagggattt gcttcattgg taaacggcct 600
tttaagagtt tcttgagca gttttagtag gactctcctg gagacattat tgattttgat 660
acacaacagg tagtcggccg acatgaagga tggcattatt atacgattgg acagcgtcga 720
gggttaaaaca taggaggaat ggaaaagcct tggtatgttc ttagcaagaa tatggaaaag 780
aatattgttt acattgtaag ggggtgaagat catcctttac tttatcgaca agagctttta 840
gctaagggaac ttaattggtt tgttcccttg caggagccta tgatctgtag tgctaaagtt 900
cggtacagat cccctgacga gaaatgttct gtatatcctt tggaagatgg aacggtaaaa 960
gtgattttcg atgtccctgt gaaagctgtc acccctggac agactgtagc tttctaccag 1020
ggggacattt gtttaggagg aggagtgtt gaagtgccta tgattcatca gctg 1074

```

<210> 93

<211> 801

<212> DNA

<213> *Chlamydia trachomatis* serovar D

<400> 93

```

atgtccagaa aaccggcttc taactcatcc cggaacacca aacggtcctc agacacttcc 60
tggaagatca ttgccaaga ttataataaa gccgttgatc gcgatggaca tttctatcat 120
aaggaagtga ttctccctaa tctcctttct aagctacata tttcccgctc atcgtctctg 180
gttgatgtag gatgtggtca agggattttg gagaagcatt tacccaaaca tctcccttat 240
ctaggaatcg atctttcccc tagtctgctg cgttttgcaa agaaaagcgc ttccctcaaaa 300
tcacgtcgct ttcttcatca cgatatgacg caaccggtag cagcagatca tcatgagcag 360
ttttcccatg ctacagcaat cctttctctt cagaatatgg aatctccaga acaagctatc 420
gcacacacag cgaatctttt ggctcctcaa ggtaggttgt ttattgttct caaccatcca 480
tgctttcgca tccctaggct ttcttcatgg ctttatgatg agcctaaaaa actcttatct 540
agaaaaaatag accgctatct ctctcctgtg gcggttccta togtttgtgca tcctggagaa 600
aaacattctg agacgacata ttctttccat ttccctttaa gctattgggt acaagcttta 660
tctaatacaca atcttctgat tgatagtatg gaagaatgga tctcccttaa aaaatcctca 720
gggaagaggg ctcgagcaga aaatctttgt cgcaaggagt ttccgctttt cttgtttatc 780
tcagcattaa aaatatcaaa a                                     801

```

<210> 94

<211> 2601

<212> DNA

<213> *Chlamydia trachomatis* serovar D

<400> 94

```

atggagaaat tttcagatgc agtaagcgaa gccttagaaa aggcgtttga gttagctaaa 60
aactctaagc attcctaogt gacagaaaaac catthtctga aaagtctttt gcaaaatcca 120
ggttccctat tttgttttgt cattaaggat gtgcacggta atcttggttt gcttacttct 180
ggtgtggacg acgccttacg cagagaacca ttgttagtgc agggaaacgc tgttgctagt 240
ccttctccaa gtttacagca gttgttgctc aatgcgcctc aagaagctag aagtatgggt 300
gacgaatata tatcagggga tcatttgtta ctagcttttt ggcatcgac taaagagcct 360
tttgcttctt ggagaaaaac tgtaaaaact acctctgaag cgttgaaaga attaattact 420
aaattaagac aaggaagtgc tatggactca cctagtgtgc aagaaaaatct gaaaggatta 480
gagaaatact gcaaaaattt gactgtactt gcaagagaag gcaagcttga tcctgtgatt 540
ggctcgagatg aagagattag acgtacgata caggttcttt ctacagcaac aaagaataat 600
cctatgttga taggggagcc cggagtggg aaaacagcaa tcgctgaagg acttgctctt 660
cgcatagtgc aaggggatgt tccagagagt ttaaaggaaa agcatctgta tgtactggat 720
atgggagctt tgattgcagg tgccaagtat cgaggagagt ttgaagagcg gttaaaaagt 780
gtattgaagg gtgtagaagc ttctgaaggc gagtgtatcc tattcattga tgaagtgcatt 840
acttttagtag gagcgggagc tacagatgga gctatggatg cagcgaatct attaaagcct 900
gcttttagcac gaggcacttt gcattgtatt ggcgctacga ctttgaatga ataccaaaaa 960
tatatagaga aagacgcggc ttgtgaacgg cgtttccagc ctatttttgt aacagaacct 1020
tctttggaag atgctgtatt cattctccgg gggtaaaggg aaaaatatga aatttttcat 1080
gggtgtgcga ttacagaagg ggctttgaat gcagctgtag ttctttctta tcgttacatc 1140
acagaccgat ttcttctctg taaggcgatt gacctaatg atgaggctgc gattttaatc 1200
cgtatgcaaa taggaagttt acctctgcct attgatgaaa aggaagaga attatcagct 1260
ttaatcgtga aacaagaagc tattaacgc gagcaagcac cagcttatca ggaagaggct 1320
gaagacatgc aaaaagcaat tgaccgggtt aaggaagagc tggccgcttt acgcttgcc 1380
tggtatgaag aaaaaggatt aattacagga ttaaagaaa agaagaatgc tttagaaaat 1440
ttaaaatttg ccgaagagga agctgagcgt actgccgatt acaatcgggt gccagaacta 1500
cgctatagtt tgattccttc ttgtgaggaa gaaattcatt tagctgagga agctttaaat 1560
caaagagatg ggccctgct tcgaaggaa gttgatgagc ggttgattgc gcaagttgtt 1620
gcgaattgga ctggaatccc tgtgcaaaaa atgttgagg gagaatctga aaagttattg 1680
gtgttgagg agtctttaga agaaagggtt gttggacaac ctttcgctat tgccgcagtc 1740
agtgattcga ttcgagctgc tcgagtagga ttgagtgtc cgcagcgtcc tctaggagtg 1800
tttctatttc ttggacctac aggggtaggg aaaactgagc ttgctaaagc attagcagag 1860
cttttattta ataagggaaga agcgatgatt cggtttgaca tgaccgaata tatggaaaaa 1920
cattccgctt ccaaatgat aggatctcct ccagggtatg taggatatga agaaggaggg 1980
agtctctcag aagctttaag aagacgacct tattctgttg ttctttttga tgagatagaa 2040
aaagcagata aagaagtatt taatatttta ttgcagattt ttgatgatgg gattcttacg 2100

```

```

gatagcaaga agcgtaagggt aaattgtaag aatgctcttt tcattatgac atcaaataatt 2160
ggttcgcaag agcttgctga ttattgtact aagaaaggaa ctatcgtaga caaagaagct 2220
gtgctatctg ttgttgcccc tgcgcttaaa aattatttta gtccagaatt tatcaatcgt 2280
atcgatgaca ttctgccttt cgttcctttg actacggaag acattgtaaa aattgtcggg 2340
attcaaataga atcgggttgc tttacgtttg ctggaaagaa aaatttcggt aacttgggat 2400
gattcttttag tgctatttct cagtgagcaa ggttatgaca gcgcttttgg agctcgccct 2460
ctgaagcgtt tgatacagca aaaagtagtg actatgttgt ctaaagctct tttgaaagga 2520
gatatcaaac ctggaatggc ggtggagctt actatggcaa aagatgtagt tgtgtttaaa 2580
attaaaacaa atccagctgt g 2601

```

<210> 95

<211> 1016

<212> PRT

<213> *Chlamydia trachomatis* serovar D

<400> 95

```

Met Pro Phe Ser Leu Arg Ser Thr Ser Phe Cys Phe Leu Ala Cys Leu
          5          10          15
Cys Ser Tyr Ser Tyr Gly Phe Ala Ser Ser Pro Gln Val Leu Thr Pro
          20          25          30
Asn Val Thr Thr Pro Phe Lys Gly Asp Asp Val Tyr Leu Asn Gly Asp
          35          40          45
Cys Ala Phe Val Asn Val Tyr Ala Gly Ala Glu Asn Gly Ser Ile Ile
          50          55          60
Ser Ala Asn Gly Asp Asn Leu Thr Ile Thr Gly Gln Asn His Thr Leu
          65          70          75          80
Ser Phe Thr Asp Ser Gln Gly Pro Val Leu Gln Asn Tyr Ala Phe Ile
          85          90          95
Ser Ala Gly Glu Thr Leu Thr Leu Lys Asp Phe Ser Ser Leu Met Phe
          100          105          110
Ser Lys Asn Val Ser Cys Gly Glu Lys Gly Met Ile Ser Gly Lys Thr
          115          120          125
Val Ser Ile Ser Gly Ala Gly Glu Val Ile Phe Trp Asp Asn Ser Val
          130          135          140
Gly Tyr Ser Pro Leu Ser Ile Val Pro Ala Ser Thr Pro Thr Pro Pro
          145          150          155          160
Ala Pro Ala Pro Ala Pro Ala Ala Ser Ser Ser Leu Ser Pro Thr Val
          165          170          175
Ser Asp Ala Arg Lys Gly Ser Ile Phe Ser Val Glu Thr Ser Leu Glu
          180          185          190
Ile Ser Gly Val Lys Lys Gly Val Met Phe Asp Asn Asn Ala Gly Asn
          195          200          205
Phe Gly Thr Val Phe Arg Gly Asn Ser Asn Asn Asn Ala Gly Ser Gly
          210          215          220
Gly Ser Gly Ser Ala Thr Thr Pro Ser Phe Thr Val Lys Asn Cys Lys
          225          230          235          240
Gly Lys Val Ser Phe Thr Asp Asn Val Ala Ser Cys Gly Gly Gly Val
          245          250          255
Val Tyr Lys Gly Thr Val Leu Phe Lys Asp Asn Glu Gly Gly Ile Phe
          260          265          270
Phe Arg Gly Asn Thr Ala Tyr Asp Asp Leu Gly Ile Leu Ala Ala Thr
          275          280          285
Ser Arg Asp Gln Asn Thr Glu Thr Gly Gly Gly Gly Gly Val Ile Cys
          290          295          300
Ser Pro Asp Asp Ser Val Lys Phe Glu Gly Asn Lys Gly Ser Ile Val
          305          310          315          320
Phe Asp Tyr Asn Phe Ala Lys Gly Arg Gly Gly Ser Ile Leu Thr Lys
          325          330          335
Glu Phe Ser Leu Val Ala Asp Asp Ser Val Val Phe Ser Asn Asn Thr
          340          345          350
Ala Glu Lys Gly Gly Gly Ala Ile Tyr Ala Pro Thr Ile Asp Ile Ser

```

355	360	365
Thr Asn Gly Gly Ser Ile	Leu Phe Glu Arg Asn Arg	Ala Ala Glu Gly
370	375	380
Gly Ala Ile Cys Val Ser	Glu Ala Ser Ser Gly Ser	Thr Gly Asn Leu
385	390	395
Thr Leu Ser Ala Ser Asp	Gly Asp Ile Val Phe Ser	Gly Asn Met Thr
405	410	415
Ser Asp Arg Pro Gly Glu	Arg Ser Ala Ala Arg	Ile Leu Ser Asp Gly
420	425	430
Thr Thr Val Ser Leu Asn	Ala Ser Gly Leu Ser	Lys Leu Ile Phe Tyr
435	440	445
Asp Pro Val Val Gln Asn	Asn Ser Ala Ala Gly	Ala Ser Thr Pro Ser
450	455	460
Pro Ser Ser Ser Ser Met	Pro Gly Ala Val Thr	Ile Asn Gln Ser Gly
465	470	475
Asn Gly Ser Val Ile Phe	Thr Ala Glu Ser Leu	Thr Pro Ser Glu Lys
485	490	495
Leu Gln Val Leu Asn Ser	Thr Ser Asn Phe Pro	Gly Ala Leu Thr Val
500	505	510
Ser Gly Gly Glu Leu Val	Val Thr Glu Gly Ala	Thr Leu Thr Thr Gly
515	520	525
Thr Ile Thr Ala Thr Ser	Gly Arg Val Thr Leu	Gly Ser Gly Ala Ser
530	535	540
Leu Ser Ala Val Ala Gly	Ala Ala Asn Asn Asn	Tyr Thr Cys Thr Val
545	550	555
Ser Lys Leu Gly Ile Asp	Leu Glu Ser Phe Leu	Thr Pro Asn Tyr Lys
565	570	575
Thr Ala Ile Leu Gly Ala	Asp Gly Thr Val Thr	Val Asn Ser Gly Ser
580	585	590
Thr Leu Asp Leu Val Met	Glu Ser Glu Ala Glu	Val Tyr Asp Asn Pro
595	600	605
Leu Phe Val Gly Ser Leu	Thr Ile Pro Phe Val	Thr Leu Ser Ser Ser
610	615	620
Ser Ala Ser Asn Gly Val	Thr Lys Asn Ser Val	Thr Ile Asn Asp Ala
625	630	635
Asp Ala Ala His Tyr Gly	Tyr Gln Gly Ser Trp	Ser Ala Asp Trp Thr
645	650	655
Lys Pro Pro Leu Ala Pro	Asp Ala Lys Gly Met	Val Pro Pro Asn Thr
660	665	670
Asn Asn Thr Leu Tyr Leu	Thr Trp Arg Pro Ala	Ser Asn Tyr Gly Glu
675	680	685
Tyr Arg Leu Asp Pro Gln	Arg Lys Gly Glu Leu	Val Pro Asn Ser Leu
690	695	700
Trp Val Ala Gly Ser Ala	Leu Arg Thr Phe Thr	Asn Gly Leu Lys Glu
705	710	715
His Tyr Val Ser Arg Asp	Val Gly Phe Val Ala	Ser Leu His Ala Leu
725	730	735
Gly Asp Tyr Ile Leu Asn	Tyr Thr Gln Asp Asp	Arg Asp Gly Phe Leu
740	745	750
Ala Arg Tyr Gly Gly Phe	Gln Ala Thr Ala Ala	Ser His Tyr Glu Asn
755	760	765
Gly Ser Ile Phe Gly Val	Ala Phe Gly Gln Leu	Tyr Gly Gln Thr Lys
770	775	780
Ser Arg Met Tyr Tyr Ser	Lys Asp Ala Gly Asn	Met Thr Met Leu Ser
785	790	795
Cys Phe Gly Arg Ser Tyr	Val Asp Ile Lys Gly	Thr Glu Thr Val Met
805	810	815
Tyr Trp Glu Thr Ala Tyr	Gly Tyr Ser Val His	Arg Met His Thr Gln
820	825	830
Tyr Phe Asn Asp Lys Thr	Gln Lys Phe Asp His	Ser Lys Cys His Trp
835	840	845

His Asn Asn Asn Tyr Tyr Ala Phe Val Gly Ala Glu His Asn Phe Leu
 850 855 860
 Glu Tyr Cys Ile Pro Thr Arg Gln Phe Ala Arg Asp Tyr Glu Leu Thr
 865 870 875 880
 Gly Phe Met Arg Phe Glu Met Ala Gly Gly Trp Ser Ser Ser Thr Arg
 885 890 895
 Glu Thr Gly Ser Leu Thr Arg Tyr Phe Ala Arg Gly Ser Gly His Asn
 900 905 910
 Met Ser Leu Pro Ile Gly Ile Val Ala His Ala Val Ser His Val Arg
 915 920 925
 Arg Ser Pro Pro Ser Lys Leu Thr Leu Asn Met Gly Tyr Arg Pro Asp
 930 935 940
 Ile Trp Arg Val Thr Pro His Cys Asn Met Glu Ile Ile Ala Asn Gly
 945 950 955 960
 Val Lys Thr Pro Ile Gln Gly Ser Pro Leu Ala Arg His Ala Phe Phe
 965 970 975
 Leu Glu Val His Asp Thr Leu Tyr Ile His His Phe Gly Arg Ala Tyr
 980 985 990
 Met Asn Tyr Ser Leu Asp Ala Arg Arg Gln Thr Ala His Phe Val
 995 1000 1005
 Ser Met Gly Leu Asn Arg Ile Phe
 1010 1015

<210> 96

<211> 346

<212> PRT

<213> *Chlamydia trachomatis* serovar D

<400> 96

Met Gln Ala Asp Ile Leu Asp Gly Lys Gln Lys Arg Val Asn Leu Asn
 5 10 15
 Ser Lys Arg Leu Val Asn Cys Asn Gln Val Asp Val Asn Gln Leu Val
 20 25 30
 Pro Ile Lys Tyr Lys Trp Ala Trp Glu His Tyr Leu Asn Gly Cys Ala
 35 40 45
 Asn Asn Trp Leu Pro Thr Glu Ile Pro Met Gly Lys Asp Ile Glu Leu
 50 55 60
 Trp Lys Ser Asp Arg Leu Ser Glu Asp Glu Arg Val Ile Leu Leu
 65 70 75 80
 Asn Leu Gly Phe Phe Ser Thr Ala Glu Ser Leu Val Gly Asn Asn Ile
 85 90 95
 Val Leu Ala Ile Phe Lys His Val Thr Asn Pro Glu Ala Arg Gln Tyr
 100 105 110
 Leu Leu Arg Gln Ala Phe Glu Glu Ala Val His Thr His Thr Phe Leu
 115 120 125
 Tyr Ile Cys Glu Ser Leu Gly Leu Asp Glu Lys Glu Ile Phe Asn Ala
 130 135 140
 Tyr Asn Glu Arg Ala Ala Ile Lys Ala Lys Asp Asp Phe Gln Met Glu
 145 150 155 160
 Ile Thr Gly Lys Val Leu Asp Pro Asn Phe Arg Thr Asp Ser Val Glu
 165 170 175
 Gly Leu Gln Glu Phe Val Lys Asn Leu Val Gly Tyr Tyr Ile Ile Met
 180 185 190
 Glu Gly Ile Phe Phe Tyr Ser Gly Phe Val Met Ile Leu Ser Phe His
 195 200 205
 Arg Gln Asn Lys Met Ile Gly Ile Gly Glu Gln Tyr Gln Tyr Ile Leu
 210 215 220
 Arg Asp Glu Thr Ile His Leu Asn Phe Gly Ile Asp Leu Ile Asn Gly
 225 230 235 240
 Ile Lys Glu Glu Asn Pro Glu Ile Trp Thr Pro Glu Leu Gln Gln Glu

64

```

      245      250      255
Ile Val Glu Leu Ile Lys Arg Ala Val Asp Leu Glu Ile Glu Tyr Ala
      260      265      270
Gln Asp Cys Leu Pro Arg Gly Ile Leu Gly Leu Arg Ala Ser Met Phe
      275      280      285
Ile Asp Tyr Val Gln His Ile Ala Asp Arg Arg Leu Glu Arg Ile Gly
      290      295      300
Leu Lys Pro Ile Tyr His Thr Lys Asn Pro Phe Pro Trp Met Ser Glu
305      310      315      320
Thr Ile Asp Leu Asn Lys Glu Lys Asn Phe Phe Glu Thr Arg Val Ile
      325      330      335
Glu Tyr Gln His Ala Ala Ser Leu Thr Trp
      340      345

```

<210> 97

<211> 1053

<212> PRT

<213> *Chlamydia trachomatis* serovar D

<400> 97

```

Met Phe Thr Arg Ile Val Met Val Asp Leu Gln Glu Lys Gln Cys Thr
      5      10      15
Ile Val Lys Arg Asn Gly Met Phe Val Pro Phe Asp Arg Asn Arg Ile
      20      25      30
Phe Gln Ala Leu Glu Ala Ala Phe Arg Asp Thr Arg Arg Ile Asp Asp
      35      40      45
His Met Pro Leu Pro Glu Asp Leu Glu Ser Ser Ile Arg Ser Ile Thr
      50      55      60
His Gln Val Val Lys Glu Val Val Gln Lys Ile Thr Asp Gly Gln Val
      65      70      75      80
Val Thr Val Glu Arg Ile Gln Asp Met Val Glu Ser Gln Leu Tyr Val
      85      90      95
Asn Gly Leu Gln Asp Val Ala Arg Asp Tyr Ile Val Tyr Arg Asp Asp
      100      105      110
Arg Lys Ala His Arg Lys Lys Ser Trp Gln Ser Leu Ser Val Val Arg
      115      120      125
Arg Cys Gly Thr Val Val His Phe Asn Pro Met Lys Ile Ser Ala Ala
      130      135      140
Leu Glu Lys Ala Phe Arg Ala Thr Asp Lys Thr Glu Gly Met Thr Pro
      145      150      155      160
Ser Ser Val Arg Glu Glu Ile Asn Ala Leu Thr Gln Asn Ile Val Ala
      165      170      175
Glu Ile Glu Glu Cys Cys Pro Gln Gln Asp Arg Arg Ile Asp Ile Glu
      180      185      190
Lys Ile Gln Asp Ile Val Glu Gln Gln Leu Met Val Val Gly His Tyr
      195      200      205
Ala Val Ala Lys Asn Tyr Ile Leu Tyr Arg Glu Ala Arg Ala Arg Val
      210      215      220
Arg Asp Asn Arg Glu Glu Asp Gly Ser Thr Glu Lys Thr Ile Ala Glu
      225      230      235      240
Glu Ala Val Glu Val Leu Ser Lys Asp Gly Ser Thr Tyr Thr Met Thr
      245      250      255
His Ser Gln Leu Leu Ala His Leu Ala Arg Ala Cys Ser Arg Phe Pro
      260      265      270
Glu Thr Thr Asp Ala Ala Leu Leu Thr Asp Met Ala Phe Ala Asn Phe
      275      280      285
Tyr Ser Gly Ile Lys Glu Ser Glu Val Val Leu Ala Cys Ile Met Ala
      290      295      300
Ala Arg Ala Asn Ile Glu Lys Glu Pro Asp Tyr Ala Phe Val Ala Ala
305      310      315      320

```

Glu Leu Leu Leu Asp Val Val Tyr Lys Glu Ala Leu Gly Lys Ser Lys
 325 330 335
 Tyr Ala Glu Asp Leu Glu Gln Ala His Arg Asp His Phe Lys Arg Tyr
 340 345 350
 Ile Ala Glu Gly Asp Thr Tyr Arg Leu Asn Ala Glu Leu Lys His Leu
 355 360 365
 Phe Asp Leu Asp Ala Leu Ala Asp Ala Met Asp Leu Ser Arg Asp Leu
 370 375 380
 Gln Phe Ser Tyr Met Gly Ile Gln Asn Leu Tyr Asp Arg Tyr Phe Asn
 385 390 395 400
 His His Glu Gly Cys Arg Leu Glu Thr Pro Gln Ile Phe Trp Met Arg
 405 410 415
 Val Ala Met Gly Leu Ala Leu Asn Glu Gln Asp Lys Thr Ser Trp Ala
 420 425 430
 Ile Thr Phe Tyr Asn Leu Leu Ser Thr Phe Arg Tyr Thr Pro Ala Thr
 435 440 445
 Pro Thr Leu Phe Asn Ser Gly Met Arg His Ser Gln Leu Ser Ser Cys
 450 455 460
 Tyr Leu Ser Thr Val Gln Asp Asn Leu Val Asn Ile Tyr Lys Val Ile
 465 470 475 480
 Ala Asp Asn Ala Met Leu Ser Lys Trp Ala Gly Gly Ile Gly Asn Asp
 485 490 495
 Trp Thr Ala Ile Arg Ala Thr Gly Ala Leu Ile Lys Gly Thr Asn Gly
 500 505 510
 Arg Ser Gln Gly Val Ile Pro Phe Ile Lys Val Thr Asn Asp Thr Ala
 515 520 525
 Val Ala Val Asn Gln Gly Gly Lys Arg Lys Gly Ala Val Cys Val Tyr
 530 535 540
 Leu Glu Val Trp His Leu Asp Tyr Glu Asp Phe Leu Glu Leu Arg Lys
 545 550 555 560
 Asn Thr Gly Asp Glu Arg Arg Arg Ala His Asp Val Asn Ile Ala Ser
 565 570 575
 Trp Ile Pro Asp Leu Phe Phe Lys Arg Leu Gln Gln Lys Gly Thr Trp
 580 585 590
 Thr Leu Phe Ser Pro Asp Asp Val Pro Gly Leu His Asp Ala Tyr Gly
 595 600 605
 Glu Glu Phe Glu Arg Leu Tyr Glu Glu Tyr Glu Arg Lys Val Asp Thr
 610 615 620
 Gly Glu Ile Arg Leu Phe Lys Lys Val Glu Ala Glu Asp Leu Trp Arg
 625 630 635 640
 Lys Met Leu Ser Met Leu Phe Glu Thr Gly His Pro Trp Met Thr Phe
 645 650 655
 Lys Asp Pro Ser Asn Ile Arg Ser Ala Gln Asp His Lys Gly Val Val
 660 665 670
 Arg Cys Ser Asn Leu Cys Thr Glu Ile Leu Leu Asn Cys Ser Glu Thr
 675 680 685
 Glu Thr Ala Val Cys Asn Leu Gly Ser Ile Asn Leu Val Gln His Ile
 690 695 700
 Val Gly Asp Gly Leu Asp Glu Glu Lys Leu Ser Glu Thr Ile Ser Ile
 705 710 715 720
 Ala Val Arg Met Leu Asp Asn Val Ile Asp Ile Asn Phe Tyr Pro Thr
 725 730 735
 Lys Glu Ala Lys Glu Ala Asn Phe Ala His Arg Ala Ile Gly Leu Gly
 740 745 750
 Val Met Gly Phe Gln Asp Ala Leu Tyr Lys Leu Asp Ile Ser Tyr Ala
 755 760 765
 Ser Gln Glu Ala Val Glu Phe Ala Asp Tyr Ser Ser Glu Leu Ile Ser
 770 775 780
 Tyr Tyr Ala Ile Gln Ala Ser Cys Leu Leu Ala Lys Glu Arg Gly Thr
 785 790 795 800
 Tyr Ser Ser Tyr Lys Gly Ser Lys Trp Asp Arg Gly Leu Leu Pro Ile

66

```

      805      810      815
Asp Thr Ile Gln Leu Leu Ala Asn Tyr Arg Gly Glu Ala Asn Leu Gln
      820      825      830
Met Asp Thr Ser Ser Arg Lys Asp Trp Glu Pro Ile Arg Ser Leu Val
      835      840      845
Lys Glu His Gly Met Arg His Cys Gln Leu Met Ala Ile Ala Pro Thr
      850      855      860
Ala Thr Ile Ser Asn Ile Ile Gly Val Thr Gln Ser Ile Glu Pro Thr
      865      870      875      880
Tyr Lys His Leu Phe Val Lys Ser Asn Leu Ser Gly Glu Phe Thr Ile
      885      890      895
Pro Asn Val Tyr Leu Ile Glu Lys Leu Lys Lys Leu Gly Ile Trp Asp
      900      905      910
Ala Asp Met Leu Asp Asp Leu Lys Tyr Phe Asp Gly Ser Leu Leu Glu
      915      920      925
Ile Glu Arg Ile Pro Asp His Leu Lys His Ile Phe Leu Thr Ala Phe
      930      935      940
Glu Ile Glu Pro Glu Trp Ile Ile Glu Cys Ala Ser Arg Arg Gln Lys
      945      950      955      960
Trp Ile Asp Met Gly Gln Ser Leu Asn Leu Tyr Leu Ala Gln Pro Asp
      965      970      975
Gly Lys Lys Leu Ser Asn Met Tyr Leu Thr Ala Trp Lys Lys Gly Leu
      980      985      990
Lys Thr Thr Tyr Tyr Leu Arg Ser Ser Ala Thr Thr Val Glu Lys
      995      1000      1005
Ser Phe Val Asp Ile Asn Lys Arg Gly Ile Gln Pro Arg Trp Met Lys
      1010      1015      1020
Asn Lys Ser Ala Ser Ala Gly Ile Ile Val Glu Arg Ala Lys Lys Ala
      1025      1030      1035      1040
Pro Val Cys Ser Leu Glu Glu Gly Cys Glu Ala Cys Gln
      1045      1050

```

<210> 98

<211> 1531

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 98

```

Met Ser Ser Glu Lys Asp Ile Lys Ser Thr Cys Ser Lys Phe Ser Leu
      5      10      15
Ser Val Val Ala Ala Ile Leu Ala Ser Val Ser Gly Leu Ala Ser Cys
      20      25      30
Val Asp Leu His Ala Gly Gly Gln Ser Val Asn Glu Leu Val Tyr Val
      35      40      45
Gly Pro Gln Ala Val Leu Leu Leu Asp Gln Ile Arg Asp Leu Phe Val
      50      55      60
Gly Ser Lys Asp Ser Gln Ala Glu Gly Gln Tyr Arg Leu Ile Val Gly
      65      70      75      80
Asp Pro Ser Ser Phe Gln Glu Lys Asp Ala Asp Thr Leu Pro Gly Lys
      85      90      95
Val Glu Gln Ser Thr Leu Phe Ser Val Thr Asn Pro Val Val Phe Gln
      100      105      110
Gly Val Asp Gln Gln Asp Gln Val Ser Ser Gln Gly Leu Ile Cys Ser
      115      120      125
Phe Thr Ser Ser Asn Leu Asp Ser Pro Arg Asp Gly Glu Ser Phe Leu
      130      135      140
Gly Ile Ala Phe Val Gly Asp Ser Ser Lys Ala Gly Ile Thr Leu Thr
      145      150      155      160
Asp Val Lys Ala Ser Leu Ser Gly Ala Ala Leu Tyr Ser Thr Glu Asp
      165      170      175

```

Leu Ile Phe Glu Lys Ile Lys Gly Gly Leu Glu Phe Ala Ser Cys Ser
 180 185 190
 Ser Leu Glu Gln Gly Gly Ala Cys Ala Ala Gln Ser Ile Leu Ile His
 195 200 205
 Asp Cys Gln Gly Leu Gln Val Lys His Cys Thr Thr Ala Val Asn Ala
 210 215 220
 Glu Gly Ser Ser Ala Asn Asp His Leu Gly Phe Gly Gly Gly Ala Phe
 225 230 235 240
 Phe Val Thr Gly Ser Leu Ser Gly Glu Lys Ser Leu Tyr Met Pro Ala
 245 250 255
 Gly Asp Met Val Val Ala Asn Cys Asp Gly Ala Ile Ser Phe Glu Gly
 260 265 270
 Asn Ser Ala Asn Phe Ala Asn Gly Gly Ala Ile Ala Ala Ser Gly Lys
 275 280 285
 Val Leu Phe Val Ala Asn Asp Lys Lys Thr Ser Phe Ile Glu Asn Arg
 290 295 300
 Ala Leu Ser Gly Gly Ala Ile Ala Ala Ser Ser Asp Ile Ala Phe Gln
 305 310 315 320
 Asn Cys Ala Glu Leu Val Phe Lys Gly Asn Cys Ala Ile Gly Thr Glu
 325 330 335
 Asp Lys Gly Ser Leu Gly Gly Gly Ala Ile Ser Ser Leu Gly Thr Val
 340 345 350
 Leu Leu Gln Gly Asn His Gly Ile Thr Cys Asp Lys Asn Glu Ser Ala
 355 360 365
 Ser Gln Gly Gly Ala Ile Phe Gly Lys Asn Cys Gln Ile Ser Asp Asn
 370 375 380
 Glu Gly Pro Val Val Phe Arg Asp Ser Thr Ala Cys Leu Gly Gly Gly
 385 390 395 400
 Ala Ile Ala Ala Gln Glu Ile Val Ser Ile Gln Asn Asn Gln Ala Gly
 405 410 415
 Ile Ser Phe Glu Gly Gly Lys Ala Ser Phe Gly Gly Gly Ile Ala Cys
 420 425 430
 Gly Ser Phe Ser Ser Ala Gly Gly Ala Ser Val Leu Gly Thr Ile Asp
 435 440 445
 Ile Ser Lys Asn Leu Gly Ala Ile Ser Phe Ser Arg Thr Leu Cys Thr
 450 455 460
 Thr Ser Asp Leu Gly Gln Met Glu Tyr Gln Gly Gly Gly Ala Leu Phe
 465 470 475 480
 Gly Glu Asn Ile Ser Leu Ser Glu Asn Ala Gly Val Leu Thr Phe Lys
 485 490 495
 Asp Asn Ile Val Lys Thr Phe Ala Ser Asn Gly Lys Ile Leu Gly Gly
 500 505 510
 Gly Ala Ile Leu Ala Thr Gly Lys Val Glu Ile Thr Asn Asn Ser Glu
 515 520 525
 Gly Ile Ser Phe Thr Gly Asn Ala Arg Ala Pro Gln Ala Leu Pro Thr
 530 535 540
 Gln Glu Glu Phe Pro Leu Phe Ser Lys Lys Glu Gly Arg Pro Leu Ser
 545 550 555 560
 Ser Gly Tyr Ser Gly Gly Ala Ile Leu Gly Arg Glu Val Ala Ile
 565 570 575
 Leu His Asn Ala Ala Val Val Phe Glu Gln Asn Arg Leu Gln Cys Ser
 580 585 590
 Glu Glu Glu Ala Thr Leu Leu Gly Cys Cys Gly Gly Gly Ala Val His
 595 600 605
 Gly Met Asp Ser Thr Ser Ile Val Gly Asn Ser Ser Val Arg Phe Gly
 610 615 620
 Asn Asn Tyr Ala Met Gly Gln Gly Val Ser Gly Gly Ala Leu Leu Ser
 625 630 635 640
 Lys Thr Val Gln Leu Ala Gly Asn Gly Ser Val Asp Phe Ser Arg Asn
 645 650 655
 Ile Ala Ser Leu Gly Gly Gly Ala Leu Gln Ala Ser Glu Gly Asn Cys

[illegible]

Ala Ser Ala Glu Ile Ser Asn Leu Ser Val Ser Asp Leu Gln Ile His
 1155 1160 1165
 Val Val Thr Pro Glu Ile Glu Glu Asp Thr Tyr Gly His Met Gly Asp
 1170 1175 1180
 Trp Ser Glu Ala Lys Ile Gln Asp Gly Thr Leu Val Ile Ser Trp Asn
 1185 1190 1195 1200
 Pro Thr Gly Tyr Arg Leu Asp Pro Gln Lys Ala Gly Ala Leu Val Phe
 1205 1210 1215
 Asn Ala Leu Trp Glu Glu Gly Ala Val Leu Ser Ala Leu Lys Asn Ala
 1220 1225 1230
 Arg Phe Ala His Asn Leu Thr Ala Gln Arg Met Glu Phe Asp Tyr Ser
 1235 1240 1245
 Thr Asn Val Trp Gly Phe Ala Phe Gly Gly Phe Arg Thr Leu Ser Ala
 1250 1255 1260
 Glu Asn Leu Val Ala Ile Asp Gly Tyr Lys Gly Ala Tyr Gly Gly Ala
 1265 1270 1275 1280
 Ser Ala Gly Val Asp Ile Gln Leu Met Glu Asp Phe Val Leu Gly Val
 1285 1290 1295
 Ser Gly Ala Ala Phe Leu Gly Lys Met Asp Ser Gln Lys Phe Asp Ala
 1300 1305 1310
 Glu Val Ser Arg Lys Gly Val Val Gly Ser Val Tyr Thr Gly Phe Leu
 1315 1320 1325
 Ala Gly Ser Trp Phe Phe Lys Gly Gln Tyr Ser Leu Gly Glu Thr Gln
 1330 1335 1340
 Asn Asp Met Lys Thr Arg Tyr Gly Val Leu Gly Glu Ser Ser Ala Ser
 1345 1350 1355 1360
 Trp Thr Ser Arg Gly Val Leu Ala Asp Ala Leu Val Glu Tyr Arg Ser
 1365 1370 1375
 Leu Val Gly Pro Val Arg Pro Thr Phe Tyr Ala Leu His Phe Asn Pro
 1380 1385 1390
 Tyr Val Glu Val Ser Tyr Ala Ser Met Lys Phe Pro Gly Phe Thr Glu
 1395 1400 1405
 Gln Gly Arg Glu Ala Arg Ser Phe Glu Asp Ala Ser Leu Thr Asn Ile
 1410 1415 1420
 Thr Ile Pro Leu Gly Met Lys Phe Glu Leu Ala Phe Ile Lys Gly Gln
 1425 1430 1435 1440
 Phe Ser Glu Val Asn Ser Leu Gly Ile Ser Tyr Ala Trp Glu Ala Tyr
 1445 1450 1455
 Arg Lys Val Glu Gly Gly Ala Val Gln Leu Leu Glu Ala Gly Phe Asp
 1460 1465 1470
 Trp Glu Gly Ala Pro Met Asp Leu Pro Arg Gln Glu Leu Arg Val Ala
 1475 1480 1485
 Leu Glu Asn Asn Thr Glu Trp Ser Ser Tyr Phe Ser Thr Val Leu Gly
 1490 1495 1500
 Leu Thr Ala Phe Cys Gly Gly Phe Thr Ser Thr Asp Ser Lys Leu Gly
 1505 1510 1515 1520
 Tyr Glu Ala Asn Thr Gly Leu Arg Leu Ile Phe
 1525 1530

<210> 99

<211> 474

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 99

Met Lys Ile Ile His Thr Ala Ile Glu Phe Ala Pro Val Ile Lys Ala
 5 10 15
 Gly Gly Leu Gly Asp Ala Leu Tyr Gly Leu Ala Lys Ala Leu Ala Ala
 20 25 30
 Asn His Thr Thr Glu Val Val Ile Pro Leu Tyr Pro Lys Leu Phe Thr

Ieu	Pro	Lys	Glu	Gln	Asp	Leu	Cys	Ser	Ile	Gln	Lys	Leu	Ser	Tyr	Phe
	50					55					60				
Phe	Ala	Gly	Glu	Gln	Glu	Ala	Thr	Ala	Phe	Ser	Tyr	Phe	Tyr	Glu	Gly
65					70					75					80
Ile	Lys	Val	Thr	Leu	Phe	Lys	Leu	Asp	Thr	Gln	Pro	Glu	Leu	Phe	Glu
				85					90						95
Asn	Ala	Glu	Thr	Ile	Tyr	Thr	Ser	Asp	Ala	Phe	Arg	Phe	Cys	Ala	
			100					105					110		
Phe	Ser	Ala	Ala	Ala	Ala	Ser	Tyr	Ile	Gln	Lys	Glu	Gly	Ala	Asn	Ile
		115					120					125			
Val	His	Leu	His	Asp	Trp	His	Thr	Gly	Leu	Val	Ala	Gly	Leu	Leu	Lys
	130					135					140				
Gln	Gln	Pro	Cys	Ser	Gln	Leu	Gln	Lys	Ile	Val	Leu	Thr	Leu	His	Asn
145					150					155					160
Phe	Gly	Tyr	Arg	Gly	Tyr	Thr	Thr	Arg	Glu	Ile	Leu	Glu	Ala	Ser	Ser
				165					170					175	
Leu	Asn	Glu	Phe	Tyr	Ile	Ser	Gln	Tyr	Gln	Leu	Phe	Arg	Asp	Pro	Gln
			180					185					190		
Thr	Cys	Val	Leu	Leu	Lys	Gly	Ala	Leu	Tyr	Cys	Ser	Asp	Phe	Val	Thr
		195					200					205			
Thr	Val	Ser	Pro	Thr	Tyr	Ala	Lys	Glu	Ile	Leu	Glu	Asp	Tyr	Ser	Asp
	210					215					220				
Tyr	Glu	Ile	His	Asp	Ala	Ile	Thr	Ala	Arg	Gln	His	His	Leu	Arg	Gly
225					230					235					240
Ile	Leu	Asn	Gly	Ile	Asp	Thr	Thr	Ile	Trp	Gly	Pro	Glu	Thr	Asp	Pro
				245					250					255	
Asn	Leu	Ala	Lys	Asn	Tyr	Thr	Lys	Glu	Leu	Phe	Glu	Thr	Pro	Ser	Ile
		260						265					270		
Phe	Phe	Glu	Ala	Lys	Ala	Glu	Asn	Lys	Lys	Ala	Leu	Tyr	Glu	Arg	Leu
		275					280					285			
Gly	Leu	Ser	Leu	Glu	His	Ser	Pro	Cys	Val	Cys	Ile	Ser	Arg	Ile	
	290					295					300				
Ala	Glu	Gln	Lys	Gly	Pro	His	Phe	Met	Lys	Gln	Ala	Ile	Leu	His	Ala
305					310					315					320
Leu	Glu	Asn	Ala	Tyr	Thr	Leu	Ile	Ile	Ile	Gly	Thr	Cys	Tyr	Gly	Asn
				325						330				335	
Gln	Leu	His	Glu	Glu	Phe	Ala	Asn	Leu	Gln	Glu	Ser	Leu	Ala	Asn	Ser
			340					345					350		
Pro	Asp	Val	Arg	Ile	Leu	Leu	Thr	Tyr	Ser	Asp	Val	Leu	Ala	Arg	Gln
		355					360					365			
Ile	Phe	Ala	Ala	Ala	Asp	Met	Ile	Cys	Ile	Pro	Ser	Met	Phe	Glu	Pro
	370					375					380				
Cys	Gly	Leu	Thr	Gln	Met	Ile	Gly	Met	Arg	Tyr	Gly	Thr	Val	Pro	Leu
385					390										

$\langle 210 \rangle$ 100

<211> 393

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 100

```

Met Lys Lys Leu Leu Lys Ser Val Leu Val Phe Ala Ala Leu Ser Ser
      5      10      15
Ala Ser Ser Leu Gln Ala Leu Pro Val Gly Asn Pro Ala Glu Pro Ser
      20      25      30
Leu Met Ile Asp Gly Ile Leu Trp Glu Gly Phe Gly Gly Asp Pro Cys
      35      40      45
Asp Pro Cys Ala Thr Trp Cys Asp Ala Ile Ser Met Arg Val Gly Tyr
      50      55      60
Tyr Gly Asp Phe Val Phe Asp Arg Val Leu Lys Thr Asp Val Asn Lys
      65      70      75      80
Glu Phe Gln Met Gly Ala Lys Pro Thr Thr Asp Thr Gly Asn Ser Ala
      85      90      95
Ala Pro Ser Thr Leu Thr Ala Arg Glu Asn Pro Ala Tyr Gly Arg His
      100      105      110
Met Gln Asp Ala Glu Met Phe Thr Asn Ala Ala Cys Met Ala Leu Asn
      115      120      125
Ile Trp Asp Arg Phe Asp Val Phe Cys Thr Leu Gly Ala Thr Ser Gly
      130      135      140
Tyr Leu Lys Gly Asn Ser Ala Ser Phe Asn Leu Val Gly Leu Phe Gly
      145      150      155      160
Asp Asn Glu Asn Gln Lys Thr Val Lys Ala Glu Ser Val Pro Asn Met
      165      170      175
Ser Phe Asp Gln Ser Val Val Glu Leu Tyr Thr Asp Thr Thr Phe Ala
      180      185      190
Trp Ser Val Gly Ala Arg Ala Ala Leu Trp Glu Cys Gly Cys Ala Thr
      195      200      205
Leu Gly Ala Ser Phe Gln Tyr Ala Gln Ser Lys Pro Lys Val Glu Glu
      210      215      220
Leu Asn Val Leu Cys Asn Ala Ala Glu Phe Thr Ile Asn Lys Pro Lys
      225      230      235      240
Gly Tyr Val Gly Lys Glu Phe Pro Leu Asp Leu Thr Ala Gly Thr Asp
      245      250      255
Ala Ala Thr Gly Thr Lys Asp Ala Ser Ile Asp Tyr His Glu Trp Gln
      260      265      270
Ala Ser Leu Ala Leu Ser Tyr Arg Leu Asn Met Phe Thr Pro Tyr Ile
      275      280      285
Gly Val Lys Trp Ser Arg Ala Ser Phe Asp Ala Asp Thr Ile Arg Ile
      290      295      300
Ala Gln Pro Lys Ser Ala Thr Ala Ile Phe Asp Thr Thr Thr Leu Asn
      305      310      315      320
Pro Thr Ile Ala Gly Ala Gly Asp Val Lys Thr Gly Ala Glu Gly Gln
      325      330      335
Leu Gly Asp Thr Met Gln Ile Val Ser Leu Gln Leu Asn Lys Met Lys
      340      345      350
Ser Arg Lys Ser Cys Gly Ile Ala Val Gly Thr Thr Ile Val Asp Ala
      355      360      365
Asp Lys Tyr Ala Val Thr Val Glu Thr Arg Leu Ile Asp Glu Arg Ala
      370      375      380
Ala His Val Asn Ala Gln Phe Arg Phe
      385      390

```

<210> 101

<211> 195

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 101

```

Met Gly Ser Leu Val Gly Arg Gln Ala Pro Asp Phe Ser Gly Lys Ala

```

72

```

      5              10              15
Val Val Cys Gly Glu Glu Lys Glu Ile Ser Leu Ala Asp Phe Arg Gly
      20              25              30
Lys Tyr Val Val Leu Phe Phe Tyr Pro Lys Asp Phe Thr Tyr Val Cys
      35              40              45
Pro Thr Glu Leu His Ala Phe Gln Asp Arg Leu Val Asp Phe Glu Glu
      50              55              60
Arg Gly Ala Val Val Leu Gly Cys Ser Val Asp Asp Ile Glu Thr His
      65              70              75              80
Ser Arg Trp Leu Ala Val Ala Arg Asn Ala Gly Gly Ile Glu Gly Thr
      85              90              95
Glu Tyr Pro Leu Leu Ala Asp Pro Ser Phe Lys Ile Ser Glu Ala Phe
      100             105             110
Gly Val Leu Asn Pro Glu Gly Ser Leu Ala Leu Arg Ala Thr Phe Leu
      115             120             125
Ile Asp Lys Tyr Gly Val Val Arg His Ala Val Ile Asn Asp Leu Pro
      130             135             140
Leu Gly Arg Ser Ile Asp Glu Glu Leu Arg Ile Leu Asp Ser Leu Ile
      145             150             155             160
Phe Phe Glu Asn His Gly Met Val Cys Pro Ala Asn Trp Arg Ser Gly
      165             170             175
Glu Arg Gly Met Val Pro Ser Glu Glu Gly Leu Lys Glu Tyr Phe Gln
      180             185             190
Thr Met Asp
      195

```

<210> 102

<211> 86

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 102

```

Met Ser Gln Asn Lys Asn Ser Ala Phe Met Gln Pro Val Asn Val Ser
      5              10              15
Ala Asp Leu Ala Ala Ile Val Gly Ala Gly Pro Met Pro Arg Thr Glu
      20              25              30
Ile Ile Lys Lys Met Trp Asp Tyr Ile Lys Lys Asn Gly Leu Gln Asp
      35              40              45
Pro Thr Asn Lys Arg Asn Ile Asn Pro Asp Asp Lys Leu Ala Lys Val
      50              55              60
Phe Gly Thr Glu Lys Pro Ile Asp Met Phe Gln Met Thr Lys Met Val
      65              70              75              80
Ser Gln His Ile Ile Lys
      85

```

<210> 103

<211> 394

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 103

```

Met Ser Lys Glu Thr Phe Gln Arg Asn Lys Pro His Ile Asn Ile Gly
      5              10              15
Thr Ile Gly His Val Asp His Gly Lys Thr Thr Leu Thr Ala Ala Ile
      20              25              30
Thr Arg Ala Leu Ser Gly Asp Gly Leu Ala Asp Phe Arg Asp Tyr Ser
      35              40              45
Ser Ile Asp Asn Thr Pro Glu Glu Lys Ala Arg Gly Ile Thr Ile Asn
      50              55              60

```

73

Ala Ser His Val Glu Tyr Glu Thr Ala Asn Arg His Tyr Ala His Val
 65 70 75 80
 Asp Cys Pro Gly His Ala Asp Tyr Val Lys Asn Met Ile Thr Gly Ala
 85 90 95
 Ala Gln Met Asp Gly Ala Ile Leu Val Val Ser Ala Thr Asp Gly Ala
 100 105 110
 Met Pro Gln Thr Lys Glu His Ile Leu Leu Ala Arg Gln Val Gly Val
 115 120 125
 Pro Tyr Ile Val Val Phe Leu Asn Lys Ile Asp Met Ile Ser Glu Glu
 130 135 140
 Asp Ala Glu Leu Val Asp Leu Val Glu Met Glu Leu Val Glu Leu Leu
 145 150 155 160
 Glu Glu Lys Gly Tyr Lys Gly Cys Pro Ile Ile Arg Gly Ser Ala Leu
 165 170 175
 Lys Ala Leu Glu Gly Asp Ala Ala Tyr Ile Glu Lys Val Arg Glu Leu
 180 185 190
 Met Gln Ala Val Asp Asp Asn Ile Pro Thr Pro Glu Arg Glu Ile Asp
 195 200 205
 Lys Pro Phe Leu Met Pro Ile Glu Asp Val Phe Ser Ile Ser Gly Arg
 210 215 220
 Gly Thr Val Val Thr Gly Arg Ile Glu Arg Gly Ile Val Lys Val Ser
 225 230 235 240
 Asp Lys Val Gln Leu Val Gly Leu Arg Asp Thr Lys Glu Thr Ile Val
 245 250 255
 Thr Gly Val Glu Met Phe Arg Lys Glu Leu Pro Glu Gly Arg Ala Gly
 260 265 270
 Glu Asn Val Gly Leu Leu Leu Arg Gly Ile Gly Lys Asn Asp Val Glu
 275 280 285
 Arg Gly Met Val Val Cys Leu Pro Asn Ser Val Lys Pro His Thr Gln
 290 295 300
 Phe Lys Cys Ala Val Tyr Val Leu Gln Lys Glu Glu Gly Gly Arg His
 305 310 315 320
 Lys Pro Phe Phe Thr Gly Tyr Arg Pro Gln Phe Phe Phe Arg Thr Thr
 325 330 335
 Asp Val Thr Gly Val Val Thr Leu Pro Glu Gly Ile Glu Met Val Met
 340 345 350
 Pro Gly Asp Asn Val Glu Phe Glu Val Gln Leu Ile Ser Pro Val Ala
 355 360 365
 Leu Glu Glu Gly Met Arg Phe Ala Ile Arg Glu Gly Gly Arg Thr Ile
 370 375 380
 Gly Ala Gly Thr Ile Ser Lys Ile Ile Ala
 385 390

<210> 104

<211> 82

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 104

Met Gly Gln Asp His Arg Arg Lys Phe Leu Lys Lys Val Ser Phe Val
 5 10 15
 Lys Lys Gln Ala Ala Phe Ala Gly Asn Phe Ile Glu Glu Ile Lys Lys
 20 25 30
 Ile Glu Trp Val Asn Lys Arg Asp Leu Lys Arg Tyr Val Lys Ile Val
 35 40 45
 Leu Met Asn Ile Phe Gly Phe Gly Phe Ser Ile Tyr Cys Val Asp Leu
 50 55 60
 Ala Leu Arg Lys Ser Leu Ser Leu Phe Gly Lys Val Thr Ser Phe Phe
 65 70 75 80
 Phe Gly

<210> 105

<211> 379

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 105

```

Met Val Ile Pro Lys Val Asp Leu Gly Glu Ser Ala Val Met Met Gly
      5      10      15
Tyr Lys Leu Thr Ser Gln Leu Ala Met Leu Ser Ile Leu Leu Thr Phe
      20      25      30
Thr His Thr Met Gly His Ala Ser Gln Met Ser Gln Thr Leu Pro Thr
      35      40      45
Ile Ile Glu Ala Gln Ala Glu Glu Ala Leu Gln Ala Asp Arg Gly Val
      50      55      60
Ala Gly Gln Ala Leu Lys Lys Leu Arg Lys Lys Arg Cys Ala Ser Arg
      65      70      75      80
Lys Ser Ala Cys Lys Ala Ser Phe Lys Lys Lys Asp Phe Phe Ser Cys
      85      90      95
Ile Thr Asn Gly Leu Phe Ser Gly Asn His Glu Gln Arg Leu Thr Ala
      100      105      110
Lys Lys Glu Asn Lys Ala Arg Gly Lys Glu Pro Arg Val Val Val Gln
      115      120      125
Thr Thr Lys Lys Arg Gln Ile Thr Gln Ser Glu Lys Glu Phe Phe Asp
      130      135      140
Trp Leu Cys Asn Ser Lys Arg Glu Arg Lys Leu Leu Lys Lys Lys Pro
      145      150      155      160
Val Asn Thr Ser Leu Ala Lys Ser Glu Glu Leu Ser Pro Lys Glu Ala
      165      170      175
Ala Ile Ala Ala Ala Arg Ala Ser Leu Ser Pro Glu Glu Lys Arg Gln
      180      185      190
Leu Ile Arg Glu Trp Leu Ala Glu Glu Lys Thr Ala Arg Lys Ser Gly
      195      200      205
Arg Ala Ala Cys Ala Val Ser Glu Asn Leu Lys Arg Asp Gly Ser Ile
      210      215      220
Thr Ser Thr Leu Arg Tyr Asp Ala Glu Lys Ala Leu Thr Thr Arg Val
      225      230      235      240
Lys Arg Asn Glu Asn Ser Val Asn Ala Arg Ala Arg Gln Arg Ala Ala
      245      250      255
Leu Gln Lys Ala Lys Lys Ala Lys Thr Glu Lys Pro Glu Ala Asp Glu
      260      265      270
Lys Ala Ala Glu Ala Val Ala Ala Ala Pro Thr Lys Gln Ala His Lys
      275      280      285
Glu Pro Glu Asn Tyr Phe Ala Ala Thr Ala Ser Thr Asn Asn Thr Asn
      290      295      300
Val Met Ser Tyr Leu Asn Ala His Gln Tyr Arg Cys Asp Ser Ser Glu
      305      310      315      320
Thr Asp Trp Pro Cys Ser Ser Cys Val Thr Lys Arg Arg Ala Asn Phe
      325      330      335
Gly Ile Ser Val Cys Thr Met Val Val Thr Val Ile Ala Met Ile Val
      340      345      350
Gly Ala Val Ile Ile Ser Asn Ala Thr Asp Ser Thr Val Ala Gly Ser
      355      360      365
Ser Gly Thr Gly Gly Gly Gly Ser Thr Gln Pro
      370      375

```

<210> 106

<211> 563

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 106

```

Met Val Tyr Phe Arg Ala His Gln Pro Arg His Thr Pro Lys Thr Phe
      5      10      15
Pro Leu Glu Val His His Ser Phe Ser Asp Lys His Pro Gln Ile Ala
      20      25      30
Lys Ala Met Arg Ile Thr Gly Ile Ala Leu Ala Ala Leu Ser Leu Leu
      35      40      45
Ala Val Val Ala Cys Val Ile Ala Val Ser Ala Gly Gly Ala Ala Ile
      50      55      60
Pro Leu Ala Val Ile Ser Gly Ile Ala Val Met Ser Gly Leu Leu Ser
      65      70      75      80
Ala Ala Thr Ile Ile Cys Ser Ala Lys Lys Ala Leu Ala Gln Arg Lys
      85      90      95
Gln Lys Gln Leu Glu Glu Ser Leu Pro Leu Asp Asn Ala Thr Glu His
      100      105      110
Val Ser Tyr Leu Thr Ser Asp Thr Ser Tyr Phe Asn Gln Trp Glu Ser
      115      120      125
Leu Gly Ala Leu Asn Lys Gln Leu Ser Gln Ile Asp Leu Thr Ile Gln
      130      135      140
Ala Pro Glu Lys Lys Leu Leu Lys Glu Val Leu Gly Ser Arg Tyr Asp
      145      150      155      160
Ser Ile Asn His Ser Ile Glu Glu Ile Ser Asp Arg Phe Thr Lys Met
      165      170      175
Leu Ser Leu Leu Arg Leu Arg Glu His Phe Tyr Arg Gly Glu Glu Arg
      180      185      190
Tyr Ala Pro Tyr Leu Ser Pro Pro Leu Leu Asn Lys Asn Arg Leu Leu
      195      200      205
Thr Gln Ile Thr Ser Asn Met Ile Arg Met Leu Pro Lys Ser Gly Gly
      210      215      220
Val Phe Ser Leu Lys Ala Asn Thr Leu Ser His Ala Ser Arg Thr Leu
      225      230      235      240
Tyr Thr Val Leu Lys Val Ala Leu Ser Leu Gly Val Leu Ala Gly Val
      245      250      255
Ala Ala Leu Ile Ile Phe Leu Pro Pro Ser Leu Pro Phe Ile Ala Val
      260      265      270
Ile Gly Val Ser Ser Leu Ala Leu Gly Met Ala Ser Phe Leu Met Ile
      275      280      285
Arg Gly Ile Lys Tyr Leu Leu Glu His Ser Pro Leu Asn Arg Lys Gln
      290      295      300
Leu Ala Lys Asp Ile Gln Lys Thr Ile Gly Pro Asp Val Leu Ala Ser
      305      310      315      320
Met Val His Tyr Gln His Gln Leu Leu Ser His Leu His Glu Thr Leu
      325      330      335
Leu Asp Glu Ala Ile Thr Ala Arg Trp Ser Glu Pro Phe Phe Ile Glu
      340      345      350
His Ala Asn Leu Lys Ala Lys Ile Glu Asp Leu Thr Lys Gln Tyr Asp
      355      360      365
Ile Leu Asn Ala Ala Phe Asn Lys Ser Leu Gln Gln Asp Glu Ala Leu
      370      375      380
Arg Ser Gln Leu Glu Lys Arg Ala Tyr Leu Phe Pro Ile Pro Asn Asn
      385      390      395      400
Asp Glu Asn Ala Lys Thr Lys Glu Ser Gln Leu Leu Asp Ser Glu Asn
      405      410      415
Asp Ser Asn Ser Glu Phe Gln Glu Ile Ile Asn Lys Gly Leu Glu Ala
      420      425      430
Ala Asn Lys Arg Arg Ala Asp Ala Lys Ser Lys Phe Tyr Thr Glu Asp
      435      440      445
Glu Thr Ser Asp Lys Ile Phe Ser Ile Trp Lys Pro Thr Lys Asn Leu

```

450	455	460
Ala Leu Glu Asp Leu Trp Arg Val His Glu Ala Cys Asn Glu Glu Gln		
465	470	475
Gln Ala Leu Leu Leu Glu Asp Tyr Met Ser Tyr Lys Thr Ser Glu Cys		480
	485	490
Gln Ala Ala Leu Gln Lys Val Ser Gln Glu Leu Lys Ala Ala Gln Lys		495
	500	505
Ser Phe Ala Val Leu Glu Lys His Ala Leu Asp Arg Ser Tyr Glu Ser		510
	515	520
Ser Val Ala Thr Met Asp Leu Ala Arg Ala Asn Gln Glu Thr His Arg		525
	530	535
Leu Leu Asn Ile Leu Ser Glu Leu Gln Gln Leu Ala Gln Tyr Leu Leu		540
545	550	555
Asp Asn His		560

<210> 107

<211> 358

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 107

Met Arg Lys Thr Val Ile Val Ala Met Ser Gly Gly Val Asp Ser Ser	
	5 10 15
Val Val Ala Tyr Leu Leu Lys Lys Gln Gly Glu Tyr Asn Val Val Gly	
	20 25 30
Leu Phe Met Lys Asn Trp Gly Glu Gln Asp Glu Asn Gly Glu Cys Thr	
	35 40 45
Ala Thr Lys Asp Phe Arg Asp Val Glu Arg Ile Ala Glu Gln Leu Ser	
	50 55 60
Ile Pro Tyr Tyr Thr Val Ser Phe Ser Lys Glu Tyr Lys Glu Arg Val	
	65 70 75 80
Phe Ser Arg Phe Leu Arg Glu Tyr Ala Asn Gly Tyr Thr Pro Asn Pro	
	85 90 95
Asp Val Leu Cys Asn Arg Glu Ile Lys Phe Asp Leu Leu Gln Lys Lys	
	100 105 110
Val Arg Glu Leu Lys Gly Asp Phe Leu Ala Thr Gly His Tyr Cys Arg	
	115 120 125
Gly Gly Ala Asp Gly Thr Gly Leu Ser Arg Gly Ile Asp Pro Asn Lys	
	130 135 140
Asp Gln Ser Tyr Phe Leu Cys Gly Thr Pro Lys Asp Ala Leu Ser Asn	
	145 150 155 160
Val Leu Phe Pro Leu Gly Gly Met Tyr Lys Thr Glu Val Arg Arg Ile	
	165 170 175
Ala Gln Glu Ala Gly Leu Ala Thr Ala Thr Lys Lys Asp Ser Thr Gly	
	180 185 190
Ile Cys Phe Ile Gly Lys Arg Pro Phe Lys Ser Phe Leu Glu Gln Phe	
	195 200 205
Val Ala Asp Ser Pro Gly Asp Ile Ile Asp Phe Asp Thr Gln Gln Val	
	210 215 220
Val Gly Arg His Glu Gly Ala His Tyr Tyr Thr Ile Gly Gln Arg Arg	
	225 230 235 240
Gly Leu Asn Ile Gly Gly Met Glu Lys Pro Cys Tyr Val Leu Ser Lys	
	245 250 255
Asn Met Glu Lys Asn Ile Val Tyr Ile Val Arg Gly Glu Asp His Pro	
	260 265 270
Leu Leu Tyr Arg Gln Glu Leu Leu Ala Lys Glu Leu Asn Trp Phe Val	
	275 280 285
Pro Leu Gln Glu Pro Met Ile Cys Ser Ala Lys Val Arg Tyr Arg Ser	
	290 295 300

77

Pro Asp Glu Lys Cys Ser Val Tyr Pro Leu Glu Asp Gly Thr Val Lys
 305 310 315 320
 Val Ile Phe Asp Val Pro Val Lys Ala Val Thr Pro Gly Gln Thr Val
 325 330 335
 Ala Phe Tyr Gln Gly Asp Ile Cys Leu Gly Gly Gly Val Ile Glu Val
 340 345 350
 Pro Met Ile His Gln Leu
 355

<210> 108

<211> 267

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 108

Met Ser Arg Lys Pro Ala Ser Asn Ser Ser Arg Asn Thr Lys Arg Ser
 5 10 15
 Ser Asp Thr Ser Trp Glu Val Ile Ala Gln Asp Tyr Asn Lys Ala Val
 20 25 30
 Asp Arg Asp Gly His Phe Tyr His Lys Glu Val Ile Leu Pro Asn Leu
 35 40 45
 Leu Ser Lys Leu His Ile Ser Arg Ser Ser Ser Leu Val Asp Val Gly
 50 55 60
 Cys Gly Gln Gly Ile Leu Glu Lys His Leu Pro Lys His Leu Pro Tyr
 65 70 75 80
 Leu Gly Ile Asp Leu Ser Pro Ser Leu Leu Arg Phe Ala Lys Lys Ser
 85 90 95
 Ala Ser Ser Lys Ser Arg Arg Phe Leu His His Asp Met Thr Gln Pro
 100 105 110
 Val Pro Ala Asp His His Glu Gln Phe Ser His Ala Thr Ala Ile Leu
 115 120 125
 Ser Leu Gln Asn Met Glu Ser Pro Glu Gln Ala Ile Ala His Thr Ala
 130 135 140
 Asn Leu Leu Ala Pro Gln Gly Arg Leu Phe Ile Val Leu Asn His Pro
 145 150 155 160
 Cys Phe Arg Ile Pro Arg Leu Ser Ser Trp Leu Tyr Asp Glu Pro Lys
 165 170 175
 Lys Leu Leu Ser Arg Lys Ile Asp Arg Tyr Leu Ser Pro Val Ala Val
 180 185 190
 Pro Ile Val Val His Pro Gly Glu Lys His Ser Glu Thr Thr Tyr Ser
 195 200 205
 Phe His Phe Pro Leu Ser Tyr Trp Val Gln Ala Leu Ser Asn His Asn
 210 215 220
 Leu Leu Ile Asp Ser Met Glu Glu Trp Ile Ser Pro Lys Lys Ser Ser
 225 230 235 240
 Gly Lys Arg Ala Arg Ala Glu Asn Leu Cys Arg Lys Glu Phe Pro Leu
 245 250 255
 Phe Leu Phe Ile Ser Ala Leu Lys Ile Ser Lys
 260 265

<210> 109

<211> 867

<212> PRT

<213> Chlamydia trachomatis serovar D

<400> 109

Met Glu Lys Phe Ser Asp Ala Val Ser Glu Ala Leu Glu Lys Ala Phe
 5 10 15

Glu Leu Ala Lys Asn Ser Lys His Ser Tyr Val Thr Glu Asn His Leu
 20 25 30
 Leu Lys Ser Leu Leu Gln Asn Pro Gly Ser Leu Phe Cys Leu Val Ile
 35 40 45
 Lys Asp Val His Gly Asn Leu Gly Leu Leu Thr Ser Ala Val Asp Asp
 50 55 60
 Ala Leu Arg Arg Glu Pro Thr Val Val Glu Gly Thr Ala Val Ala Ser
 65 70 75 80
 Pro Ser Pro Ser Leu Gln Gln Leu Leu Leu Asn Ala His Gln Glu Ala
 85 90 95
 Arg Ser Met Gly Asp Glu Tyr Leu Ser Gly Asp His Leu Leu Leu Ala
 100 105 110
 Phe Trp Arg Ser Thr Lys Glu Pro Phe Ala Ser Trp Arg Lys Thr Val
 115 120 125
 Lys Thr Thr Ser Glu Ala Leu Lys Glu Leu Ile Thr Lys Leu Arg Gln
 130 135 140
 Gly Ser Arg Met Asp Ser Pro Ser Ala Glu Glu Asn Leu Lys Gly Leu
 145 150 155 160
 Glu Lys Tyr Cys Lys Asn Leu Thr Val Leu Ala Arg Glu Gly Lys Leu
 165 170 175
 Asp Pro Val Ile Gly Arg Asp Glu Glu Ile Arg Arg Thr Ile Gln Val
 180 185 190
 Leu Ser Arg Arg Thr Lys Asn Asn Pro Met Leu Ile Gly Glu Pro Gly
 195 200 205
 Val Gly Lys Thr Ala Ile Ala Glu Gly Leu Ala Leu Arg Ile Val Gln
 210 215 220
 Gly Asp Val Pro Glu Ser Leu Lys Glu Lys His Leu Tyr Val Leu Asp
 225 230 235 240
 Met Gly Ala Leu Ile Ala Gly Ala Lys Tyr Arg Gly Glu Phe Glu Glu
 245 250 255
 Arg Leu Lys Ser Val Leu Lys Gly Val Glu Ala Ser Glu Gly Glu Cys
 260 265 270
 Ile Leu Phe Ile Asp Glu Val His Thr Leu Val Gly Ala Gly Ala Thr
 275 280 285
 Asp Gly Ala Met Asp Ala Ala Asn Leu Leu Lys Pro Ala Leu Ala Arg
 290 295 300
 Gly Thr Leu His Cys Ile Gly Ala Thr Thr Leu Asn Glu Tyr Gln Lys
 305 310 315 320
 Tyr Ile Glu Lys Asp Ala Ala Leu Glu Arg Arg Phe Gln Pro Ile Phe
 325 330 335
 Val Thr Glu Pro Ser Leu Glu Asp Ala Val Phe Ile Leu Arg Gly Leu
 340 345 350
 Arg Glu Lys Tyr Glu Ile Phe His Gly Val Arg Ile Thr Glu Gly Ala
 355 360 365
 Leu Asn Ala Ala Val Val Leu Ser Tyr Arg Tyr Ile Thr Asp Arg Phe
 370 375 380
 Leu Pro Asp Lys Ala Ile Asp Leu Ile Asp Glu Ala Ala Ser Leu Ile
 385 390 395 400
 Arg Met Gln Ile Gly Ser Leu Pro Leu Pro Ile Asp Glu Lys Glu Arg
 405 410 415
 Glu Leu Ser Ala Leu Ile Val Lys Gln Glu Ala Ile Lys Arg Glu Gln
 420 425 430
 Ala Pro Ala Tyr Gln Glu Glu Ala Glu Asp Met Gln Lys Ala Ile Asp
 435 440 445
 Arg Val Lys Glu Glu Leu Ala Ala Leu Arg Leu Arg Trp Asp Glu Glu
 450 455 460
 Lys Gly Leu Ile Thr Gly Leu Lys Glu Lys Lys Asn Ala Leu Glu Asn
 465 470 475 480
 Leu Lys Phe Ala Glu Glu Glu Ala Glu Arg Thr Ala Asp Tyr Asn Arg
 485 490 495
 Val Ala Glu Leu Arg Tyr Ser Leu Ile Pro Ser Leu Glu Glu Glu Ile


```

      500      505      510
His Leu Ala Glu Glu Ala Leu Asn Gln Arg Asp Gly Arg Leu Leu Gln
      515      520      525
Glu Glu Val Asp Glu Arg Leu Ile Ala Gln Val Val Ala Asn Trp Thr
      530      535      540
Gly Ile Pro Val Gln Lys Met Leu Glu Gly Glu Ser Glu Lys Leu Leu
545      550      555      560
Val Leu Glu Glu Ser Leu Glu Glu Arg Val Val Gly Gln Pro Phe Ala
      565      570      575
Ile Ala Ala Val Ser Asp Ser Ile Arg Ala Ala Arg Val Gly Leu Ser
      580      585      590
Asp Pro Gln Arg Pro Leu Gly Val Phe Leu Phe Leu Gly Pro Thr Gly
      595      600      605
Val Gly Lys Thr Glu Leu Ala Lys Ala Leu Ala Glu Leu Leu Phe Asn
      610      615      620
Lys Glu Glu Ala Met Ile Arg Phe Asp Met Thr Glu Tyr Met Glu Lys
625      630      635      640
His Ser Val Ser Lys Leu Ile Gly Ser Pro Pro Gly Tyr Val Gly Tyr
      645      650      655
Glu Glu Gly Gly Ser Leu Ser Glu Ala Leu Arg Arg Arg Pro Tyr Ser
      660      665      670
Val Val Leu Phe Asp Glu Ile Glu Lys Ala Asp Lys Glu Val Phe Asn
      675      680      685
Ile Leu Leu Gln Ile Phe Asp Gly Ile Leu Thr Asp Ser Lys Lys
      690      695      700
Arg Lys Val Asn Cys Lys Asn Ala Leu Phe Ile Met Thr Ser Asn Ile
705      710      715      720
Gly Ser Gln Glu Leu Ala Asp Tyr Cys Thr Lys Lys Gly Thr Ile Val
      725      730      735
Asp Lys Glu Ala Val Leu Ser Val Val Ala Pro Ala Leu Lys Asn Tyr
      740      745      750
Phe Ser Pro Glu Phe Ile Asn Arg Ile Asp Asp Ile Leu Pro Phe Val
      755      760      765
Pro Leu Thr Thr Glu Asp Ile Val Lys Ile Val Gly Ile Gln Met Asn
      770      775      780
Arg Val Ala Leu Arg Leu Leu Glu Arg Lys Ile Ser Leu Thr Trp Asp
785      790      795      800
Asp Ser Leu Val Leu Phe Leu Ser Glu Gln Gly Tyr Asp Ser Ala Phe
      805      810      815
Gly Ala Arg Pro Leu Lys Arg Leu Ile Gln Gln Lys Val Val Thr Met
      820      825      830
Leu Ser Lys Ala Leu Leu Lys Gly Asp Ile Lys Pro Gly Met Ala Val
      835      840      845
Glu Leu Thr Met Ala Lys Asp Val Val Val Phe Lys Ile Lys Thr Asn
      850      855      860
Pro Ala Val
865

```

<210> 110

<211> 1170

<212> DNA

<213> Chlamydia pneumoniae

<400> 110

```

atgaaaaaac tcttaaagtc ggcgttatta tccgccgcat ttgctgggttc tgttggtcc 60
ttacaagcct tgccgttagg gaacccttct gatccaagct tattaattga tggtagaata 120
tggaaggtg ctgcaggaga tccttgcgat ccttgcgcta cttggtgcga cgctattagc 180
ttacgtgctg gattttacgg agactatggt ttgcaccgta tcttaaaagt agatgcacct 240
aaaacatttt ctatgggagc caagcctact ggatccgctg ctgcaaaacta tactactgcc 300
gtagatagac ctaaccgggc ctacaataag catttacacg atgcagagtg gttcactaat 360

```

```

gcaggcttca ttgccttaaa catttgggat cgctttgatg ttttctgtac tttaggagct 420
tctaattggtt acatttagagg aaactctaca gcgttcaatc tcgttggttt attcggagtt 480
aaaggtaacta ctgtaaatgc aaatgaacta ccaaacgttt cttaagtaa cggagttggt 540
gaacttttaca cagacacctc tttctcttgg agcgtaggcg ctogtggagc cttatgggaa 600
tgcggttggtg caactttggg agctgaatc caatatgcac agtccaaacc taaagttgaa 660
gaacttaatg tgatctgtaa cgtatcgcaa ttctctgtaa acaaacccaa gggctataaa 720
ggcgttgctt tccccttgcc aacagacgct ggcgtagcaa cagctactgg aacaaagtct 780
gcgaccatca attatcatga atggcaagta ggagcctctc tatcttacag actaaactct 840
ttagtgccat acattggagt acaatggtct cgagcaactt ttgatgctga taacatccgc 900
attgctcagc caaaactacc tacagctggt ttaaacttaa ctgcatggaa cccttcttta 960
ctaggaaatg ccacagcatt gtctactact gattcgttct cagacttcat gcaaattggt 1020
tctgtcaga tcaacaagtt taaatctaga aaagcttggt gagttactgt aggagctact 1080
ttagttgatg ctgataaatg gtcacttact gcagaagctc gtttaattaa cgagagagct 1140
gtcacggtat ctggtcagtt cagattctaa 1170

```

<210> 111

<211> 2601

<212> DNA

<213> *Chlamydia pneumoniae*

<400> 111

```

atggagaaat tttccgatgc tgtctctgaa gctttagaga aggccttccga acttgctaaa 60
tcttcgaaac atacctatgt cacagaaaat cacctattac tggctttatt agaaaatata 120
gagtctctct tttatttggg aattaaggac attcatggga accctggttt gctcaataacg 180
gcagttaaaag atgcgctctc acgagagccg actgtagttg aaggagaggt ggatcctaaa 240
ccttctccgg gtttacaac ccttcttagg gatgccaaac aagaggcaaa gacattagga 300
gatgaatata tttctggaga tcatctgctg cttgcttttt ggagttcaaa caaagagcct 360
tttaattctt ggaagcaaac aacaaaagtt agttttaaag atcttaagaa tctgattact 420
aaaatacgac gaggaaatcg tatggattcg ccaagcgctg aaagtaattt tcagggttta 480
gaaaagtatt gtaaaaaattt aacagcatta gctcgtgaag gttaaactgga tcctgtgatc 540
ggtagagatg aagaaattcg tagaaccatc caagtgcctt cccgtagaac taaaaataac 600
cctatgctta ttggtgagcc ggggtgtagg aaaactgcta tagcagaagg attagctctt 660
aggcttatcc aggggtgatgt tcctgaatct ctcaaaggta aacagcttta tgtcttagat 720
atgggagctt tgattgcagg agctaagtat cgagggtgagt ttgaagaaag actaaagagt 780
gttttaaaag atgtagaatc tggagatggc gaggcacatta totttattga tgaggtgcat 840
actcttgttg gagcaggagc tactgatgga gctatggatg ctgcgaatct tttaaagcct 900
gcattagcaa gagggacgct acactgtatt ggcgogacga ctttgaatga gtatcagaag 960
tatattgaaa aagatgctgc tttggaacgt cgatttcagc ctatttttgt gacagagcct 1020
tctttggagg atgctgtctt tattcttcgt ggactaagag aaaaatatga aattttccat 1080
ggagtcaggg ttacagaggg ggctttgaat gccgcagtcct tactttcota tcgttatatc 1140
ccagatcgct ttcttccaga taaggctatc gatctgatag atgaagcggc aagtttaatt 1200
cgcatgcaaa ttggtagtct tcctcttctc attgatgaaa aggagagaga gcttgctgct 1260
ttgatcggtt agcaagaggc tataaaacgc gagcaatctc cttcctatca agaagaggcg 1320
gatgctatgc agaagtctat agatgctttg agagaggaat tagcatctct acgtttgggt 1380
tgggatgaag agaagaagtt gatttcgggg ctcaaggaaa aaaagaatc cttggaaagt 1440
atgaaatttt ctgaagagga ggcggagcgt gttgcagact ataatcgtgt agctgagctt 1500
cggtatagtt taattcccca acttgaagaa gaaatcaaac aggatgaagc ctctttaaat 1560
caaagagata accgtctcct tcaagaagaa gttgacgagc gattgattgc gcaagtggta 1620
gctaattgga cagggattcc tgtgcaaaaa atgctagaag ggaagctga gaaactgta 1680
attcttgaag aatccttaga agaacgtgtg gtaggacagc cttttgcagt ctctgcgggt 1740
agtgattcta ttcgtgctgc acgtgtaggt ttaaattgatc ctcaacgtcc cttaggagtc 1800
tttttatttt tagggccaac aggggtagga aaaaccgagc ttgcaaaagc tcttgcatat 1860
cttcttttca ataaagagga agctatgggt cgcttcgata tgtcagagta tatggaaaag 1920
cattccattt ccaagcttat aggatcttct ccagggtatg tgggttatga ggaaggtggg 1980
agtctttctg aggcctcttg acgacgtccc tattcagtag ttctctttga tgagatagag 2040
aaagcagata aggaagtctt aaatatcctt ttacaggttt ttgatgatgg gattottacg 2100
gatgggaaaa aacgcaaaat aaattgtaaa aatgccttgt ttatcatgac atcaaatata 2160
ggttctccag aacttcgaga ttattgttca aaaaaaggaa gtgagcttac gaaagaagcg 2220
attctttctg tagtctctcc agtattgaaa agatacttga gccctgaatt tatgaaccga 2280
attgatgaga tacttccctt tgttccatta acgaaagaag atatcgtgaa aatagttggc 2340
attcaaatgc gaaggattgc ccagagatta aaggcacggc ggatcaattt atcttgggat 2400

```

```

gattctgtaa tattatttct tagtgaacag ggttatgaca gtgctttcgg agcccgccct 2460
ttaaaccgtt tgatccaaca aaaagtgtg atcttgcttt ctaaggcttt gcttaaagga 2520
gatattaaac ctgatacatc gattgagttg acgatggcaa aagaggtgct cgtatttaaa 2580
aaagtggaaa ctcttctta g                                     2601

```

<210> 112

<211> 389

<212> PRT

<213> Chlamydia pneumoniae

<400> 112

```

Met Lys Lys Leu Leu Lys Ser Ala Leu Leu Ser Ala Ala Phe Ala Gly
      5      10      15
Ser Val Gly Ser Leu Gln Ala Leu Pro Val Gly Asn Pro Ser Asp Pro
      20      25      30
Ser Leu Leu Ile Asp Gly Thr Ile Trp Glu Gly Ala Ala Gly Asp Pro
      35      40      45
Cys Asp Pro Cys Ala Thr Trp Cys Asp Ala Ile Ser Leu Arg Ala Gly
      50      55      60
Phe Tyr Gly Asp Tyr Val Phe Asp Arg Ile Leu Lys Val Asp Ala Pro
      65      70      75      80
Lys Thr Phe Ser Met Gly Ala Lys Pro Thr Gly Ser Ala Ala Ala Asn
      85      90      95
Tyr Thr Thr Ala Val Asp Arg Pro Asn Pro Ala Tyr Asn Lys His Leu
      100     105     110
His Asp Ala Glu Trp Phe Thr Asn Ala Gly Phe Ile Ala Leu Asn Ile
      115     120     125
Trp Asp Arg Phe Asp Val Phe Cys Thr Leu Gly Ala Ser Asn Gly Tyr
      130     135     140
Ile Arg Gly Asn Ser Thr Ala Phe Asn Leu Val Gly Leu Phe Gly Val
      145     150     155     160
Lys Gly Thr Thr Val Asn Ala Asn Glu Leu Pro Asn Val Ser Leu Ser
      165     170     175
Asn Gly Val Val Glu Leu Tyr Thr Asp Thr Ser Phe Ser Trp Ser Val
      180     185     190
Gly Ala Arg Gly Ala Leu Trp Glu Cys Gly Cys Ala Thr Leu Gly Ala
      195     200     205
Glu Phe Gln Tyr Ala Gln Ser Lys Pro Lys Val Glu Leu Asn Val
      210     215     220
Ile Cys Asn Val Ser Gln Phe Ser Val Asn Lys Pro Lys Gly Tyr Lys
      225     230     235     240
Gly Val Ala Phe Pro Leu Pro Thr Asp Ala Gly Val Ala Thr Ala Thr
      245     250     255
Gly Thr Lys Ser Ala Thr Ile Asn Tyr His Glu Trp Gln Val Gly Ala
      260     265     270
Ser Leu Ser Tyr Arg Leu Asn Ser Leu Val Pro Tyr Ile Gly Val Gln
      275     280     285
Trp Ser Arg Ala Thr Phe Asp Ala Asp Asn Ile Arg Ile Ala Gln Pro
      290     295     300
Lys Leu Pro Thr Ala Val Leu Asn Leu Thr Ala Trp Asn Pro Ser Leu
      305     310     315     320
Leu Gly Asn Ala Thr Ala Leu Ser Thr Thr Asp Ser Phe Ser Asp Phe
      325     330     335
Met Gln Ile Val Ser Cys Gln Ile Asn Lys Phe Lys Ser Arg Lys Ala
      340     345     350
Cys Gly Val Thr Val Gly Ala Thr Leu Val Asp Ala Asp Lys Trp Ser
      355     360     365
Leu Thr Ala Glu Ala Arg Leu Ile Asn Glu Arg Ala Ala His Val Ser
      370     375     380
Gly Gln Phe Arg Phe

```

385

<210> 113

<211> 866

<212> PRT

<213> Chlamydia pneumoniae

<400> 113

```

Met Glu Lys Phe Ser Asp Ala Val Ser Glu Ala Leu Glu Lys Ala Phe
      5      10      15
Glu Leu Ala Lys Ser Ser Lys His Thr Tyr Val Thr Glu Asn His Leu
      20      25      30
Leu Leu Ala Leu Leu Glu Asn Thr Glu Ser Leu Phe Tyr Leu Val Ile
      35      40      45
Lys Asp Ile His Gly Asn Pro Gly Leu Leu Asn Thr Ala Val Lys Asp
      50      55      60
Ala Leu Ser Arg Glu Pro Thr Val Val Glu Gly Glu Val Asp Pro Lys
      65      70      75      80
Pro Ser Pro Gly Leu Gln Thr Leu Leu Arg Asp Ala Lys Gln Glu Ala
      85      90      95
Lys Thr Leu Gly Asp Glu Tyr Ile Ser Gly Asp His Leu Leu Leu Ala
      100      105      110
Phe Trp Ser Ser Asn Lys Glu Pro Phe Asn Ser Trp Lys Gln Thr Thr
      115      120      125
Lys Val Ser Phe Lys Asp Leu Lys Asn Leu Ile Thr Lys Ile Arg Arg
      130      135      140
Gly Asn Arg Met Asp Ser Pro Ser Ala Glu Ser Asn Phe Gln Gly Leu
      145      150      155      160
Glu Lys Tyr Cys Lys Asn Leu Thr Ala Leu Ala Arg Glu Gly Lys Leu
      165      170      175
Asp Pro Val Ile Gly Arg Asp Glu Glu Ile Arg Arg Thr Ile Gln Val
      180      185      190
Leu Ser Arg Arg Thr Lys Asn Asn Pro Met Leu Ile Gly Glu Pro Gly
      195      200      205
Val Gly Lys Thr Ala Ile Ala Glu Gly Leu Ala Leu Arg Leu Ile Gln
      210      215      220
Gly Asp Val Pro Glu Ser Leu Lys Gly Lys Gln Leu Tyr Val Leu Asp
      225      230      235      240
Met Gly Ala Leu Ile Ala Gly Ala Lys Tyr Arg Gly Glu Phe Glu Glu
      245      250      255
Arg Leu Lys Ser Val Leu Lys Asp Val Glu Ser Gly Asp Gly Glu His
      260      265      270
Ile Ile Phe Ile Asp Glu Val His Thr Leu Val Gly Ala Gly Ala Thr
      275      280      285
Asp Gly Ala Met Asp Ala Ala Asn Leu Leu Lys Pro Ala Leu Ala Arg
      290      295      300
Gly Thr Leu His Cys Ile Gly Ala Thr Thr Leu Asn Glu Tyr Gln Lys
      305      310      315      320
Tyr Ile Glu Lys Asp Ala Ala Leu Glu Arg Arg Phe Gln Pro Ile Phe
      325      330      335
Val Thr Glu Pro Ser Leu Glu Asp Ala Val Phe Ile Leu Arg Gly Leu
      340      345      350
Arg Glu Lys Tyr Glu Ile Phe His Gly Val Arg Ile Thr Glu Gly Ala
      355      360      365
Leu Asn Ala Ala Val Leu Leu Ser Tyr Arg Tyr Ile Pro Asp Arg Phe
      370      375      380
Leu Pro Asp Lys Ala Ile Asp Leu Ile Asp Glu Ala Ala Ser Leu Ile
      385      390      395      400
Arg Met Gln Ile Gly Ser Leu Pro Leu Pro Ile Asp Glu Lys Glu Arg
      405      410      415

```

Glu Leu Ala Ala Leu Ile Val Lys Gln Glu Ala Ile Lys Arg Glu Gln
 420 425 430
 Ser Pro Ser Tyr Gln Glu Glu Ala Asp Ala Met Gln Lys Ser Ile Asp
 435 440 445
 Ala Leu Arg Glu Glu Leu Ala Ser Leu Arg Leu Gly Trp Asp Glu Glu
 450 455 460
 Lys Lys Leu Ile Ser Gly Leu Lys Glu Lys Lys Asn Ser Leu Glu Ser
 465 470 475 480
 Met Lys Phe Ser Glu Glu Glu Ala Glu Arg Val Ala Asp Tyr Asn Arg
 485 490 495
 Val Ala Glu Leu Arg Tyr Ser Leu Ile Pro Gln Leu Glu Glu Glu Ile
 500 505 510
 Lys Gln Asp Glu Ala Ser Leu Asn Gln Arg Asp Asn Arg Leu Leu Gln
 515 520 525
 Glu Glu Val Asp Glu Arg Leu Ile Ala Gln Val Val Ala Asn Trp Thr
 530 535 540
 Gly Ile Pro Val Gln Lys Met Leu Glu Gly Glu Ala Glu Lys Leu Leu
 545 550 555 560
 Ile Leu Glu Glu Ser Leu Glu Glu Arg Val Val Gly Gln Pro Phe Ala
 565 570 575
 Val Ser Ala Val Ser Asp Ser Ile Arg Ala Ala Arg Val Gly Leu Asn
 580 585 590
 Asp Pro Gln Arg Pro Leu Gly Val Phe Leu Phe Leu Gly Pro Thr Gly
 595 600 605
 Val Gly Lys Thr Glu Leu Ala Lys Ala Leu Ala Asp Leu Leu Phe Asn
 610 615 620
 Lys Glu Glu Ala Met Val Arg Phe Asp Met Ser Glu Tyr Met Glu Lys
 625 630 635 640
 His Ser Ile Ser Lys Leu Ile Gly Ser Ser Pro Gly Tyr Val Gly Tyr
 645 650 655
 Glu Glu Gly Gly Ser Leu Ser Glu Ala Leu Arg Arg Arg Pro Tyr Ser
 660 665 670
 Val Val Leu Phe Asp Glu Ile Glu Lys Ala Asp Lys Glu Val Leu Asn
 675 680 685
 Ile Leu Leu Gln Val Phe Asp Asp Gly Ile Leu Thr Asp Gly Lys Lys
 690 695 700
 Arg Lys Val Asn Cys Lys Asn Ala Leu Phe Ile Met Thr Ser Asn Ile
 705 710 715 720
 Gly Ser Pro Glu Leu Ala Asp Tyr Cys Ser Lys Lys Gly Ser Glu Leu
 725 730 735
 Thr Lys Glu Ala Ile Leu Ser Val Val Ser Pro Val Leu Lys Arg Tyr
 740 745 750
 Leu Ser Pro Glu Phe Met Asn Arg Ile Asp Glu Ile Leu Pro Phe Val
 755 760 765
 Pro Leu Thr Lys Glu Asp Ile Val Lys Ile Val Gly Ile Gln Met Arg
 770 775 780
 Arg Ile Ala Gln Arg Leu Lys Ala Arg Arg Ile Asn Leu Ser Trp Asp
 785 790 795 800
 Asp Ser Val Ile Leu Phe Leu Ser Glu Gln Gly Tyr Asp Ser Ala Phe
 805 810 815
 Gly Ala Arg Pro Leu Lys Arg Leu Ile Gln Gln Lys Val Val Ile Leu
 820 825 830
 Leu Ser Lys Ala Leu Leu Lys Gly Asp Ile Lys Pro Asp Thr Ser Ile
 835 840 845
 Glu Leu Thr Met Ala Lys Glu Val Leu Val Phe Lys Lys Val Glu Thr
 850 855 860
 Pro Ser
 865

<211> 1179
 <212> DNA
 <213> Homo sapiens

<400> 114
 taactctccc ctctcttctt aaaaaagagg ggagcctttt ttctttacaa agatacgccta 60
 gcttttttct gaagaatctc atcaagagat atttgcatit tcccacggat aaaggcatcc 120
 caaggaagcc ctggaatcac ttcatattct cccgttgcta gcattcgaca agggaaacca 180
 aagattaaat cttccggtaa tccataggga ttgtgggtccg aacacactcc ggaagaaaac 240
 cattctcctt cttttggctg atatattgat cgagcagcct ctgctaaga cctgtgctgca 300
 gaagctgccg aagacttccc tctgtcttcg attactgcac taccacgact ctgtacagaa 360
 ggccaccataa tattctctaa ccaatcacga tccgctatcg tctctgcgat aggacggtca 420
 ttaatcagag cttgcgtaaa atcaggcact tgtttggcgg agtgatttcc ccaaaccaca 480
 acttgtgata cagccgataa aggtacttct gctctatgcy ataactctgg agcatgattc 540
 ttctgggtcca atcgtagcat cgcagaaaag ttcttttcta ataactctgg agcatgattc 600
 attgctatcc agcaattggt attcacaggg ttcccaacaa caaaaatctt tgcattccgc 660
 ttggctgttg tgttcaaaag ttttcttgc gtagcaaaaa tctccctatt tttctttaga 720
 agatcccttc tctccattcc tgggcctcta ggaactgacc ctataaggaa tgcgcgatca 780
 atgccatcaa aagcatcatg caatgatgtc gttacctgca cagcgtgtaa taaagggaaa 840
 gcaccatcat ctagctccat gcgcacacca gataaagccc tttctgttcc aggaatatcg 900
 tagatacgca gatcgatgcc acaatcaagg ccaaaaacat ctccatgagc cagagaaaaat 960
 agaaagctat aggctatttg cctgttctct cctgttactg ctacactcac tgtttgagaa 1020
 accataagcc accctctctt tacttttaca aaacgcacat actctcaaca ctacgtttgc 1080
 aactaactaa ttttgggtccc aacatacggt tggatgataa aagaatcaag tacctagatt 1140
 cottagtaaa agcttttggc aaaaaaagc tcatctatt 1179

<210> 115
 <211> 772
 <212> DNA
 <213> Homo sapiens

<400> 115
 gcaaaactgc tgacaaagct ggagacggaa ctacaacagc tactgttctt gctgaagcta 60
 tctatacaga aggattacgc aatgtaacag ctggagcaaa tccaatggac ctcaaacgag 120
 gtattgataa agctgttaag gttgttgttg atcaaatacag aaaaatcagc aaacctgttc 180
 agcatcataa agaaattgct caagttgcaa caatttctgc taataatgat gcagaaatcg 240
 ggaatctgat tgctgaagca atggagaaaag ttggtaaaaa cggctctatc actgttgaag 300
 aagcaaaaag atttgaacc gttttggatg ttgttgaagg aatgaatttc aatagaggtt 360
 acctctctag ctacttcgca acaaatccag aaactcaaga atgtgtatta gaagacgctt 420
 tggttctaat ctacgataag aaaatttctg ggatcaaaga tttccttctt gttttacaac 480
 aagttgctga atccggcctt cctcttctta ttatagcaga agacattgaa ggcgaagctt 540
 tagctacttt ggtcgtgaac agaattcgtg gaggattccg ggtttgcgca gttaaagctc 600
 caggcttttg agatagaaga aaagctatgt tgggaagacat cgctatctta actggcggtc 660
 aactcattag cgaagagttg ggcatgaaat tagaaaacgc taacttagct atgttaggta 720
 aagctaaaaa agttatcgtt tctaaggaag acacgaccat cgtcgaagga at 772

<210> 116
 <211> 487
 <212> DNA
 <213> Homo sapiens

<400> 116
 gcagctcctg caaagccaca agctcctgtc gcacaaacac ggcatittta aaagagccat 60
 cagattttct ctcttaattt tacgcagtct tcccaacagg tgaataaacc tgaggaaaga 120
 agacgtcctt tggagtctcg atacttacaa ggccgaggta agcaggcagc tgcgtcaaag 180
 gaaaaaaaag ctcttgaaca ggaagtatcc aaacaagaag aagaagcttc taaactctgg 240
 gaagagaaac agagttatgc tctgtctgct gtgaatgcca tcaatttcag tgtaagaaag 300
 caaatagaag agcaacagaa aaccatttcc aatccaggaa atgaccagac tcttctctgg 360
 aagaaagatc cacatacatc cggagaacct gttatccaaa cggtaacaaga ctgttctcag 420

85

gatcaagaag aagagaaaa agttctagag cgattaaaca aacgttctct gacgtgtcag 480
gatctta 487

<210> 117

<211> 1014

<212> DNA

<213> Homo sapiens

<400> 117

ctcgtgccga atctttctaac aagagaacaa gctcctttct ttctttttcta aacaagggtc 60
agcgcctttct attaaaaagaa accctattca gaccctatgc agcacatagt ttataaaaaa 120
attttttctat taacagagga aaaataacct attgataaac agagcggtag aaggagatgc 180
aaataaagct gcttttaggat ccttacctag attctagaaa atggttgcat gaatttgaac 240
aaacaaacta attaaaaatt aaaactgaaa aaaatagttt aaacaacaa ctagaggata 300
ttttttcatg gcgctaaaag atacggcaaa aaaaatgact gacttggttg aaagtatcca 360
acaaaatttg cttaaagcag aaaaaggaaa taaagccgca gcacaaagag ttcgtacaga 420
atctatcaaa ttagaaaaaga tcgcgaaggt atatcgtaaa gagtccatta aagcagaaaa 480
aatgggctta atgaaaaaaa gcaagccgc tgcataaaaa gctaaagctg ctgctaagaa 540
gcctgttcgc gctacaaaaa cagtggctaa aaaagcttgt acaaaaagaa cttgtgctac 600
taaagcaaa gtcaaaccaa caaaaaagc cgtcctctaa acaaaagtta aaacagcgaa 660
aaaaaactgc tcaacaaaaa aataatattt tagcgcttct tcttttttat agagggcact 720
tttatcaaca gggccctctt tctctttctc attgatccct tctctttttt ttgttatcct 780
ttccgttctc gcaaaagcaa gtccttgcaa ataaaagtac aacctcacac ctcccttgga 840
ggaaaaacct ttcactttct ttaggattca agttgctctc ctgctatcgt aactgtaaac 900
attttggcgt ctgtggaggc tgttcattct ctcaaattga atatgcatcc tctttaaaaa 960
caaaagagct tgcgctccat aatttatattg caccctttat cccatcccaa aata 1014

<210> 118

<211> 287

<212> DNA

<213> Homo sapiens

<400> 118

atgcaaataa agctgcttta ggatccttac ctagattcta gaaaaatgggt gcatgaattt 60
gaacaaacaa actaattaaa aattaaaact gaaaaaaata gtttaaaaca acaactagag 120
gatatttttt catggcgcta aaagatacgg caaaaaaaat gactgacttg ttggaaagta 180
tccaacaaaa tttgcttaaa gcagaaaaag gaaataaagc cgcagcacia agagttccta 240
cagaatctat caaattagaa aagatcgca aggtatatcg taaagag 287

<210> 119

<211> 1002

<212> DNA

<213> Homo sapiens

<400> 119

catatgcac accatcacca tcacatgagt attcgaccta ctaatgggag tggaaatgga 60
taccgctcta ttaatccttc taacgataat caatacggtc ttgtgcaatc gacctctggg 120
cctaattacg gaggcatac ggtatcttct cgaggaggat ttcaagggat atgcgtacga 180
atagccgatt tattccgtaa ctgtttctct cgtaatagag gcactactac tacgccatct 240
cgaactgtta tcaactcaggc agatatttat catccgacta tttctggaca aggagctcaa 300
cctattgtct ctacaggaga taagaaatta gatagcgcaa ttattcaagc agatttgct 360
ggcgagaata aacagacttt ggctacacat attcaaagta agctagggtc tatggaggga 420
caatctcttc aagattataa agctgggtgc tatagtgcgc taagattgat gctgtttact 480
ccaggcgaaa ctactgtgag tagcgagcgg gaacgtcaag cgtgcgttac gggtcgggat 540
ctctgggaac aggtcgcagg agatcttgct accaatggga atacagatgg gcttatgtta 600
atggctaacc tatctgtggg agggaagcat gtgcctgcgg ggcatttaag agaatacatg 660
gatactgtaa aggtacggt tactgatgag aacgaggcta cagatcctac ggtagatgcc 720
attttgatt tagcagcaaa aatcgatgc acggaattct ctagtccctg ttcaggggcaa 780
gtcattctta attatatagg aaattatgga caagtcgttt tagaaaacga ggagatgaac 840
cttctgttt tagaagatca aaatgggcaa gatcctcaac gtgttcaaga taactcaaaa 900

86

gagttacaaa aactgttaga aaatgctcga aaaacagatc ctgagttata tttccaaaca 960
 ctaactgtca taacttcttc tgttttctta gactaaggat cc 1002

<210> 120
 <211> 1218
 <212> DNA
 <213> Homo sapiens

<400> 120
 atgcatcacc atcaccatca cgtgagtagc ataagcccta taggggggaa ttctgggcca 60
 gagggatttt ctagtgcacg tcgaggcgat gagattgatg atgtaccaga tagtgaagag 120
 ggagagctag aagagcgcgt ttccgatcat gcagagtcta tcattaccga gagctcggaa 180
 acgctgtttc gtactacttc ttcatcaggg gtcagtgaag atcttcagca acacgttagc 240
 ttggaggaat ctccacgaca acgaggtttc cttggacgga tccgtgatgc agtagcttct 300
 atttggaagc gtcgtgttgc acgaaggaat gaaaactatg atgtgaaaaa agcagaagag 360
 cagcaaggga ttgtgcaata tctgcaggat tcgaaaatgc ctgctttaac gcgtgcctat 420
 cgccatctcc gtgctttcaa ttctgcatgc ttaogtacga ttctgtagtt ttctgctacc 480
 atttttcgtg ctttaaggga tgcgtattat cgacattgta cacgttctgg gatcaacttt 540
 tgtggagctg ataaagactc tttagaagtt cttgtttgog tgggtttgct ttgctgtatg 600
 gctaccttac gctcttttga acatgtcggg gggaattacg aagatcgatt agtaaataat 660
 gatgctccgg tgacaggtgc ggggagaact cttgttgatg atgctgtaga cgatattgaa 720
 tcgattttta atacgagaac caactggcct caacatgtca tgatagggtt ttctcgtggg 780
 ctgcttcaat tatgtgcgac tccttataat ggcacttctc aagaatgttt caagtcgatt 840
 gttcgttttag aaaaagaaga cccttcttca gattattctc aagctttatt attagcaggg 900
 ataatagatc gcttgccgga gaaagcccct atggctgcaa agtatgtttt ggatgcattg 960
 cgtgttcgaa cttcggagct cataggagaa ctcatattc tcgatttgct tcctcctgta 1020
 tggaaggttg gccgcggagg cgtattccct cctgtgaatg agcagctcgt tgtgcaaatt 1080
 gttaatgcaa acgtagaacg attgcattcc actttcgtc atgagccaca agcttatttg 1140
 cgtatgatcg aagggttggt aaccaatttc tttttcttac ctagcgagga agatccttct 1200
 tcggttgga atatctaa 1218

<210> 121
 <211> 726
 <212> DNA
 <213> Homo sapiens

<400> 121
 catatgcac accatcacca tcacacaaag catggaaaac gcattcgtgg tatccaagag 60
 acttacgatt tagctaagtc gtattctttg ggtgaagcga tagatatattt aaaacagtgt 120
 cctactgtgc gtttcgatca aacggttgat gtgtctgtta aattagggat cgatccaaga 180
 aagagtgatc agcaaattcg tggttcgggt tctttacctc acggtacagg taaagttttg 240
 cgaatttttag tttttgctgc tggagataag gctgcagagg ctattgaagc aggagcggac 300
 tttgttggtg ggcacgactt ggtagaaaaa atcaaagggtg gatgggttga cttcgatggt 360
 gcggttgcca ctcccgatat gatgagagag gtccgaaaagc taggaaaagt tttaggtcca 420
 agaaacctta tgcctacgcc taaagccgga actgtaacaa cagatgtggt taaaactatt 480
 gcggaactgc gaaaaggtaa aattgaattt aaagctgatc gagctgggtg atgcaacgtc 540
 ggagttgcga agctttcttt cgatagtgcg caaatcaaag aaaatgttga agcgtttgtg 600
 gcagccttag ttaaagctaa gcccgcaact gctaaaggac aatatattag taatttctact 660
 atttcctcga ccatggggcc aggggttacc gtggatacta gggagttgat tgcgttataa 720
 gaatto 726

<210> 122
 <211> 330
 <212> PRT
 <213> Homo sapiens

<400> 122
 Met His His His His His Met Ser Ile Arg Pro Thr Asn Gly Ser

87

```

      5              10              15
Gly Asn Gly Tyr Pro Ser Ile Asn Pro Ser Asn Asp Asn Gln Tyr Gly
      20              25              30
Leu Val Gln Ser Thr Ser Gly Pro Asn Tyr Gly Gly His Thr Val Ser
      35              40              45
Ser Arg Gly Gly Phe Gln Gly Ile Cys Val Arg Ile Ala Asp Leu Phe
      50              55              60
Arg Asn Cys Phe Ser Arg Asn Arg Gly Thr Thr Thr Thr Pro Ser Arg
      65              70              75              80
Thr Val Ile Thr Gln Ala Asp Ile Tyr His Pro Thr Ile Ser Gly Gln
      85              90              95
Gly Ala Gln Pro Ile Val Ser Thr Gly Asp Lys Lys Leu Asp Ser Ala
      100              105              110
Ile Ile Gln Ala Asp Leu Arg Ala Gln Asn Lys Gln Thr Leu Ala Thr
      115              120              125
His Ile Gln Ser Lys Leu Gly Ser Met Glu Gly Gln Ser Pro Gln Asp
      130              135              140
Tyr Lys Ala Gly Ala Tyr Ser Ala Leu Arg Leu Met Leu Phe Thr Pro
      145              150              155              160
Gly Glu Thr Thr Val Ser Ser Glu Arg Glu Arg Gln Ala Cys Val Thr
      165              170              175
Gly Arg Asp Leu Trp Glu Gln Ala Ala Gly Asp Leu Ala Thr Asn Gly
      180              185              190
Asn Thr Asp Gly Leu Met Leu Met Ala Asn Leu Ser Val Gly Gly Lys
      195              200              205
His Val Pro Ala Gly His Leu Arg Glu Tyr Met Asp Thr Val Lys Gly
      210              215              220
Thr Phe Thr Asp Glu Asn Glu Ala Thr Asp Pro Thr Val Asp Ala Ile
      225              230              235              240
Leu Asp Leu Ala Ala Lys Ile Asp Ala Thr Glu Phe Ser Ser Pro Gly
      245              250              255
Ser Gly Gln Val Ile Leu Asn Tyr Ile Gly Asn Tyr Gly Gln Val Val
      260              265              270
Leu Glu Asn Glu Glu Met Asn Leu Leu Val Leu Glu Asp Gln Asn Gly
      275              280              285
Gln Asp Pro Gln Arg Val Gln Asp Asn Ser Lys Glu Leu Gln Lys Leu
      290              295              300
Leu Glu Asn Ala Arg Lys Thr Asp Pro Glu Leu Tyr Phe Gln Thr Leu
      305              310              315              320
Thr Val Ile Thr Ser Ser Val Phe Leu Asp
      325              330

```

<210> 123

<211> 405

<212> PRT

<213> Homo sapiens

<400> 123

```

Met His His His His His His Val Ser Ser Ile Ser Pro Ile Gly Gly
      5              10              15
Asn Ser Gly Pro Glu Gly Phe Ser Ser Ala Ser Arg Gly Asp Glu Ile
      20              25              30
Asp Asp Val Pro Asp Ser Glu Glu Gly Glu Leu Glu Glu Arg Val Ser
      35              40              45
Asp His Ala Glu Ser Ile Ile Thr Glu Ser Ser Glu Thr Leu Phe Arg
      50              55              60
Thr Thr Ser Ser Ser Gly Val Ser Glu Asp Leu Gln Gln His Val Ser
      65              70              75              80
Leu Glu Glu Ser Pro Arg Gln Arg Gly Phe Leu Gly Arg Ile Arg Asp
      85              90              95

```

88

Ala Val Ala Ser Ile Trp Lys Arg Arg Val Ala Arg Arg Asn Glu Asn
 100 105 110
 Tyr Asp Val Lys Lys Ala Glu Glu Gln Gln Gly Ile Val Gln Tyr Leu
 115 120 125
 Gln Asp Ser Lys Met Pro Ala Leu Thr Arg Ala Tyr Arg His Leu Arg
 130 135 140
 Ala Phe Asn Ser Ala Cys Leu Arg Thr Ile Arg Glu Phe Phe Ala Thr
 145 150 155 160
 Ile Phe Arg Ala Leu Arg Asp Ala Tyr Tyr Arg His Cys Thr Arg Ser
 165 170 175
 Gly Ile Asn Phe Cys Gly Ala Asp Lys Asp Ser Leu Glu Val Leu Val
 180 185 190
 Ala Val Gly Leu Leu Leu Arg Met Ala Thr Leu Arg Ser Phe Glu His
 195 200 205
 Val Gly Gly Asn Tyr Glu Asp Arg Leu Val Asn Asn Asp Ala Pro Val
 210 215 220
 Thr Gly Ala Gly Arg Thr Leu Val Asp Asp Ala Val Asp Asp Ile Glu
 225 230 235 240
 Ser Ile Leu Asn Thr Arg Thr Asn Trp Pro Gln His Val Met Ile Gly
 245 250 255
 Phe Ser Arg Gly Leu Val Gln Leu Cys Ala Thr Pro Tyr Asn Ala Thr
 260 265 270
 Ser Gln Glu Cys Phe Lys Ser Ile Val Arg Leu Glu Lys Glu Asp Pro
 275 280 285
 Ser Ser Asp Tyr Ser Gln Ala Leu Leu Leu Ala Gly Ile Ile Asp Arg
 290 295 300
 Leu Ala Glu Lys Ala Pro Met Ala Ala Lys Tyr Val Leu Asp Ala Leu
 305 310 315 320
 Arg Val Arg Thr Ser Glu Leu Ile Gly Glu Leu Ile Ile Leu Asp Leu
 325 330 335
 Leu Pro Pro Val Trp Lys Val Gly Arg Gly Gly Val Phe Pro Pro Val
 340 345 350
 Asn Glu Gln Leu Val Val Gln Ile Val Asn Ala Asn Val Glu Arg Leu
 355 360 365
 His Ser Thr Phe Ala His Glu Pro Gln Ala Tyr Leu Arg Met Ile Glu
 370 375 380
 Gly Leu Val Thr Asn Phe Phe Phe Leu Pro Ser Glu Glu Asp Pro Ser
 385 390 395 400
 Ser Val Gly Asn Ile
 405

<210> 124

<211> 238

<212> PRT

<213> Homo sapiens

<400> 124

Met His His His His His His Thr Lys His Gly Lys Arg Ile Arg Gly
 5 10 15
 Ile Gln Glu Thr Tyr Asp Leu Ala Lys Ser Tyr Ser Leu Gly Glu Ala
 20 25 30
 Ile Asp Ile Leu Lys Gln Cys Pro Thr Val Arg Phe Asp Gln Thr Val
 35 40 45
 Asp Val Ser Val Lys Leu Gly Ile Asp Pro Arg Lys Ser Asp Gln Gln
 50 55 60
 Ile Arg Gly Ser Val Ser Leu Pro His Gly Thr Gly Lys Val Leu Arg
 65 70 75 80
 Ile Leu Val Phe Ala Ala Gly Asp Lys Ala Ala Glu Ala Ile Glu Ala
 85 90 95
 Gly Ala Asp Phe Val Gly Ser Asp Asp Leu Val Glu Lys Ile Lys Gly

<400> 125							
ataacaatcc	ctcccaatca	tgttgaacg	tacaaggagg	agccatctat	gccaaaacct	60	
ctttgtctat	tggatcttcc	gatgtcgaa	cctctctat	tttctcgggg	aacagtgct	120	
cactgggaa	atctcaaaa	acaggcaca	tacggggagg	agcgatctac	tcccctactg	180	
ttacattgaa	ttgtcctgcg	acattctcta	acaatacagc	ctctatagct	acaccgaaga	240	
cttctttctga	agatggatcc	tcaggaaatt	ctattaaaga	taccattgga	ggagccattg	300	
cagggacagc	cattacccta	tctggagct	ctcgattttc	aggggaatag	gctgatttaag	360	
gagctgcaat	aggaactcta	gctaattgca	atacaccgac	tgcaactacg	ggatctcaaa	420	
atagcattac	agaaaaaatt	acttttagaa	acggttcttt	tatttttgaa	agaaaccaag	480	
ctaataaaag	tggagcgatt	tactctccta	gcgtttccat	taaagggaat	aatattacct	540	
ttcaatcaaaa	tacatccaact	catgatggaa	gcgcctatcta	ctttacaaaa	gatgctaaga	600	
ttagtctctt	aggattctgt	ctttttacag	gaaaataacgt	tacagctaca	caagctagtt	660	
ctgcaacatc	ttgacaaaaa	acaaaatactg	ccaactatgg	ggcagccatc	ttt	713	

<400>	126						
ccttctcctt	actcaggagt	tttaaaagaa	aacgcaccgt	ttttacgttt	cctcacacaa	60	
taaactaaga	agcatactca	ttctggattt	cattgcoctc	taaaattctt	agtcaaatcc	120	
gaagaagacc	gacactcgag	cgctcttctc	ctaaaaatct	tgttttttct	ctgcttcoga	180	
gttataacgc	ggctgtctca	taaccacac	taacatgatg	aaacctctac	gtttcggtta	240	
tttcttttgc	acaatctatt	ttactttggt	acaggcagcg	tttgctaaag	aaccgaattc	300	
ttgtcccgac	tgccagaata	attggaaaga	agtcacccac	acggatcaac	tcctgaaaa	360	
catcattcat	gctgatgatg	cttggttatc	ctctggttat	gtacaggctc	tcattgatat	420	
gcatttctta	gatagctgct	gcgaggtcat	cgttgaaaac	caaactgctt	acttattttc	480	
ttttcctaca	gatgatgtta	ggcgcaacgc	cattatcaac	ctaattaaag	accttccatt	540	
cattcaactc	gtagaaatct	gccaaagcatc	ctatcaaacc	tgtcatcatc	aaggccctca	600	
tggaagaact	ctctctccag	aacaacgttc	ttctgtgaca	aaggtctgtg	gaaaagaagc	660	
tatttggtta	ccacgaata	ccatcctatt	ctgcgctctt	gtagcagata	catatcaagc	720	
aactaatagt	gcaggtatcc	gttttaacga	cgaagtcgta	ggaaacctgt	ttggctctgc	780	

<210> 127
<211> 433

<212> DNA

<213> Chlamydia trachomatis

<400> 127

```

ctttaaagat tcgtcgtcct tttgttacta cgagagaagt tcgtgtgaaa tggcgttatg 60
ttcctgaagg tgtaggagat ttggctacca tagctccttc tatcagggtt ccacagttac 120
agaaatcgat gagaagcttt ttccctaaga aagatgatgc gtttcatcgg tctagtctgc 180
tattctactc tccaatgggt ccgcattttt ggcagagct tcgcaatcat tatgcaacga 240
gtggtttgaa aagcgggtac aatattggga gtaccgatgg gtttctccct gtcattgggc 300
ctgttatatg ggagtcggag ggtcttttcc gcgcttata ttcttcggtg actgatgggg 360
atggtaagag ccataaagta ggatttctaa gaattcctac atatagtggg caggacatgg 420
aagattttga tcc 433

```

<210> 128

<211> 803

<212> DNA

<213> Chlamydia trachomatis

<400> 128

```

atctattaat taatagcaag cttgaaacta aaaacctaatt ttattttaaag ctcaaaataa 60
aaaagagttt taaaatggga aattctgggt tttatttgta taacactgaa aactgcgtct 120
ttgctgataa tatcaaagt gggcaaatga cagagccgct caaggaccag caaataatcc 180
ttgggacaac atcaacacct gtccgagcca aaatgacagc ttctgatgga atatctttaa 240
cagtctccaa taattcatca accaatgctt ctattacaat tggtttggat gcggaaaaag 300
cttaaccagct tattctagaa aagttgggag atcaaaattct tgatggaatt gctgatacta 360
ttgttgatag tacagtccaa gatattttag acaaaatcaa aacagaccct tctctaggtt 420
tgttgaaagc ttttaacaac tttccaatca ctaataaaat tcaatgcaac gggttattca 480
ctcccagtaa cattgaaact ttattaggag gaactgaaat aggaaaattc acagtccacac 540
ccaaaagctc tgggagcatg ttcttagtct cagcagatat tattgcata agaatggaag 600
gcggcggttg totagctttg gtacgagaag gtgattctaa gccctgcgcg attagtattg 660
gatactcatc aggcattcct aatttatgta gtctaagaac cagtattact aatacaggat 720
tgactccgac aacgtattca ttacgtgtag gcggtttaga aagcgggtgtg gtatgggtta 780
atgccctttc taatctcgtg ccg 803

```

<210> 129

<211> 842

<212> DNA

<213> Chlamydia trachomatis

<400> 129

```

tgggaatgtc gaagaatacg attacgttct cgtatctata ggacgccgtt tgaatacaga 60
aaatattggc ttggataaag ctggtgttat ttgtgatgaa cgcggagtca tccctaccga 120
tgccacaatg cgcacaaacg tacctaacat ttatgctatt ggagatatca caggaaaatg 180
gcaacttgcc catgtagctt ctcatcaagg aatcattgca gcacggaata tagctggcca 240
taaagaggaa atcgattact ctgccgtccc ttctgtgatc tttaccttcc ctgaagtcgc 300
ttcagtaggc ctctcccaa cagcagctca acaacaaaaa atccccgtca aagtaacaaa 360
attccatttt cgagctattg gaaaagcggg cgaatggggc gaggccgatg gatttgcagc 420
cattatcagc catgagacta ctacgcagat cctaggagct tatgtgattg gccctcatgc 480
ctcatcactg atttccgaaa ttaccctagc agttcgtaat gaactgactc ttccttgtat 540
ttacgaaact atccacgcac atccaacctt agcagaagtt tgggctgaaa gtgcgttggt 600
agctgctgat accccattac atatgcccc tgcataaaaa tgaccgatcc agaattctct 660
actcctaata aatctatacc cgcagattc cctaagtggc tacgccagaa actcccttta 720
ggcggggtat ttgctcaaac tgataatact atcaaaaaa aagggcttcc tacagtctgt 780
gaggaagcct cttgtccgaa tcgcacccat tgttggtcta gacatacagc tacctatcta 840
gc 842

```

<210> 130

<211> 813

<212> DNA

<213> Chlamydia trachomatis

<400> 130

```

aaaatacttt gagctgcaca agctccccc tgttctagag aagaacatga tgcaaattcc 60
aatccaccct taatcttttc aaagataaga tcttctgtag aatataaagc cgctccagac 120
aaagaagcct tcacgtcagt taatgtgatt ccagccttac tactatcccc aacaaaagca 180
atacctaaaa aagattctcc gtcacgagga gaatcaaggt tgctgctcgt aaaactacaa 240
attaaccctt gggaagagac ttgatcctgt tgggtccacac cttggaaaac tacgggattg 300
gttactgaga acaaagtact ttgctctacc ttaccgggaa gagtatccgc atctttctct 360
tggaagaac ttggatctcc tacaattaac ctatactgtc ottcagcctg actatcttta 420
gacccaacga atagatctcg aatttgggtct aacaataaaa ccgcttgagg gcctacatat 480
accagctcat ttacagactg tctccagca tgaagatcta cgcaactagc taacccgcta 540
acagaggcaa ggatagctgc tactacagac aaagaaaact tagaacaggt gctttttata 600
tctttctcgg aactcatttc aaacctgcga aatagcactt ttttgacaaa ctagcgtacc 660
gaaacaatcg gtccaacaac gcgttctgcc tatgatttca caaagacaaa acgaccata 720
gacaagctcc agagacgaca ttagagcttt agaccgtgga atgtacaatg ctgactgctt 780
tttgagaaag attttttata aagaacaggc cct 813

```

<210> 131

<211> 1947

<212> DNA

<213> Chlamydia trachomatis

<400> 131

```

tcttttgcct atagagcaat ctcttatcat tgggtctgat ccaccagact atttcttcta 60
gatagagatt ctactacccc atccatggca ttcaacctct catcagtaaa cactttatta 120
gagttgttta tctgccatc atcgatgata tcttctgaag tctttaatac cttcttacat 180
aagatccatc tctccggaga acagtgtcct tctatggata aaattcctac gcagatattc 240
acgcattcca aaatagcagg aatacctaga tagatggcat ttacaaaoga agctgccgaa 300
actaggaata tcaaagcagt aatcaactaaa agtagtecta tcaccaactaa tcccacctta 360
aatgcagtyg aagatagaag attcgatata cgctctttca gtgttaatgg tgcagaacta 420
gtggaaatat cctgtgccga attggaagat ccagctcctt gaacaacggg tacagtgtc 480
atatttttaca ttctttttt gggtgtgagc agggagtcta cacaacact ttttttttc 540
aaaaaccctg ctgaatatg gaaatgaac tcttttattt tcatatagat 600
aacaaaaaaa agccgccag gaatccctgg acggcaccta cacatcgata aaatcaaaga 660
ttaatagatg tgtgtattct ctgtatcaga aactggaaca gtcaatgtat cggagaaaag 720
aatcgcttcc ccacgagcat ctccagctga tactgtttc aatgttacag aaaactctac 780
agtttcttta gaacctaatc taggtaacga atogaatact actgtattgc ctgtaatcgt 840
tcttttagtt ggtccagaga aggatacagg ttgcagttct ttagagaatt taagcattaa 900
agaaacattt gtatctctg cagaacctct gttggtgaca caaatcgggt aaacagtatt 960
ttctcctaca caaacagggt cacaagtatc tactacgcac atatgagtag cagcaactcc 1020
tttccagtaa gttgtcgtt ctgcgcaaga agtacaagta ccacagtcag agcagctctt 1080
cacaacaaca ttatttgtga attgtccagg agtttgtgct cttactagaa ctttatactg 1140
tagagactct ccaggattca gttctttcac agtccaaact actttattac aagaaatttg 1200
agctcctgca gcttcaagaa ctgtgactcc gggagaaaga gtgtcttcaa cgaogacatc 1260
tcgcaacaca agatctccag gatlggaaac ggagatcaca tattctacag gcttacaac 1320
ataagaccaa tctgtcctg caatacttac ttgtacgcaa ggctcattga tcacagttgt 1380
tacgcttgct gtatttttat gtctccaca gtaagaaaacc gttgtatata tggtagcacg 1440
accacgttta agcggacaaa actctacagt aattgttctg tgcctccag gttgcatact 1500
tccaagagta aacgtcagta cacgctgtcc agaagagtga gcgtaacct ctggaacagg 1560
attttcaaca acaacgttac gagctattgc tgttccttgg ttactacat taattttgta 1620
aactactggg caacgcaaac aagcattctc tgggccttct tgtttaacac agatagcagg 1680
ttgtccacat tttgtaaccg aacggatctc tggacaagcg catactgttg cagctgtaaa 1740
gcagcaacct tctttaagag gttttaccca tacagtaatt ttactctttt cgccttgtcc 1800
taagcggta attttccaaa ctagcttacc atcagcagta ggagtgtcg ctggatcact 1860
gcgtacgaac tctgcttcac atggtaattg ctgagtaatg ataacatcaa cacaatccct 1920
tttacctgta gcagtaattt caatagg 1947

```

<210> 132

<211> 1278

<212> DNA

<213> Chlamydia trachomatis

<400> 132

```

gataacaaaa aaaagccgcc caggaatccc tggacggcac ctacacatcg ataaaaatcaa 60
agattaatag atgtgtgtat tctctgtatc agaaactgga acagtcaatg tatcggaaga 120
aagaatcgct tccccacgag catctocagc tgatactgct ttcaatgtta cagaaaactc 180
tacagtttct ttagaacctt atctaggtaa cgaatcgaat actactgtat tgcttgaat 240
cgttccttta gttggtccag agaaggatac aggttgagcgt tctttagaga atttaagcat 300
taaagaaaca tttgtatctt ctgcagaacc tctgttggtg acacaaatac ggtaaacagt 360
atcttctcct acacaaacag ggtcacaagt atctactacg cacatatgag tagcagcaac 420
tcctttccag taagttgtcg cttctgcgca agaagtacaa gtaccacagt cagagcagct 480
cttcacaaca acattatttg tgaattgtcc aggagtttgt gctcttacta gaactttata 540
ctgtagagac tctccaggat tcagttcttt cacagtccaa actactttat tacaagaaat 600
ttgagctcct gcagcttcaa gaactgtgac tccgggagaa agagtgtctt caacgacgac 660
atctcgcaac acaagatctc caggattgga aacggagatc acatatctta caggcttaca 720
aacataagac caatctgctc ctgcaatact tacttgtacg caaggctcat tgatcacagt 780
tgttacgctt gctgtatttt tatgtcctcc acagtaagaa accgttgcta tattggtagc 840
acgaccacgt ttaagcggac aaaactctac agtaattgtt ctgtgctctc cagggttgcat 900
atctccaaga gtaaacgtca gtacacgctg tccagaagag tgagcgtaac catctggaac 960
aggattttca acaacaacgt tacgagctat tgctgttcct tggttcacta cattaatttt 1020
gtaactact gggcaacgca aacaagcatt ctctgggcct tcttgtttaa cacagatagc 1080
aggttgtcca cattttgtaa ccgaacggat ctctggacaa gcgcatactg ttgcagctgt 1140
aaagcagcaa ccttctttta gaggttttac ccatacagta attttactct ttctgccttg 1200
tcctaagcgg tcaattttcc aaactagctt accatcagca gtaggagttg tcgctggatc 1260
actgcgtacg aactctgc                                     1278

```

<210> 133

<211> 916

<212> DNA

<213> Chlamydia trachomatis

<400> 133

```

atggcgacaa ttttaacgatt accggacaaa accatacatt atcattttaca gattctcaag 60
ggccagttct tcaaaattat gccttcattt cagcaggaga gacacttact ctgaaagatt 120
tttcgagttt gatgttctcg aaaaatgttt cttgcggaga aaagggaatg atctcaggga 180
aaaccgtgag tatttccgga gcaggcgaag tgattttttg ggataactct gtgggggtatt 240
ctcctttgtc tatttggcca gcctogactc caactctctc agcaccagca ccagctcctg 300
ctgcttcaag ctctttatct ccaacagtta gtgatgctcg gaaagggctt attttttctg 360
tagagactag tttggagatc tcaggcgtca aaaaaggggt catgttogat aataatgccg 420
ggaatttttg aacagttttt cgaggtaata gtaataataa tgctggtagt gggggtagtg 480
ggtctgtctac aacaccaagt tttacagtta aaaactgtaa agggaaagt tctttcacag 540
ataacgtagc ctctgttgga gccggagtag tctacaaagg aactgtgctt ttcaaagaca 600
atgaaggagg catattcttc cgaggggaaca cagcatacga tgatttaggg attcttgctg 660
ctactagtcg ggatcagaat acggagacag gaggcgtgg aggagttatt tgctctccag 720
atgattctgt aaagtgtgaa ggcaataaag gttctattgt ttttgattac aactttgcaa 780
aaggcagagg cggaagcatc ctaacgaaag aattctctct tgtagcagat gattcggttg 840
tcttttagtaa caatacagca gaaaaggcg gtggagctat ttatgctcct acgtatcgat 900
ataagcacga atggag                                     916

```

<210> 134

<211> 751

<212> DNA

<213> Chlamydia trachomatis

<220>

<221> misc_feature

<222> (1)...(751)

<223> n = A,T,C or G

<400> 134

```

agcctctggc gaaggagagc cataaaaagt gcctaccagc ggagaaacaa taaaatctcc 60
ctgagcagggc aactcacttt ctttcttctc gatactctct ttaacaatag gattcccaag 120
gttttgatct gaggataagt tttgaaatcc agcaaacagt ctgttatcat aaaagactgg 180
ctcctgaata cttgggactg tatcccttcc taactctaac tccaaacctt cacgcttgat 240
aacaatgcgc ttcacgtgcc gaattcggca cgaggctctt tcttacgagg atctcgagtc 300
aagaagcctt gagccttcaa ttcttgcttc atgtcttctt tctcttgagc aacagctcta 360
gctaaaccca atcgagtagc aataacctga ccttgaaccc ctccctccact tactcggata 420
atcaaactga aactgttgac atcaccgagc attctgagcg gagctaagat ggttgctctt 480
tgaaacttcaa gagggaaata ttgctctaaa gtctttccat ttacgtcaat ttttccattc 540
ccagaacgaa gacgaacgca cacctgcttt cttctgcctg ttgcaacaga ctcttgatc 600
atattctttg tcacaaatta ccccaaatta cgcgtctaaa acaattgggt tgatagcttc 660
atactgtgcy taagaactac ctttcaaac tcttaaagat ttcatttgac gtcttccaag 720
ttttgtttta ggcaacattc nttaacagca t 751

```

<210> 135

<211> 410

<212> DNA

<213> Chlamydia trachomatis

<400> 135

```

ataatccaga ctcttctcca tctggagata gcgctggaga ctctgaagaa ctgactgaga 60
cagaagctgg ttctacaaca gaaactccta ctttaatagg aggaggtgct atctatggag 120
aaactgttaa gattgagaac ttctctggcc aaggaatatt ttctggaaac aaagctatcg 180
ataacaccac agaaggtcc tcttccaaat ctgacgtcct cggaggtgcy gtctatgcta 240
aaacattggt taatctcgat agcgggagct ctgacgaac tgtcaccttc tccgggaata 300
ctgtctcttc tcaatctaca acaggtcagg ttgctggagg agctatctac tctcctactg 360
taaccattgc tactcctgta gtattttcta aaaactctgc aacaaacaat 410

```

<210> 136

<211> 2719

<212> DNA

<213> Chlamydia trachomatis

<400> 136

```

ctcgtgcgga aaagctttct gctctaccaa agagattcgt tttttaaatt cttcattctc 60
tctaagagat ttagtttctt tcgcagaaca attgatagat actccgtacg tttgggggtg 120
ccggtgcatt cataaacagc ttctctgtaa tgggtgtagat tgttcggggg atattcaact 180
actttaccaa gtcacaggaa gaaatatccc tcgcaatgct agagatcaat acagagactg 240
ttctccagta aaagatttct cgtctctacc tataggagga cttatcttcc tcaagaaagc 300
aagcacggga caaatcaacc atgttatgat gaaaatctcg gagcatgaat tcattcatgc 360
tgccgaaaaa ataggggaaag tagaaaaagt aatcctagga aatagggtt tctttaaagg 420
gaatctattc tgctcattag gtgaaccgcc tatagaagct gtttttggcg ttcctaaaaa 480
tagaaaagcc ttcttttgaa agaaggcttt tctgaaacgc actccaatat atggacaagc 540
aatagcttat cgttttgaga attggaaact cttacgagct ttcttacgac cgtatttttt 600
acgctcttcc ttacgaggat ctcgagtcga gaagccttga gccttcaatt cttgcttcat 660
gtcttcttcc tcttgagaa cagctctagc taaacccaat cgagtagcaa taacctgacc 720
ttgaaccctt cctccactta ctcgataaat caaatcgaaa ctggtgacat caccgagcat 780
tctgagcgga gctaagatgg ttgctctttg aacttcaaga gggaaatatt gctctaaagt 840
ctttccattt acgtcaattt ttccattccc agaacgaaga cgaacgctag aaacagcctg 900
ctttctctcg cctggtgcaa cagactcttg tatcatattc tttgtcacia attaccccaa 960
attacgcgtc taaaacaatt ggtttgatag cttcatactg tgcgtaagaa ctacctttca 1020
aaactcttaa agatttcatt tgacgtcttc caagttttgt tttaggcaac attcctttta 1080
cagcatgctc gataacataa gcaggctttc gcgcaatcat gttttcaaaa ggaacttctc 1140
gcatcccaga aataaagcct gtgtaatagt gatcacctt ctgagttcct tttgocccag 1200
tcaaacgcac tttctcagca ttgatcacia tccacccatc tcccatcgct acgtgaggag 1260
taaaagtcac cttatgctta cctctcagga tottcgcaac ttctgaagat aatctcccta 1320
aggtcttccc ttcagcatta actacatacc aggtcttgtt tcgatcgtcc gaagccttag 1380

```

```

ctagggtcgt tttcgtatct tttctttttt ccataactta aatcacotta tcagagggaa 1440
tgattataat tttgatgatt atttttttcca aacaaaaagc agctgtattt gccttctaaa 1500
gaatttagaa aagaaaaaat ttcaaaaaga tctcttttct ttttgcttc aaaaacagcc 1560
ttacacttct atacttcttt cgaaaaaata ttttagggaa gttcttgaat catgatttac 1620
ataataaaaa aaatagtttag ctgccatcag ctaaatttaa aaaggtgcta ccagacgcta 1680
aaagctggtc cacgtaatta atatcataat cagaaagaag aaacttcgga ttatccaaca 1740
tgaactgatg aaaaggaatt gtagaatgca cccaccaat atggaactct tttaaagctc 1800
ttttcataat ggctatcgct tctctctgat tctttccttt tgtgattacc ttagcaatca 1860
tggaaatcata ataaggaggt atcgcataac cactgtagca agcccgctct actcgcacag 1920
caggacctgc aggagggaga taataatcta atctaccagg ggaaggagta aagttattaa 1980
ttggatcctc tgcattgatt cggcattgaa tcacgtgcc tttaaactct atattctttt 2040
gcttccaagg cagtttttct ccttagcga cactaatctg agcctttaac aaatcgatcc 2100
ctgtcacttc ttccgtaata gtatgttcca cttggatacg cgtattcatc tccatgaaat 2160
aaaaacgctt ctcccttatct aacagaaatt ctactgttcc aacagagaaa taccgggcac 2220
tcogagctaa atccactgct acttttccaa ctttagctcg catttctgga gttaaaatag 2280
gacttgaggt ctcttctatt aatttttgcc gacgcctttg tactgacaat ctcgctctcc 2340
aagatacacg taatttccgt gcttatctcc aattacttga acttctaaat gtcttgatt 2400
ttcaataaat ttttcaatat acacgtcagg attattaaat ccgcttctg cttcagcccg 2460
agcggcagta aaagccctat agaattcgct tttttctcta acaatccgta ttccctgctc 2520
accgcctcca gcaacagctt tgatgacgat gggaatccg atcttttctg caattctaat 2580
cccttccacc tcactcttca ctacaccttc agatccaggg attacaggc acttaattct 2640
tttagccaac tgcttagctg cgactttatc tcccatagtc gctatcgact cagcactagg 2700
accgataaat ctcggtgcg 2719

```

<210> 137

<211> 2354

<212> DNA

<213> Chlamydia trachomatis

<400> 137

```

gtgcaagatg ggacgagttt gaagtttaat actagcacat aacttccctt ctggaggttt 60
aggagagagc ccttttatta gggctctctt tttttgtgtg tgaggaaagc tagcgtctaa 120
ctaaatgtct ctaagtaagg atgttttttag gggaaatagc gattttcagt gttgagaagc 180
ttagttacaa gacaataaac aaggctaaga aaaacctttc tttagcctgt ttctcaacga 240
atcgccctata gaagactaat cttccagcgt tgcctatgg ctcagcttca actggccttt 300
ttcggttaatg ctaaggagtt taacagcaag cttgtctcct totttgacaa agccagagat 360
attgtctact ttttgtttag acaattcaga aatatgacag agcccttctt ttccctggag 420
gacttctacg aatactcaa atgttgcgat agatgtaaca cggccattat aaactttacc 480
gaacttcaact tctccagtta atccttcgat aagttcttta gctttgttaa tctgattcttg 540
gggtgcttgca gctatgttaa tgacgcgctc atcattgatg tcaacttgcg caccagaacg 600
ctcgataaatt tgacggattt gttttcctcc gggaccaatg accgttgcg tttttgaggt 660
attgatctgc atagtttcaa tgcgcggagc atatttagaa acagttccct taggggaggc 720
cagaacctgt gtcataagat taaggatatg actacgccct tgttttagctt gcgctagagc 780
ttgtctcata atcttatgag tgattccctc tatcttgata tccatttga aagctgtaat 840
accttttagct gttccggcta ctttaaagtc catatctcct agatgatctt ctataccgga 900
aatatcgac aagatgatgg cttgatctcg atctaagatt aagcccatag caatacctgc 960
cacgggagct ttgataggaa ctccagcatc catgagtga agacagcctc cacatacga 1020
tgccatggag gaagatccat tagactcagt aatattagat tctaggcgaa tgatataagg 1080
gaatcgcgat gtctcaggaa gaacatgact taaagctttc tcagctaatt tcccatgtcc 1140
aatttcacgt ctccctgggg aaccaattct gccaaactct cctacggaga aaggagggaa 1200
gaaatactgt agatagaagc gagcggtccc atctccattc agatcttcga atcgctgtgc 1260
catatttttc cctccaagcg tacatacggc catgctttgc gtctctccgc gaggtaataa 1320
gcaacttcog tgtgttcttg gaagaaaagg agtctctatg gaaatggggc gaatctctgt 1380
gggtggttcgt ccatctacac gaataccaag atcttgata agagctcgca tttgattgga 1440
ttttgctgtc ttaaatgcag ccttaacgtt caacaaagaa aaatcactgt tttcttcttg 1500
aaccaagtta gcaataacgg attcctctaa ttcttttcgag gcttgcctta gagcttcttt 1560
atctctaaaa gacaatgctt tttogaattt ttctctaata aaatctgaaa ctacattttg 1620
tacgtcttct ggcatatcaa gaacggcaga gaaattcttt tgtttgcga tagctttctg 1680
ccatgcttca atagcatcgc atatttttagc tatataggtt tgcccaaaaa caatagcttc 1740
tagaacttgc tcttctgtta aaaagtgcga atgtccttca atcattaaaa ctgcagaagc 1800

```



```

tggttcctgcc atgacgagat ccagcctgga ggcacttaac tcatctctgg ttgggttaat 1860
gacccacttt cctccgacga gcccaacgcg tacacccgca acgatacaat ttgaggaac 1920
ctctgagata gctaaagcgg cagaagctcc gcaaatagct agaggatcag gtaaagtfff 1980
cccgctcgtaa gaccaaactg aggacaagac ttgaatatct tgcattgagtc tattaggaaa 2040
cgacgggacgc aaagagcgat ccattagccg agaaacaaga atttctctct cggaaggccg 2100
tctttcacgt tttagaaatc ctccagaggt tcttcctgcg gaggaaaact tctcttgata 2160
gtctactctg aaaggcagaa aatcgacagc ctctgacaag gaggctgcac acgctgaaga 2220
aaaaacccaa gtctcggtca ttttgacgag aacagcccca ctggcctggc gagctatfff 2280
cctgtctctg aaaattaatg ttttattfff gtctaacgca acagaaaaag tctcaaaagc 2340
catggagttg tctt                                     2354

```

<210> 138

<211> 898

<212> DNA

<213> Chlamydia trachomatis

<400> 138

```

tcatcttctg tgatatttcc ggtatagaag atcatctagg agatatggac tttaaagtag 60
ccggaacagc taaaggtatt acagctttcc aaatggatat caagatagag ggaatcactc 120
ataagattat ggagcaagct ctacgcgaag cttaaacaagg gcgtagtcac atccttaatc 180
ttatgacaca gggtctggcc tcccctaagg gaactgttcc taaatatgct ccgcgcattg 240
aaactatgca gatcaatacc tcaaaaatcg caacgggtcat tgggtcccga ggaaaacaaa 300
tccgtcaaat tatcgagcgt tctggtgccc aagttgacat caatgatgac ggcgtcatta 360
acatagctgc aagcaccocaa gaatcgatta acaaagctaa agaacttacc gaaggattaa 420
ctggagaagt tgaagtcggt aaagtttata atggccgtgt tacatctatc gcaacatttg 480
gagttattct agaagtcctc ccaggaaaag aagggtcttg tcatatttct gaattgtcta 540
aacaataagt agacaatacc tctggttttg tcaaagaagg agacaagctt gctgttaaac 600
tcttagcat taacgaaaaa ggccagttga agctgagcca tagggcaacg ctggaagatt 660
agtcttctat aggcgattcg ttgagaaaca aggctaagaa aggtttttct tagccttgg 720
tattgtcttg taactaagct tctcaacact gaaaatcgct atttccccta aaaacatcct 780
tacttagaga catttagtta gacgctagct ttcctcacac acaaaaaaag agagccctaa 840
taaaagggct ctctcctaaa cctccagaag ggaagttatg tgctagtatt aaacttca 898

```

<210> 139

<211> 660

<212> PRT

<213> Chlamydia trachomatis

<400> 139

```

Met His His His His His His Met Glu Ser Gly Pro Glu Ser Val Ser
          5          10          15
Ser Asn Gln Ser Ser Met Asn Pro Ile Ile Asn Gly Gln Ile Ala Ser
          20          25          30
Asn Ser Glu Thr Lys Glu Ser Thr Lys Ala Ser Glu Ala Ser Pro Ser
          35          40          45
Ala Ser Ser Ser Val Ser Ser Trp Ser Phe Leu Ser Ser Ala Lys Asn
          50          55          60
Ala Leu Ile Ser Leu Arg Asp Ala Ile Leu Asn Lys Asn Ser Ser Pro
          65          70          75          80
Thr Asp Ser Leu Ser Gln Leu Glu Ala Ser Thr Ser Thr Ser Thr Val
          85          90          95
Thr Arg Val Ala Ala Lys Asp Tyr Asp Glu Ala Lys Ser Asn Phe Asp
          100          105          110
Thr Ala Lys Ser Gly Leu Glu Asn Ala Lys Thr Leu Ala Glu Tyr Glu
          115          120          125
Thr Lys Met Ala Asp Leu Met Ala Ala Leu Gln Asp Met Glu Arg Leu
          130          135          140
Ala Asn Ser Asp Pro Ser Asn Asn His Thr Glu Glu Val Asn Asn Ile
          145          150          155          160
Lys Lys Ala Leu Glu Ala Gln Lys Asp Thr Ile Asp Lys Leu Asn Lys

```

				165					170					175	
Leu	Val	Thr	Leu	Gln	Asn	Gln	Asn	Lys	Ser	Leu	Thr	Glu	Val	Leu	Lys
			180					185					190		
Thr	Thr	Asp	Ser	Ala	Asp	Gln	Ile	Pro	Ala	Ile	Asn	Ser	Gln	Leu	Glu
			195					200					205		
Ile	Asn	Lys	Asn	Ser	Ala	Asp	Gln	Ile	Ile	Lys	Asp	Leu	Glu	Arg	Gln
							215					220			
Asn	Ile	Ser	Tyr	Glu	Ala	Val	Leu	Thr	Asn	Ala	Gly	Glu	Val	Ile	Lys
225						230					235				240
Ala	Ser	Ser	Glu	Ala	Gly	Ile	Lys	Leu	Gly	Gln	Ala	Leu	Gln	Ser	Ile
				245						250				255	
Val	Asp	Ala	Gly	Asp	Gln	Ser	Gln	Ala	Ala	Val	Leu	Gln	Ala	Gln	Gln
			260					265					270		
Asn	Asn	Ser	Pro	Asp	Asn	Ile	Ala	Ala	Thr	Lys	Glu	Leu	Ile	Asp	Ala
			275					280					285		
Ala	Glu	Thr	Lys	Val	Asn	Glu	Leu	Lys	Gln	Glu	His	Thr	Gly	Leu	Thr
						295					300				
Asp	Ser	Pro	Leu	Val	Lys	Lys	Ala	Glu	Glu	Gln	Ile	Ser	Gln	Ala	Gln
305						310					315				320
Lys	Asp	Ile	Gln	Glu	Ile	Lys	Pro	Ser	Gly	Ser	Asp	Ile	Pro	Ile	Val
				325						330				335	
Gly	Pro	Ser	Gly	Ser	Ala	Ala	Ser	Ala	Gly	Ser	Ala	Ala	Gly	Ala	Leu
			340					345					350		
Lys	Ser	Ser	Asn	Asn	Ser	Gly	Arg	Ile	Ser	Leu	Leu	Leu	Asp	Asp	Val
			355					360					365		
Asp	Asn	Glu	Met	Ala	Ala	Ile	Ala	Leu	Gln	Gly	Phe	Arg	Ser	Met	Ile
						375					380				
Glu	Gln	Phe	Asn	Val	Asn	Asn	Pro	Ala	Thr	Ala	Lys	Glu	Leu	Gln	Ala
385						390					395				400
Met	Glu	Ala	Gln	Leu	Thr	Ala	Met	Ser	Asp	Gln	Leu	Val	Gly	Ala	Asp
				405						410				415	
Gly	Glu	Leu	Pro	Ala	Glu	Ile	Gln	Ala	Ile	Lys	Asp	Ala	Leu	Ala	Gln
			420					425					430		
Ala	Leu	Lys	Gln	Pro	Ser	Ala	Asp	Gly	Leu	Ala	Thr	Ala	Met	Gly	Gln
			435					440				445			
Val	Ala	Phe	Ala	Ala	Ala	Lys	Val	Gly	Gly	Gly	Ser	Ala	Gly	Thr	Ala
			450				455				460				
Gly	Thr	Val	Gln	Met	Asn	Val	Lys	Gln	Leu	Tyr	Lys	Thr	Ala	Phe	Ser
465						470				475				480	
Ser	Thr	Ser	Ser	Ser	Ser	Tyr	Ala	Ala	Ala	Leu	Ser	Asp	Gly	Tyr	Ser
				485						490				495	
Ala	Tyr	Lys	Thr	Leu	Asn	Ser	Leu	Tyr	Ser	Glu	Ser	Arg	Ser	Gly	Val
			500					505					510		
Gln	Ser	Ala	Ile	Ser	Gln	Thr	Ala	Asn	Pro	Ala	Leu	Ser	Arg	Ser	Val

Gly Tyr Leu Ser
660

<210> 140
<211> 598
<212> PRT
<213> Chlamydia trachomatis

<400> 140
Met His His His His His His Met Ser Ile Arg Gly Val Gly Gly Asn
5 10 15
Gly Asn Ser Arg Ile Pro Ser His Asn Gly Asp Gly Ser Asn Arg Arg
20 25 30
Ser Gln Asn Thr Lys Gly Asn Asn Lys Val Glu Asp Arg Val Cys Ser
35 40 45
Leu Tyr Ser Ser Arg Ser Asn Glu Asn Arg Glu Ser Pro Tyr Ala Val
50 55 60
Val Asp Val Ser Ser Met Ile Glu Ser Thr Pro Thr Ser Gly Glu Thr
65 70 75 80
Thr Arg Ala Ser Arg Gly Val Leu Ser Arg Phe Gln Arg Gly Leu Val
85 90 95
Arg Ile Ala Asp Lys Val Arg Arg Ala Val Gln Cys Ala Trp Ser Ser
100 105 110
Val Ser Thr Ser Arg Ser Ser Ala Thr Arg Ala Ala Glu Ser Gly Ser
115 120 125
Ser Ser Arg Thr Ala Arg Gly Ala Ser Ser Gly Tyr Arg Glu Tyr Ser
130 135 140
Pro Ser Ala Ala Arg Gly Leu Arg Leu Met Phe Thr Asp Phe Trp Arg
145 150 155 160
Thr Arg Val Leu Arg Gln Thr Ser Pro Met Ala Gly Val Phe Gly Asn
165 170 175
Leu Asp Val Asn Glu Ala Arg Leu Met Ala Ala Tyr Thr Ser Glu Cys
180 185 190
Ala Asp His Leu Glu Ala Lys Glu Leu Ala Gly Pro Asp Gly Val Ala
195 200 205
Ala Ala Arg Glu Ile Ala Lys Arg Trp Glu Lys Arg Val Arg Asp Leu
210 215 220
Gln Asp Lys Gly Ala Ala Arg Lys Leu Leu Asn Asp Pro Leu Gly Arg
225 230 235 240
Arg Thr Pro Asn Tyr Gln Ser Lys Asn Pro Gly Glu Tyr Thr Val Gly
245 250 255
Asn Ser Met Phe Tyr Asp Gly Pro Gln Val Ala Asn Leu Gln Asn Val
260 265 270
Asp Thr Gly Phe Trp Leu Asp Met Ser Asn Leu Ser Asp Val Val Leu
275 280 285
Ser Arg Glu Ile Gln Thr Gly Leu Arg Ala Arg Ala Thr Leu Glu Glu
290 295 300
Ser Met Pro Met Leu Glu Asn Leu Glu Glu Arg Phe Arg Arg Leu Gln
305 310 315 320
Glu Thr Cys Asp Ala Ala Arg Thr Glu Ile Glu Glu Ser Gly Trp Thr
325 330 335
Arg Glu Ser Ala Ser Arg Met Glu Gly Asp Glu Ala Gln Gly Pro Ser
340 345 350
Arg Val Gln Gln Ala Phe Gln Ser Phe Val Asn Glu Cys Asn Ser Ile
355 360 365
Glu Phe Ser Phe Gly Ser Phe Gly Glu His Val Arg Val Leu Cys Ala
370 375 380
Arg Val Ser Arg Gly Leu Ala Ala Ala Gly Glu Ala Ile Arg Arg Cys
385 390 395 400
Phe Ser Cys Cys Lys Gly Ser Thr His Arg Tyr Ala Pro Arg Asp Asp

[illegible]